

CLINICAL GUIDELINES

**MICHAEL T. CIBULKA, DPT • DOUGLAS M. WHITE, DPT • JUDITH WOHRLE, PT, PHD
 MARCIE HARRIS-HAYES, DPT • KEELAN ENSEKI, PT, MS • TIMOTHY L. FAGERSON, DPT
 JAMES SLOVER, MD, MS • JOSEPH J. GODGES, DPT**

Hip Pain and Mobility Deficits – Hip Osteoarthritis:*Clinical Practice Guidelines*

*Linked to the International Classification
 of Functioning, Disability, and Health
 from the Orthopaedic Section of the
 American Physical Therapy Association*

J Orthop Sports Phys Ther. 2009;39(_):A_-A_. doi:.....

RECOMMENDATIONS.....	Ax
INTRODUCTION.....	Ax
METHODS.....	Ax
CLINICAL GUIDELINES: <i>Impairment/Function-Based Diagnosis</i>	Axx
CLINICAL GUIDELINES: <i>Examinations</i>	Axx
CLINICAL GUIDELINES: <i>Interventions</i>	Axx
SUMMARY OF RECOMMENDATIONS.....	Axx
AUTHOR/REVIEWER AFFILIATIONS & CONTACTS.....	Axx
REFERENCES.....	Axx

Reviewers: Roy D. Altman, MD • Anthony Delitto, PT, PhD • John Dewitt, DPT • Amanda Ferland, DPT • Helene Fearon, PT • Joy MacDermid, PT, PhD • James W. Matheson, DPT • Kathleen Kline Mangione, PT, PhD • Philip McClure, PT, PhD • Marian A. Minor, PT, PhD • Paul Shekelle, MD, PhD • A. Russell Smith, Jr., PT, EdD • Leslie Torburn, DPT

For author, coordinator, and reviewer affiliations, see end of text. ©2009 Orthopaedic Section American Physical Therapy Association (APTA), Inc, and the *Journal of Orthopaedic & Sports Physical Therapy*. The Orthopaedic Section, APTA, Inc., and the *Journal of Orthopaedic & Sports Physical Therapy* consent to the photocopying of this guideline for educational purposes. Address correspondence to: Joseph J. Godges, DPT, ICF Practice Guidelines Coordinator, Orthopaedic Section, APTA Inc., 2920 East Avenue South, Suite 200; La Crosse, WI 54601. Email: icf@orthopt.org

Recommendations*

Pathoanatomical Features: Clinicians should assess for impairments in mobility of the hip joint and the strength of the surrounding muscles, especially the hip abductor muscles, when a patient presents with hip pain. (Recommendation based on weak evidence.)

Risk Factors: Clinicians should consider age, hip developmental disorders, and previous hip joint injury as risk factors for hip osteoarthritis. (Recommendation based on strong evidence.)

Diagnosis/Classification: Moderate lateral or anterior hip pain during weight bearing, in adults over the age of 50 years, with morning stiffness less than 1 hour, with limited hip internal rotation and hip flexion by more than 15° when comparing the painful to the non painful side are useful clinical findings to classify a patient with hip pain into the International Statistical Classification of Diseases and Related Health Problems (ICD) category of unilateral coxarthrosis and the associated International Classification of Functioning, Disability, and Health (ICF) impairment-based category of hip pain (b2816 Pain in joints) and mobility deficits (b7100 Mobility of a single joint). (Recommendation based on strong evidence.)

Differential Diagnosis: Clinicians should consider diagnostic classifications other than osteoarthritis of the hip when the patient's history, reported activity limitations, or impairments of body function and structure are not consistent with those presented in the diagnosis/classification section of this guideline - or - when the patient's symptoms are not diminishing with interventions aimed at normalization of the patient's impairments of body function. (Recommendation based on expert opinion.)

Examination – Outcome Measures: Clinicians should use validated functional outcome measures, such as the Western Ontario and McMaster Universities Osteoarthritis Index, Lower Extremity Functional Scale, and the Harris Hip Score before and after interventions intended to alleviate the impairments of body function and structure, activity limitations, and participation restrictions associated with hip osteoarthritis. (Recommendation based on strong evidence.)

Examination – Activity Limitation and Participation Restriction Measures: Clinicians should utilize easily reproducible physical performance measures, such as the 6 Minute Walk, Self-Paced Walk, Stair Measure, and Timed Up and Go to assess activity limitation and participation restrictions associated with their patient's hip pain and to assess the changes in the patient's level of function over the episode of care. (Recommendation based on strong evidence.)

Interventions - Patient Education: Clinicians should consider the use of patient education to teach activity modification, exercise, weight reduction when overweight, and methods of unloading the arthritic joints. (Recommendation based on moderate evidence.)

Interventions – Functional, Gait, and Balance Training: Functional, gait, and balance training, including the use of assistive devices such as canes, crutches, and walkers, can be used in patients with hip osteoarthritis to improve function associated with weight bearing activities. (Recommendation based on weak evidence.)

Interventions – Manual Therapy: Clinicians should consider the use of manual therapy procedures to provide short-term pain relief and improve hip mobility and function in patients with mild hip osteoarthritis. (Recommendation based on moderate evidence.)

Interventions – Flexibility, Strengthening, and Endurance Exercises: Clinicians should consider the use of flexibility, strengthening, and endurance exercises in patients with hip osteoarthritis. (Recommendation based on moderate evidence.)

*These recommendations and clinical practice guidelines are based on the scientific literature published prior to August 2008.

Introduction

AIM OF THE GUIDELINE

The Orthopaedic Section of the American Physical Therapy Association (APTA) has an ongoing effort to create evidence-based practice guidelines for orthopaedic physical therapy management of patients with musculoskeletal impairments described in the World Health Organization's International Classification of Functioning, Disability, and Health (ICF).⁹²

The purposes of these clinical guidelines are to:

- Describe evidence-based physical therapy practice including diagnosis, prognosis, intervention, and assessment of outcome for musculoskeletal disorders commonly managed by orthopaedic physical therapists
- Classify and define common musculoskeletal conditions using the World Health Organization's terminology related to impairments of body function and body structure, activity limitations, and participation restrictions
- Identify interventions supported by current best evidence to address impairments of body function and structure, activity limitations, and participation restrictions associated with common musculoskeletal conditions
- Identify appropriate outcome measures to assess changes resulting from physical therapy interventions in body function and structure as well as in activity and participation of the individual
- Provide a description to policy makers, using internationally accepted terminology, of the practice of orthopaedic physical therapists
- Provide information for payors and claims reviewers regarding the practice of orthopaedic physical therapy for common musculoskeletal conditions
- Create a reference publication for orthopaedic physical therapy clinicians, academic instructors, clinical instructors, students, interns, residents, and fellows regarding the best current practice of orthopaedic physical therapy

STATEMENT OF INTENT

This guideline is not intended to be construed or to serve as a standard of medical care. Standards of care are determined on the basis of all clinical data available for an individual patient and are subject to change as scientific knowledge and technology advance and patterns of care evolve. These parameters of practice should be considered guidelines only. Adherence to them will not ensure a successful outcome in every patient, nor should they be construed as including all proper methods of care or excluding other acceptable methods of care aimed at the same results. The ultimate judgment regarding a particular clinical procedure or treatment plan must be made in light of the clinical data presented by the patient, the diagnostic and treatment options available, and the patient's values, expectations, and preferences. However, we suggest the rationale for significant departures from accepted guidelines be documented in the patient's medical records at the time the relevant clinical decision is made.

Methods

The Orthopaedic Section, APTA appointed content experts as developers and authors of clinical practice guidelines for musculoskeletal conditions of the hip which are commonly treated by physical therapists. These content experts were given the task to identify impairments of body function and structure, activity limitations, and participation restrictions, described using ICF terminology, which could 1) categorize patients into mutually exclusive impairment patterns upon which to base intervention strategies, and 2) serve as measures of changes in function over the course of an episode of care. The second task given to the content experts was to describe interventions and supporting evidence for specific subsets of patients based upon the previously chosen patient categories. It was also acknowledged by the Orthopaedic Section, APTA content experts that a systematic search and review of the evidence solely related to diagnostic categories based on International Statistical Classification of Diseases and Health Related Problems (ICD)⁹¹ terminology would not be useful for these ICF-based clinical practice guidelines as most of the evidence associated with changes in levels of impairment or function in homogeneous populations is not readily searchable using the ICD terminology. The authors of this guideline (MTC, DMW, JW) independently performed a systematic search of the MEDLINE, CINAHL, and the Cochrane Database of Systematic Reviews (1967 through August 2008) for any relevant articles related to classification, examination, and intervention for musculoskeletal conditions of the hip region commonly treated by physical therapists. As relevant articles were identified, their reference lists were hand-searched in an attempt to identify additional articles that might contribute to the outcome of this guideline. Articles from the searches were compiled by three of the authors (MTC, DMW, JW) and this compilation was reviewed for accuracy and completeness by three other authors (MHH, KE, TLF). Articles with the highest levels of evidence that were most relevant to classification, examination, and intervention for patients with hip pain, mobility deficits, and osteoarthritis were included in this guideline.

This guideline was issued in 2009 based upon publications in the scientific literature prior to September 2008. This guideline will be considered for review in 2013, or sooner if new evidence becomes available. Any updates to the guideline in the interim period will be noted on the Orthopaedic Section of the APTA website: www.orthopt.org

Levels of Evidence

Individual clinical research articles were graded according to criteria described by the Center for Evidence-Based Medicine, Oxford, United Kingdom (Table 1).

I	Evidence obtained from high-quality randomized controlled trials, prospective studies, or diagnostic studies
II	Evidence obtained from lesser quality randomized controlled trials, prospective studies, or diagnostic studies (eg, improper randomization, no blinding, < 80% follow-up)
III	Case controlled studies or retrospective studies
IV	Case series
V	Expert opinion

Grades of Evidence

The overall strength of the evidence supporting recommendations made in this guideline were graded according to guidelines described by Guyatt et al,⁷⁵ as modified by MacDermid and adopted by the coordinator and reviewers of this project. In this modified system, the typical A, B, C, and D grades of evidence have been modified to include the role of consensus expert opinion and basic science research to demonstrate biological or biomechanical plausibility (Table 2).

Grades of Recommendation		Strength of Evidence
A	Strong evidence	A preponderance of level I and/or level II studies support the recommendation. This must include at least 1 level I study
B	Moderate evidence	A single high-quality randomized controlled trial or a preponderance of level II studies support the recommendation
C	Weak evidence	A single level II study or a preponderance of level III and IV studies including statements of consensus by content experts support the recommendation
D	Conflicting evidence	Higher-quality studies conducted on this topic disagree with respect to their conclusions. The recommendation is based on these conflicting studies
E	Theoretical/foundational evidence	A preponderance of evidence from animal or cadaver studies, from conceptual models/principles, or from basic sciences/bench research support this conclusion
F	Expert opinion	Best practice based on the clinical experience of the guidelines development team

Review Process

The Orthopaedic Section, APTA also selected consultants from the following areas to serve as reviewers of the early drafts of this clinical practice guideline:

- Arthritis Foundation
- Claims review
- Coding
- Epidemiology
- Rheumatology
- Section on Geriatrics of the APTA
- Medical practice guidelines
- Orthopaedic physical therapy residency education
- Physical therapy academic education
- Sports physical therapy residency education

Comments from these reviewers were utilized by the authors to edit this clinical practice guideline prior to submitting it for publication to the *Journal of Orthopaedic & Sports Physical Therapy*.

In addition, several physical therapists practicing in orthopaedic and sports physical therapy settings were sent initial drafts of this clinical practice guideline along with feedback forms to determine its usefulness, validity, and impact. All returned feedback forms from these practicing clinicians described this clinical practice guideline as:

- “Moderately useful” or “extremely useful”
- An “accurate representation of the peer-reviewed literature”
- A guideline that will have a “substantial positive impact on orthopaedic physical therapy patient care”

CLASSIFICATION

The primary ICD-10 code and condition associated with hip pain and mobility deficits is M16.1 Primary coxarthrosis, unilateral. In the ICD, the term osteoarthritis (OA) is used as a synonym for arthrosis or osteoarthritis. Other, secondary codes associated with hip OA are M16.0 Primary coxarthrosis, bilateral; M16.2 Coxarthrosis resulting from dysplasia, bilateral; M16.3 Dysplastic coxarthrosis, unilateral; M16.4 Post-traumatic coxarthrosis, bilateral; M16.5 Post-traumatic coxarthrosis, unilateral; M16.7 Secondary coxarthrosis, not otherwise specified. The corresponding ICD-9 CM codes and conditions, which are used in the USA, are 715.15 Osteoarthritis of the pelvic region and thigh, localized, primary; 715.25 Osteoarthritis of the pelvic region and thigh, localized, secondary; 715.85 Osteoarthritis of the pelvic region and thigh involving more than 1 site but not specified as generalized.

The primary ICF body function codes associated with the above noted primary ICD-10 condition are the sensory functions related to pain and the movement-related functions related to joint mobility. These body function codes are **b2816 Pain in joints** and **b7100 Mobility of a single joint**.

The primary ICF body structure codes associated with hip pain and mobility deficits are **s75001 Hip joint**, **s7402 Muscles of pelvic region**, and **s7403 Ligaments and fascia of pelvic region**.

The primary ICF activities and participation codes associated with hip pain and mobility deficits are **d4154 Maintaining a standing position**, **d4500 Walking short distances**, and **d4501 Walking long distances**.

The ICD-10 and primary and secondary ICF codes associated with hip pain and mobility deficits are provided in Table 3 (below).

ICD-10 and ICF Codes Associated with Hip Pain and Mobility Deficits

International Statistical Classification of Diseases and Related Health Problems

Primary ICD-10	M16.1	Primary coxarthrosis, unilateral
Secondary ICD-10	M16.0	Primary coxarthrosis, bilateral
	M16.2	Coxarthrosis resulting from dysplasia, bilateral
	M16.3	Dysplastic coxarthrosis, unilateral
	M16.4	Post-traumatic coxarthrosis, bilateral
	M16.5	Post-traumatic coxarthrosis, unilateral
	M16.7	Secondary coxarthrosis, not otherwise specified

International Classification of Functioning, Disability and Health

PRIMARY ICF CODES

Body functions	b28016	Pain in joints
	b7100	Mobility of a single joint
Body structure	s75001	Hip joint
	s7402	Muscles of pelvic region
	s7403	Ligaments and fascia of pelvic region
Activities and participation	d4154	Maintaining a standing position
	d4500	Walking short distances
	d4501	Walking long distances

SECONDARY ICF CODES

Body functions	b7201	Mobility of the pelvis
	b7300	Power of isolated muscles and muscle groups
	b7401	Endurance of muscle groups
	b770	Gait pattern functions
	b7800	Sensation of muscle stiffness
Body structure	s7401	Joints of pelvic region
Activities and participation	d4101	Squatting
	d4103	Sitting
	d4106	Shifting the body's center of gravity
	d4350	Pushing with lower extremities
	d4351	Kicking
	d4502	Walking on different slopes
	d4503	Walking around obstacles
	d4551	Climbing
	d4552	Running
	d4553	Jumping
	d4600	Moving around within the home
	d4601	Moving around within buildings other than home
d4602	Moving around outside the home or other building	

CLINICAL GUIDELINES

Impairment/Function-based Diagnosis**PREVALENCE**

Hip pain associated with OA is the most common cause of hip pain in older adults.^{41, 177} Prevalence studies have shown the rates for adult hip OA range from 0.4% to 27%.^{7, 41, 74, 89, 128, 213}

PATHOANATOMICAL FEATURES

The proximal femur articulates with the acetabulum to form the hip joint. The femoral head is two thirds of a sphere covered with hyaline cartilage and enclosed in a fibrous capsule.^{50, 170} The femoral head is connected to the femoral shaft via the femoral neck. In the frontal plane the femoral neck lies at an angle to the shaft of the femur. This “angle of inclination” is normally 120-125° in the adult population.⁵⁰ In the transverse plane the proximal femur is oriented anterior to the distal femoral condyles as a result of a medial torsion of the femur with a normal range between 14-18° of anteversion.²⁸ The hip joint is a “ball and socket” synovial joint with articular cartilage and a fully developed joint capsule allowing movement in all 3 body planes.¹⁷⁰ The joint capsule attaches around the acetabular rim proximally and distally at the inter-trochanteric line. Three strong ligaments reinforce the joint capsule, the ilio-femoral and pubo-femoral ligaments anteriorly and ischio-femoral ligament posteriorly.⁵⁰

In OA of the hip the entire joint structure and function is affected, with joint capsular changes (shortening and lengthening) creating limitation in hip joint range of motion (ROM) along with subsequent articular cartilage degeneration.^{122, 150} Later in the disease process osteophytes or spurs may develop from excessive tensile force on the hip joint capsule or from abnormal pressure on the articular cartilage.^{6, 122} Other changes also develop including sclerosis of the subchondral bone from increased focal pressure, and sometimes the formation of cysts.¹⁰⁰ Muscle weakness often develops around a joint with OA,¹⁷³ specifically the abductor muscles in the hip.¹⁵⁸ Most significantly hip abductor muscle progressively weakens in the later stages of hip osteoarthritis, which may create a Trendelenberg gait pattern over time.¹⁰ Clinicians must not overlook the function of the hip muscles at different hip positions, for example the gluteus medius is an abductor when at 0° of flexion (standing) but an external rotator at 90° of flexion (sitting).¹³¹

II

C

Clinicians should assess for impairments in the mobility of the hip joint and the strength of the surrounding muscles, especially the hip abductor muscles, when a patient presents with hip pain.

RISK FACTORS

Age: The most common predisposing factor for hip OA is age. The condition affecting primarily middle-aged and elderly people, most often in adults over 60 years of age.^{2, 156, 180} Tepper and Hochberg^{2, 156, 180} found that age was significantly associated with hip OA (adjusted odds ratio = 1.30 (95% CI 0.60-2.81) for ages 60-64, 2.38 (95% CI 0.83-3.44) for ages 65-69, and 2.38 (95% CI 1.15-4.92) for ages 70-74.

I

Developmental Disorders: Many studies have demonstrated a link between developmental disorders, such as Legg-Calve-Perthes disease, congenital hip dislocation, or slipped capital femoral epiphysis, and premature OA of the hip.^{1, 61, 62, 93, 94, 141, 216} The majority of the evidence also shows that dysplasia of the femur and the acetabulum is associated with hip OA.^{49, 52, 94, 96, 164, 181, 188-191}

Dysplasia is defined as any change in orientation of the acetabulum or the proximal femur, which creates a change in how the femur and the acetabulum articulate with each other.⁹⁵ Types of dysplasia include coxa vara, coxa valga, femoral anteversion, femoral retroversion, acetabular anteversion, acetabular retroversion, coxa plana, and coxa profundus.^{20, 94, 95} However, a few authors show limited association between hip dysplasia and OA of the hip.^{20, 90, 110, 210}

I

Race: Studies of non Caucasian populations including Asian, African, and East Indian populations indicate a very low prevalence of primary hip OA when compared to that of Caucasians of European ancestry.^{81, 213} Thus, race likely plays some sort of role in the development of hip OA. This role, however, is unknown at this time.

III

Gender: Few studies have been performed that examine gender and hip OA.^{39, 99, 147, 156} Tepper and Hochberg found that males have a slightly greater prevalence of hip OA compared to females (3.2% to 3.0%).¹⁸⁰ Although little if any difference in incidence exists between genders, men and women appear to have different patterns of hip OA. Women have a more superiomedial femoral migration while men have a more superiolateral migration.^{44, 113} There is strong evidence that superiolateral migration is an important prognostic factor in the progression of hip OA.¹⁵

I

Genetics: Siblings show a high association of hip OA, suggesting a possible genetic role.¹¹¹ While the nature of the genetic influence is still speculative, it has been postulated that hip OA involves either a structural defect (ie, collagen) or alterations in cartilage or bone metabolism.^{108, 123, 124} Genetics is often linked to hip OA because of the low prevalence of hip OA in Asian and African populations in their native countries and the familial association of hip OA in Caucasians.⁸¹ Some studies have shown that genetic factors may play a role in the development of hip OA,^{108, 126} or in reducing the risk of hip OA in women.¹¹⁵ Although there has been much interest and speculation in looking for a genetic link to hip OA, currently there is insufficient evidence to explain how genetics is related to the development of hip OA.

II

Occupation: Numerous researchers in Europe and the United States have found a higher prevalence of hip OA in male workers whose occupation involved lifting very heavy loads over a prolonged period of time.^{127, 204, 206, 207} Farming, in particular, has been identified as a high-risk occupation for the development of hip OA. However, a specific aspect of farming which leads to the development of hip OA has not been identified. Suspected risk factors have been suggested, including regular heavy lifting, tractor driving (vibration), and walking on uneven ground.^{36, 84, 127, 182-185, 211} Vibration was specifically studied and reported to not be associated with the development of hip OA.⁹⁸ In summary, there is weak evidence linking the development of hip OA to some occupations.¹¹⁶

III

Sports Exposure: Epidemiological studies have demonstrated participation in certain competitive sports to increase the risk for OA.^{25, 117} Running has low risk for OA,¹⁰⁹ but high-intensity, direct impact activities, such as American football and hockey, appear to increase the risk of hip OA.^{24, 25}

II

Previous injury: Proximal hip fracture results in changes to the articular surfaces of the hip joint that creates abnormal joint load bearing and has been shown to be related to the development of hip OA.⁶⁰ A history of a previous hip injury is also associated with hip OA.^{35, 180} Cooper et al³⁵ reported an odds ratio for hip OA when having a previous hip injury as 4.3 (95% CI: 2.2-8.4). In addition, patients with OA of 1 hip are at increased risk of developing OA in the opposite hip.²⁰⁹

I

Body mass index: A few studies have shown body mass index (BMI) to be related to hip OA.^{35, 118, 203} Other studies, however, have shown little to no association between hip OA and BMI.^{59, 73, 94, 97, 119} The most current evidence shows BMI is not related to hip OA.¹⁶³ What seems apparent is obesity is probably associated with the progression of hip OA rather than onset,¹²⁹ and the therapeutic value of weight loss is important.^{52, 205}

III

Leg length disparity: Several studies have suggested a difference in leg lengths may be an important etiological factor in hip OA.^{63-65, 68, 142} A few studies have demonstrated the biomechanical and clinical problem of leg length disparity and its relationship to hip OA.⁶³⁻⁶⁵ Nahoda¹⁴³ reported on the importance of correcting leg length disparity in the prevention of hip OA. Golightly⁶⁵ noted an association between radiographic hip OA and leg length disparity. The few papers on leg length disparity suggest a relationship with hip; however more research is needed before leg length disparity can be considered an important risk factor for hip. It should be pointed out that leg length disparity can also occur as a result of OA of the hip, particularly when there is superior migration of the hip in the acetabulum.

II

A

Clinicians should consider age, hip developmental disorders, and previous hip joint injury as risk factors for hip osteoarthritis.

NATURAL HISTORY

The natural history of individual hip OA is imperfectly understood. Many different factors contribute to this. The clinical manifestations that develop in patients with hip OA include changes in the shape, density, length, and function of the bones, cartilage, and fibrous tissue surrounding the hip joint itself as well as the muscles surrounding the hip joint. The changes that occur around the arthritic hip joint include a decrease in the joint space between the femur and acetabulum (more common superior and lateral than medial), shortening of the fibrous joint capsule, flattening of the femoral head, the appearance of osteophytic growth around the margins of the femoral head and acetabulum (in some individuals bony over growth does not occur), a superior-lateral or medial migration of the femoral head, and the development of subchondral sclerosis or cysts in the femoral head and acetabulum.^{4, 6, 42, 72, 100, 169, 164} Changes that occur outside of the hip joint include a decreased amount of hip joint ROM (usually mostly affecting internal rotation and then flexion) and muscle weakness (particularly the abductor muscles); which eventually may result in difficulty with ambulation.^{118, 120, 150, 177} The progression of these changes are usually slow but in some cases quite rapid.²¹ Currently, there is no reliable, generally accepted classification of the stages or severity of hip OA and the rate of progression varies from patient to patient, even when the demographics of the patients are similar.¹⁷⁷

DIAGNOSIS/CLASSIFICATION

The diagnosis of hip OA can be made with a reasonable level of certainty on the basis of the history and physical examination.^{2, 16} Joint space narrowing along with other radiographic features including osteophytes and subchondral sclerosis on plain film radiographs is considered the definitive diagnosis.^{6, 19, 37, 42, 45} The following clinical criteria are typically present in individuals who have radiographic findings consistent with hip OA^{2, 16, 18}

- Reports of moderate pain in the lateral or anterior hip with weight bearing. This pain may progress to the anterior thigh or knee region.
- Adults, greater than 50 years of age.
- Limited passive hip joint ROM in at least 2 of its 6 directions (flexion, extension, abduction, adduction, internal and external rotation).
- Morning stiffness, which improves in less than 1 hour.

I

Clinical criteria for the classification of patients with hip pain associated with OA were developed through a multicenter study by the American College of Rheumatology.² One hundred fourteen patients with a mean age of 64 and 87 controls with a mean age of 57 years were included in the study. Patients were classified as having hip OA if the patient 1) reports experiencing hip pain, and 2) presents with one of the following clusters of clinical findings:

- Hip internal rotation $< 15^\circ$, along with
 - Hip flexion $\leq 115^\circ$
 - Age > 50 years
- Or,
- Hip internal rotation $\geq 15^\circ$, along with
 - Pain with hip internal rotation
 - Duration of morning stiffness of the hip is ≤ 60 minutes
 - Age > 50 years

I

When patients were classified using these clinical criteria compared to a radiographic reference standard of joint space narrowing and osteophytes, the following diagnostic accuracy statistics were reported: sensitivity 86 %; specificity 75 %; positive likelihood ratio (LR+) 3.44; negative likelihood ratio (LR-) 0.19.²

Hip OA is classified as primary, in the absence of any obvious underlying joint abnormality, or secondary if degeneration occurs as a result of a pre-existing abnormal joint problem.⁸¹ Some suggest that all hip OA is secondary to some pre-existing problem (eg, dysplasia).¹²¹ The clinical and/or radiological criteria presented above are usually sufficient to diagnose a patient with OA of the hip and the associated ICF impairment-based category of hip pain (b2816 Pain in joints) and mobility deficits (b7100 Mobility of a single joint).

A recent preliminary study of patients with hip symptoms identified 5 possible clinical predictors for diagnosis: pain aggravated with squatting, lateral or anterior hip pain with the scour test, active hip flexion causing lateral hip pain,

II

pain with active extension, and passive range of internal rotation less than 25° .¹⁷⁸ The LR+ of having hip OA when all 5 predictors were present was 7.3 while the LR- was .87.¹⁷⁸ One limitation of this study was the small sample of patients (21 of 79) who had hip OA on radiographs. This could have resulted in spurious findings. Future studies are needed to validate these results prior to clinical use.

Birrell et al¹⁸ also used a standard clinical and radiographic examination to assess the predictability of hip OA from hip ROM. In their study, 195 patients with recent onset of hip pain with radiographic evidence of OA had restricted movement at the hip compared with those without radiographic evidence of hip OA. Restriction in internal rotation was the most predictive and flexion the least predictive of radiographic OA. When comparing sides, a ROM difference of more than 15° between the painful and non-painful side was considered a limitation of joint mobility. Restriction in hip ROM was predictive of the presence of OA. The diagnostic accuracy for restriction in a single plane of hip motion for patients with severe hip OA was: sensitivity 100%, specificity 54%, LR+ 2.17, LR- 0.01. The diagnostic accuracy for restriction in a single plane of hip motion for patients with moderate hip OA was: sensitivity 86%, specificity 42%, LR+ 1.48, LR- 0.33. The diagnostic accuracy for restriction in all 3 planes of hip motion for patients with moderate hip OA was: sensitivity 33%, specificity 98%, LR+ 16.5, LR- 0.67.¹⁸

A

Moderate lateral or anterior hip pain during weight bearing, in adults over the age of 50 years, with morning stiffness less than 1 hour, with limited hip internal rotation and hip flexion by more than 15° when comparing the painful to the non painful side are useful clinical findings to classify a patient with hip pain into the International Statistical Classification of Diseases and Related Health Problems (ICD) category of unilateral coxarthrosis and the associated International Classification of Functioning, Disability, and Health (ICF) impairment-based category of hip pain (b2816 Pain in joints) and mobility deficits (b7100 Mobility of a single joint).

I

DIFFERENTIAL DIAGNOSIS

The following differential diagnoses should be considered in an individual with hip OA:

- Bursitis or tendinitis
- Chondral damage or loose bodies
- Femoral neck or pubic ramus stress fracture
- Labral tear
- Muscle strain
- Neoplasm
- Osteonecrosis of the femoral head
- Paget's disease
- Piriformis syndrome
- Psoriatic arthritis
- Rheumatoid arthritis
- Sacroiliac joint dysfunction
- Septic hip arthritis
- Referred pain as a result of an L2-3 radiculopathy

The following physical examination measures may be helpful in the differential diagnostic process when differentiating hip pain from other sources of pain:

- The Scour test for labral tears¹⁷⁸
- FABER (Patrick's) test¹³⁹
- Fitzgerald's test for labral tears⁵⁴
- Flexion-adduction-internal rotation test for labral tears¹¹⁴
- Sacroiliac joint provocation test for sacroiliac joint pain¹¹²
- Femoral nerve stretch test for L2-3 radiculopathy¹⁸⁷

F

Clinicians should consider diagnostic classifications other than osteoarthritis of the hip when the patient's history, reported activity limitations, or impairments of body function and structure are not consistent with those presented in the diagnosis/classification section of this guideline - or - when the patient's symptoms are not diminishing with interventions aimed at normalization of the patient's impairments of body function.

V

IMAGING STUDIES

Imaging studies, specifically plain film radiographs, are confirmatory for moderate to severe hip joint OA, however, radiographs are less useful in demonstrating early osteoarthritic joint changes.^{53, 100} Joint space narrowing detected on radiographs may be a relatively late stage phenomenon of OA.²³ Joint space narrowing has been advocated as the best indicator and best predictor of arthritic change in patients with hip OA, with joint space narrowing occurring more superiolateral than superiomedial.^{37, 42, 45} The normal hip joint space is 3-5mm. A reduction of greater than or equal to 0.5 mm represents a clinically relevant and significant reduction in joint space width.⁴ Hip joint OA is considered moderate when joint space is less than 2.5mm and severe when joint space is less than 1.5mm.¹⁶ The development of newer imaging techniques, such as gadolinium enhanced magnetic resonance imaging, has been suggested as a method to detect deficiencies in cartilage structure that may represent early arthritic changes in young patients.¹⁰⁵

In addition to joint space narrowing, other criteria including osteophytic spurs and subchondral sclerosis also are used to identify patients with hip OA.^{6, 19} The Kellgren/Lawrence scale has been used to classify degenerative findings associated with hip OA. The scale consists of 4 grades; Grade 1: no radiographic evidence of OA; Grade 2: doubtful narrowing of joint space and possible (minute) osteophytes; Grade 3: moderate definite osteophytes, definite moderate narrowing of joints space; Grade 4: large osteophytes, severe joint space narrowing, subchondral sclerosis, and definite deformity of bone contour.^{101, 102} A potential caveat when using the Kellgren/Lawrence scale is spurs or osteophytes are emphasized,^{140, 146} and not all patients with hip OA have osteophytes.

CLINICAL GUIDELINES

Examination**OUTCOME MEASURES**

The most commonly used outcome measure for hip OA is the Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC).¹⁴ The WOMAC has been validated and its reliability has been shown in many different studies and in many different countries.^{11, 171, 186, 201, 214} The index consists of 24 questions (5 about pain, 2 about stiffness, and 17 about physical function), which are assigned a score of between 0 (extreme) and 4 (none). Individual question scores are then summed to form a raw score ranging from 0 (best) to 96 (worst). Finally, raw scores are normalized by multiplying the raw score by 100/96 creating a score of between 0% (best) to 100% (worst). Test-retest reliability of the WOMAC as measured by intraclass correlation coefficients has been shown to be good with ICCs ranging between 0.74 and 0.89.¹⁶⁸ The MDC₉₅ for the WOMAC ranges from 20-25%.¹² The minimal clinically important difference (MCID) for the WOMAC score, as a change from the baseline score, has been reported in the range of 12% to 22%.^{8, 9, 46, 212} Angst et al⁷ found the MCID for improvement in a sample of 192 patients with lower extremity OA to be in the range of 17 to 22% change from the baseline score. A prospective cohort study consisting of 223 patients diagnosed with hip or knee OA from a rehabilitation clinic found an 18% change from the baseline score as the MCID for the WOMAC.⁹

I

The Lower Extremity Functional Scale (LEFS) is another reliable and valid outcome measure that is often administered to patients with hip OA.¹⁷ The LEFS uses an ordinal scale from 0 (“extreme difficulty or unable to perform the activity”) to 4 (“no difficulty”) for the patient to rate their ability to perform 20 different activities, such as for example: getting into or out of the bath tub, sitting for 1 hour, squatting, and rolling over in bed. The total score range from 0 to 80 with 80 representing maximum function based on the scale. The reliability and validity of the LEFS have been shown to be good when determined using a sample of 107 patients with lower extremity musculoskeletal problems. In that same study, the minimal detectable change and minimal clinical important difference was 9 scale points.¹⁷

I

Another often used functional outcome measures is the Harris Hip Score.⁷⁷ The Harris Hip Score is derived from scoring 10 different variables including pain, ROM, gait/limp, gait distance, function, activities of daily living, and deformity. Scores range from 0-100 (0 worst to 100 best).⁷⁷

I

A

Clinicians should use validated functional outcome measures, such as the Western Ontario and McMaster Universities Osteoarthritis Index, Lower Extremity Functional Scale, and the Harris Hip Score before and after interventions intended to alleviate the impairments of body function and structure, activity limitations, and participation restrictions associated with hip osteoarthritis.

ACTIVITY LIMITATION AND PARTICIPATION RESTRICTION MEASURES

6-MINUTE WALK TEST

ICF category:	Measurement of activity limitation – walking long distances
Description:	A physical performance measure which assesses how far a person can walk in 6 minutes. ⁴⁸
Measurement Method	During the performance of the 6 Minute Walk Test (6MWT), patients are instructed to cover as much distance as possible during the 6-minute time frame with the opportunity to stop and rest if required. The test is conducted on an unobstructed level surface. The course is marked off in meters and the distance traveled by each subject is measured to the nearest meter. Standardized verbal encouragement, "You are doing well, keep up the good work" is provided at 60-second intervals. Patients are permitted to use their regular walking aids if needed. ¹⁰⁴
Nature of variable	Continuous
Units of measurement	Meters
Measurement properties	The 6MWT showed high test-retest reliability ($ICC_{2,1}$ of 0.95-0.97). ¹⁷² Kennedy et al ¹⁰⁰ also showed high test-retest reliability for the 6MWT with $ICC_{2,1}$ of 0.94 [95% CI=0.88-0.98]. The minimal detectable change (MDC_{90}) for the 6MWT determined from a sample of 150 subjects with hip and knee OA of which 69 underwent total hip arthroplasty is 61.34 meters. ¹⁰⁴

SELF-PACED WALK TEST

ICF category:	Measurement of activity limitation – walking short distances
Description:	A physical performance measure which assesses how fast a person can walk, for 4 meters and for 40 meters
Measurement Method	During the performance of the Self-Paced Walk Test, (SPWT) patients are instructed to "walk as quickly as you can without overexerting yourself" and are timed with a stopwatch while they walk 2 lengths (turn excluded) of a 20 meter indoor course. ¹⁰⁴
Nature of variable	Continuous
Units of measurement	Seconds
Measurement properties	The test-retest reliability of the SPWT for 40 meters has been examined by Kennedy et al. ¹⁰⁴ They found ICC of 0.91 (95% CI=0.81-0.97). Kennedy et al. ¹⁰⁴ also showed the SPWT was responsive in detecting deterioration and improvement in the early postoperative time period following arthroplasty. The MDC ₉₀ for the 40 meter SPWT determined from a sample of 150 subjects with hip and knee OA of which 69 underwent total hip arthroplasty is 4.04 seconds. ¹⁰⁴ The MDC ₉₅ for the 4 meter SPWT is 0.1 m/second in a cohort of 492 older adults. ¹⁵¹

STAIR MEASURE

ICF category:	Measurement of activity limitation – climbing
Description:	A physical performance measure, which assesses how well a person, can ascend and descend a flight of stairs.
Measurement Method	During the performance of the Stair Measure (SM) patients are instructed to ascend and descend 9 steps (step height, 20 cm) in their usual manner, and at a safe and comfortable pace. ¹⁰⁴
Nature of variable	Continuous
Units of measurement	Seconds
Measurement properties	The test-retest reliability of the SM has been examined by Kennedy et al. ¹⁰⁴ They found an ICC of 0.90 (95% CI=0.79-0.96). ¹⁰⁴ Kennedy et al. ¹⁰⁰ also showed the SM was responsive in detecting deterioration and improvement in the early postoperative time period following arthroplasty. ¹⁰⁴ The MDC ₉₀ for the SM, determined from a sample of 150 subjects with hip and knee OA of which 69 underwent total hip arthroplasty, is 5.5 seconds. ¹⁰⁴

TIMED UP AND GO TEST

ICF category:	Measurement of activity limitations – getting in and out of a seated position, walking short distances
Description:	A physical performance measure which assesses how well a person can get up from a chair with arm rests walk a short distance (3 meters), turn around, return, and then sit down again. ¹³⁰
Measurement Method	During the performance of the Timed Up and Go Test (TUG) the patient sits in a chair with arm rests and is asked to stand up from the chair and walk as quickly and safely as possible to a cone 3 meters away, turn, walk back to the chair and sit down again. The performance of this test is timed.
Nature of variable	Continuous
Units of measurement	Seconds
Measurement properties	There was good agreement among observers on the subjective scoring of the TUG, and good correlation with the Berg Balance Scale, gait speed, Barthel's Index of activities of daily living, and predicted patient's ability to walk outside safely. ¹⁵³ Podsiadlo ¹⁴⁷ showed the TUG had good intertester and intratester reliability (ICC=.99). Steffen et al ¹⁶⁶ also showed the TUG had high test-retest reliability (ICC _{2,1} =.95-.97). Podsiadlo ¹⁴⁷ provided evidence for the criterion-related validity of the TUG by showing it correlates well with other functional scales. Kennedy et al ¹⁰⁰ showed the TUG was responsive in detecting deterioration and improvement in postoperative time period following arthroplasty. The MDC ₉₀ for the TUG, determined from a sample of 150 subjects with hip and knee OA of which 69 underwent total hip arthroplasty, is 2.5 seconds. ¹⁰⁴

PHYSICAL IMPAIRMENT MEASURES

PASSIVE HIP INTERNAL AND EXTERNAL ROTATION AND HIP FLEXION

ICF category:	Measurement of impairment of body function – mobility of a single joint
Description:	The amount of passive hip rotation and passive hip flexion measured prone and supine respectively. Although assessing the range in all 6 directions (3 planes) of hip motion is important in patients with hip OA, for brevity we included the 3 most commonly limited hip motions.
Measurement Method	<p><u>Hip Internal and External Rotation</u></p> <p>The patient is positioned prone with feet over the edge of the treatment table. The hip measured is placed in 0° of abduction, and the contralateral hip is placed in about 30° of abduction. The reference knee is flexed to 90°, and the leg is passively moved to produce hip rotation. The movement arm of the goniometer is aligned vertically along the shaft of the tibia while the stationary arm is aligned along an imaginary vertical line. Manual stabilization is applied to the pelvis to prevent pelvic movement and also at the tibiofemoral joint to prevent motion (rotation or abduction/adduction), which could be construed as hip rotation.⁷⁸ The tibia is then moved in the frontal plane to produce hip internal and external rotation. The motion is stopped and measurements taken when the extremity achieves its end range of passive hip rotation or when pelvic movement is necessary for additional movement of the leg. An inclinometer may also be used to measure hip rotation. The inclinometer is first “calibrated” by placing it along the distal shaft of the vertically aligned tibia, just proximal to the medial malleolus and then setting the inclinometer dial to zero. Then, leg is passively moved to produce hip rotation and inclinometer measure is taken when the extremity achieves its end range of passive internal and external rotation.⁴⁷</p> <p><u>Hip Flexion</u></p> <p>With the patient in the supine position, the hip is passively flexed with the movement arm of the goniometer along the long axis of the femur and the stationary arm of the goniometer along the long axis of the trunk, while stabilizing the lumbar spine to avoid any posterior pelvic tilt.⁸³</p>
Nature of variable	Continuous and Ordinal
Units of measurement	Degrees and 0-10 NPRS
Measurement properties	Limited ROM is associated with high levels of disability in patients with hip OA. ¹⁷⁴ The reliability for hip rotation and hip flexion ROM measurements has been shown to be excellent, ICC of 0.95 to 0.97 ⁴⁷ for rotation and ICC = 0.94 [95% CI= 0.89-0.97] ³⁰ for flexion. ROM measurements in 22 individuals with hip OA demonstrated excellent intrarater test-retest reliability (ICC, .97) for hip flexion. ¹⁵⁴ Croft et al ³⁸ showed good agreement among 6 testers when assessing for hip rotation

and hip flexion in patients with hip OA. Steultjens et al¹⁷⁴ also showed good reliability when assessing the hip joint in patients with OA. The MDC₉₅ for hip flexion, determined using 22 patients with knee OA and 17 subjects without lower extremity symptoms or known pathology, is 5°, meaning any change more than 5° is considered to be change beyond measurement error.³⁰ The MDC₉₅ for pain with hip flexion is 1.2 on the 0-10 NPRS.³⁰

HIP ABDUCTOR MUSCLES STRENGTH TEST

ICF category:	Measurement of impairment of body function – power of isolated muscles and muscle groups
Description:	A test to determine the strength of the hip abductor muscles.
Measurement Method	The hip abductor muscles strength test is performed with the subject in the supine position and the hip in a neutral position of flexion/extension, abduction/adduction, and external/internal rotation. A “make” test using a hand held dynamometer is used by asking the subject to push the most they can against the hand held dynamometer applied on the lateral aspect of the distal thigh, just above the knee. The hip abductor muscles may also be tested in the side lying position with the hip in abduction and slight extension. A “break” test is performed by the tester applying force via the hand-held dynamometer applied on the lateral aspect of the distal thigh just above the knee. The direction of force application is toward adduction and slight flexion while the pelvis is stabilized with the other hand. ¹⁰³
Nature of variable	Continuous
Units of measurement	Force in Newton’s
Measurement properties	Interrater and intrarater reliability of force measurements obtained from college age women were excellent using a hand held dynamometer were excellent for the abductor muscle (intrarater ICC, .88-.96; interrater ICC, .90-.95). ³¹ Force measurements of hip abductors in 22 individuals with hip OA demonstrated good intrarater test-retest reliability (ICC, .84). ¹⁵⁴ The MDC ₉₅ , determined from a sample of 90 subjects (age 22-70 years) without any previous musculoskeletal problems, was 5.4% of body weight for males and 5.3% for females. ²¹⁵

THE FABER (PATRICK'S) TEST

ICF category:	Measurement of impairment of body function – pain in joints.
Description:	A test to determine the irritability of the hip joint.
Measurement Method	The FABER test is administered with the subject in supine, the heel of the lower extremity to be tested placed over the opposite knee. The hip joint is passively externally rotated and abducted by the examiner applying manual pressure over the ipsilateral knee, while stabilizing the contralateral innominate with the opposite hand. After being zeroed against a wall, the inclinometer is placed on the medial aspect of the tibia of the tested lower extremity, just distal to the medial tibial condyle. ROM measurement is taken at the point of maximal passive resistance or at the point where the patient stopped the test secondary to pain. ³⁰ The patient is also asked to rate the location of the pain as well as the amount of pain experienced during the movement on a 0-10 NPRS.
Nature of variable	Nominal and Continuous
Units of measurement	Degrees and 0-10 NPRS
Measurement properties	Intrarater reliability of ROM (ICC, 0.96) [95% CI=0.92-0.98] and pain (ICC, 0.87) [95% CI= 0.78-0.94] measurements was excellent for the FABER test. ³⁰ Cibulka et al ²⁵ found the FABER test was responsive in detecting improvement in ROM and in report of pain in patients with hip pain. The MDC ₉₅ , determined from a sample of 22 patients with knee OA and 17 subjects without lower extremity symptoms or known pathology is 8° for ROM, and 1.6 points ³⁰ on the NPRS. The clinically important difference for the NPRS has been shown to be a reduction of 2 points. ^{27, 51}

THE SCOUR TEST

ICF category:	Measurement of impairment of body function – pain in joints.
Description:	A test to determine the irritability of the hip joint.
Measurement Method	The hip Scour test is performed with the patient lying in the supine position while the clinician flexes and adducts the hip until resistance to movement is detected. The clinician then maintains flexion into resistance and gently moves the hip into abduction, then bringing the hip through 2 full arcs of motion. If the patient reports no pain, then the examiner repeats the test while applying long-axis compression through the femur. This test must be administered with some caution so as to not irritate the hip joint. The patient is asked to rate the pain experienced during the movement on a 0-10 NPRS. ³⁰
Nature of variable	Nominal
Units of measurement	0-10 NPRS
Measurement properties	The intratester reliability of the Scour test is good (ICC, 0.87) [95%CI= .76-.93) for rating of hip pain. ³⁰ The clinically important difference for the NPRS has been shown to be a reduction of 2 points. ^{27, 51} The only MDC ₉₅ determined from a sample of 22 patients with knee OA and 17 subjects without lower extremity symptoms or known pathology for the Scour test for pain is a change of more than 1.6 points on the 0-10 NPRS. ³⁰

PROGNOSIS

In most cases, OA of the hip usually progresses slowly²⁰⁰ with total hip arthroplasty (THA) being the primary clinical endpoint for individuals with severe hip OA.⁶⁹ The prognosis of hip OA depends primarily on the extent of radiographic evidence of hip OA.¹⁶⁹ The severity and progression of hip OA is commonly assessed with the Kellgren/Lawrence scale of joint space narrowing on plain film radiographs. A patient's baseline Kellgren/Lawrence radiographic grade is an important predictive factor for having THA.^{44, 162, 202} Reijman et al¹⁵⁷ found that a Kellgren/Lawrence score of II or higher is a strong predictor of progression in patients with hip OA. Gossec et al⁷⁰ reported that a Kellgren/Lawrence grade of III had an odds ratio of 3.3 and a grade of IV had an odds ratio of 5.3 that patients would have a total hip replacement. Gossec et al⁷⁰ also reported that the most important predictive factors of having a total hip replacement include Kellgren/Lawrence radiographic grades of III or higher, a high global assessment of pain, and previous trial of non-steroidal anti-inflammatory drugs.^{70, 4, 72} Altman et al⁵ has suggested that the measurement of individual radiographic features may be superior to the Kellgren/Lawrence global measurement in detecting arthritic progression. In OA of the hip, a single anteroposterior radiograph assessing for joint space narrowing and cyst formation yielded high sensitivity in detecting change. The minimal clinically important difference for joint space narrowing progression is $> \text{or} = 0.5\text{mm/year}$.^{4, 43} The rate of joint space narrowing change in patients with slow developing hip OA is $< 0.2\text{mm/year}$ and in patients with rapid developing hip OA $> 0.2\text{mm/year}$.⁶⁶ In summary, joint space narrowing and the Kellgren/Lawrence scale are important prognostic predictors of OA while joint space narrowing may be the best indicator of structural OA progression in patients with hip OA.

I

Other indicators for performing a THA include previous use of non steroidal anti-inflammatory drugs (NSAID's) and pain of at least 47 mm on a 100 mm pain scale for over 6 months.⁷⁰ Pain and function appear to be important criteria when considering THA.²¹⁷

I

THA is the most common surgical procedure for end-stage hip OA. Despite the success of total joint replacement of the hip and knee over the last 30 plus years, the criteria for when to perform such surgery are not clear.³ There have been several attempts to develop guidelines to determine the appropriate time to perform joint replacement surgery, however few are supported by research.³ Currently, there is no consensus on the appropriate time to recommend surgery as a clinical endpoint.³ However, the Group for the Respect of Ethics and Excellence in Science (GREES) suggests that conservative intervention has failed if a patient does not experience a reduction in symptoms, such as a 20-25% improvement on the Pain subscale of the WOMAC, and has a progressive loss of joint space of between .3-.7 mm/year.³

V

CLINICAL GUIDELINES

Interventions

A variety of interventions have been described for the treatment of hip OA and there is fair evidence from randomized clinical trials and systematic reviews to support the benefits of physical therapy intervention in these patients.

ANTI-INFLAMMATORY AGENTS

Both over-the-counter and prescribed anti-inflammatory agents including non-steroidal anti-inflammatory drugs (NSAIDs), Cox-2 inhibitors, and steroid injections are recommended as part of a multidisciplinary treatment approach to hip OA.²¹⁸ Randomized clinical trials evaluating the use of NSAIDs have shown NSAIDs can be effective for the temporary relief of symptoms and improvement in function in patients with hip OA.¹⁹² However, it should be noted this class of drugs is not without risk for serious adverse events including increased gastrointestinal bleeding.⁶⁷

I

There is evidence to support the use of depot corticosteroid injection in patients with hip OA to provide short-term pain relief.^{157, 165} A recent placebo-controlled trial confirmed depot corticosteroid injection can be an effective treatment of pain in hip OA, with benefits lasting up to 3 months.¹⁰⁷

I

Some evidence does suggest some NSAIDs may increase the progression of hip OA by decreasing glycosaminoglycan synthesis,^{22, 86, 149, 161} however further studies are needed.

I

ALTERNATIVE/COMPLEMENTARY MEDICATION

Glucosamine and other similar supplements are commonly suggested for individuals with hip OA. To date, randomized controlled trials evaluating the use of glucosamines have shown mixed results.^{132, 133, 208} Most of the positive results are for short term improvement in pain and in function.¹³⁴ A recent meta-analysis of chondroitin (a specific glucosamine commonly found in articular cartilage) for OA of the hip indicates the symptomatic benefit of chondroitin is minimal or non-existent and the use of chondroitin in clinical practice should be discouraged.¹⁶⁰ Glucosamines (also called glycosaminoglycans) are an important component of normal connective tissue physiology, however, the short or long-term use of glucosamines is not recommended at this time in patients with hip OA.

I

There is some evidence to support the short-term use of injectable viscosupplementation with hyaluronic acid into hip joint of patients with hip OA.³⁴ Despite a paucity of evidence, the use of injectable synthetic hyaluronic acid (hyaluronan) into the hip joint has been shown to be an effective treatment for symptomatic hip OA.^{33, 197, 198} Evidence also shows injectable hyaluronan works best in mild to moderate hip OA, especially when conservative therapy has failed.⁴⁰ A recent published meta-analysis suggests the benefit of hyaluronan for the treatment of hip osteoarthritis,³⁴ but so far it is only approved by the FDA for the knee. More controlled studies are needed to show its effectiveness in patients with hip OA.³⁴

PATIENT EDUCATION

Studies have shown the benefit of patient education in the self-management of patients with arthritis in decreasing pain, improving function, reducing stiffness and fatigue, and reducing medical usage.^{26, 58, 87, 88, 218} A meta-analysis has shown patient education can provide on average 20% more pain relief when compared to NSAIDs alone in patients with hip OA or rheumatoid arthritis.¹⁷⁶

I

An approach, called Hip School, that includes primarily patient education as an intervention has been shown to be effective in a preliminary study for patients with signs and symptoms of hip OA.¹⁰⁶ The Hip School highlights the need for educating patients with hip OA especially understanding the importance of preserving hip ROM and muscle function, understanding what therapy is effective and what is not, and when surgery is likely indicated.

II

B

Clinicians should consider the use of patient education to teach activity modification, exercise, weight reduction when overweight, and methods of unloading the arthritic joints.

FUNCTIONAL, GAIT, AND BALANCE TRAINING

Patients with hip OA often have gait abnormalities such as asymmetry in weight bearing and step length.²⁹ Assistive device are often used in patients with hip OA to reduce the pain and activity limitations associated with this condition.¹⁹⁹ A cane in the contralateral hand and choosing to carry loads in the ipsilateral hand has been shown to be effective in reducing hip abductor muscle activity^{135, 145} and acetabular contact pressures.^{135, 145} One study has shown a cane in the opposite hand can reduce hip load, reduce hip pain, and improve function in patients with hip OA.¹⁴⁴

II

Functional, gait, and balance training is recommended to address impairments of proprioception, balance, and strength, which are all commonly found in individuals with lower extremity arthritis. These deficits can contribute to higher fall risk scores in older individuals.¹⁷⁵ Functional training of a small cohort of elderly individuals with lower extremity impairment demonstrated improved functional performance. Subjects underwent a program consisting of exercises simulating activities of daily living (such as gait, rising from a chair, reaching, stepping, and squatting down) performed at 3 different speeds (self selected, fast, and slow) with progressive levels of difficulty. When subjects completed 1 task level correctly and without fatigue the next level was introduced.¹³⁶

II

C

Functional, gait, and balance training, including the use of assistive devices such as canes, crutches, and walkers, can be used in patients with hip osteoarthritis to improve function associated with weight bearing activities.

MANUAL THERAPY

Some evidence exists for using manual therapy to increase hip joint ROM and short-term pain reduction in patients with hip OA, especially in patients, who do not have signs of severe hip OA (eg, osteophytes and significant joint space narrowing).⁸²

One study has recommended mobilization/manipulation as a component of the management program for patients with hip OA.⁸² This randomized controlled trial compared the use of manual therapy and therapeutic exercises in patients with hip OA.⁸² The manual therapy session consisted of 1) stretching techniques of shortened muscles surrounding the hip joint, 2) traction of the hip joint, 3) traction manipulation in each limited position (a high velocity thrust technique). All manipulations were repeated during each session until the therapist concluded optimal results of the session were achieved. The focus of the therapeutic exercise intervention was to improve hip ROM, muscle length, and strength along with walking endurance. The outcomes for hip function (Harris Hip Score), ROM, and pain as measured by the visual analogue scale were compared for specific subgroups of hip OA depending on limited function, ROM or level of pain.⁸² After 5 weeks of intervention, the success rate (primary outcome) of manual therapy was 81% versus 50% for exercise therapy (odds ratio: 1.92; 95% CI: 1.30-2.60).⁸² Manual therapy was found to be superior to exercise therapy in some patients with hip OA but was not shown to be any more effective than exercise in patients with highly limited function, ROM, or high levels of pain.⁸² When intervention stopped, the improvements in function declined after 5 weeks. However, some improvement lasted up to 29 weeks for the patients in the manual therapy group.⁸²

I

MacDonald et al¹²⁵ published a case series describing the outcomes of individual patients with hip OA treated with manual physical therapy and exercise. The series included 7 patients diagnosed with hip OA on the basis of the clinical examination. All patients were treated with manual physical therapy followed by exercises to increase hip strength and ROM. Six of 7 patients completed a Harris Hip Score at initial examination and discharge from physical therapy. Patients exhibited reductions in pain and increases in passive ROM, as well as a clinically meaningful improvement in function.¹²⁵

IV

Harding et al,⁷⁶ in a study using cadaveric models, showed that a posterior-anterior (P/A) mobilization of the hip produced about 1 mm of movement in the hip joint when using a force of 356N. Distal distraction of the hip, however, created motion ranging from 2-7 mm of displacement when using forces between 89 to 356N.⁷⁶ This cadaveric study suggests that when attempting to mobilize the hip joint, the amount of movement produced at the hip most likely depends on the direction the joint is mobilized.⁷⁶

Risks of adverse events associated with manual therapy of the hip typically include self-limiting soreness of the hip region. There are no studies documenting an increased risk for serious adverse events associated with manual therapy of the hip.

B

Clinicians should consider the use of manual therapy procedures to provide short-term pain relief and improve hip mobility and function in patients with mild hip osteoarthritis.

FLEXIBILITY, STRENGTHENING, AND ENDURANCE EXERCISES

There are 3 categories of exercise therapy employed for OA: ROM / flexibility exercises, muscle strengthening exercises, and aerobic conditioning / endurance exercises. Often all 3 types of exercises are utilized jointly for patients with hip OA. Adequate joint motion and elasticity of periarticular tissues are necessary for cartilage nutrition and health, protection of joint structures from damaging impact loads, function, and comfort in daily activities. Exercise to regain or maintain motion and flexibility is achieved by routines of low-intensity, controlled movements which do not cause increased pain.⁵² Muscle weakness around an osteoarthritic joint is a common finding.¹⁷³ Progressive resistive / strengthening exercises load muscles in a graduated manner to allow for strengthening while limiting tissue injury. Aerobic exercise has been shown to be helpful in patients with hip OA.²¹⁸ Aerobic exercises are usually designed to provide a workload to the cardiovascular and pulmonary system at 60% to 80% of maximal capacity and sustained for duration of at least 20 minutes.²¹⁸

Range of motion and strengthening exercises have been advocated by many authors as a component of the management for patients with OA of the hip.^{32, 56, 80, 85, 125, 137, 152, 159, 166, 167, 179, 193, 195, 196, 218} Van Baar et al¹⁹⁴ showed that exercise is effective in patients with OA of the hip using an exercise program previously reported by Oostendorp et al¹⁴⁸ that consisted of flexibility, hip muscle strengthening, and an aerobic exercise program.¹⁴⁸ The emphasis of the stretching was on hip muscles including the iliopsoas, rectus femoris, and hip adductors. Before stretching it was advised to heat the specific muscle and then stretch gently without excessive force for 15-30 seconds performed 5 to 10 times preferably daily but at least 3 times a week. Hip muscle strengthening was performed with either free weights or exercise machines.^{148, 194} Depending on the therapist's findings on evaluation and the patient's needs and goals, the specific type, intensity, frequency, and duration of exercise was determined. The typical exercise prescription ranged from 1 to 3 times per week with a duration of 30 minutes for each exercise session for 12 weeks.¹⁹⁴ Exercise treatment could be discontinued within the 12 week period if according to the therapist, the treatment goals were achieved. When patients stopped exercising after 12 weeks, Van Baar et al¹⁹⁴ reported that the beneficial effects of reduced pain, less use of medication, and improved function declined, losing any earlier gain that were made.¹⁹⁵ A confounding factor of the Van Baar et al¹⁹⁵ trial was that data from patients with hip OA were pooled with data from patients with knee OA. Therefore, determining specific treatment effect in patients with hip OA was not possible.

Many of the published articles on exercise report findings that combines subjects with either hip or knee OA. However, in a recent meta-analysis,

II

II

Hernandez-Molina et al⁷⁹ contacted authors of several studies and obtained data which only pertained to hip OA. They found 9 articles where hip strengthening exercises showed a beneficial effect in reducing pain and improving function in patients with hip OA.⁷⁹ In another study moderate evidence was found for long-term effectiveness on reduced pain, improved self-reported physical function, and improved observed physical function with exercise for patients with hip OA.¹⁵² Studies have also shown aerobic exercise may offer additional improvement in function when combined with stretching and strengthening.^{155,}
218

Minor et al¹³⁸ studied 120 patients with well-defined rheumatoid arthritis (n = 40) or OA (n = 80) of the hip, knee, or ankle. Patients received 1 of 3 interventions: a stretching and strengthening exercise program alone, or the same program combined with 1 of 2 aerobic exercises: pool activities or walking. All patients participated in a 12-week program, which met 3 times each week for 1 hour, and all performed supervised ROM and isometric exercises. The two groups also performed up to 30 minutes of walking or pool exercises to increase their heart rates to 60% to 80% of each person's estimated baseline maximum. At 12 weeks, 80% of study participants were assessed for aerobic capacity, flexibility (trunk bending), function tests (exercise endurance on a treadmill test, time to walk 50 feet, and reported daily activity), and self-reported health status using the Arthritis Impact Measurement Scale (AIMS), an arthritis-specific functional status instrument that reliably measures psychological health, physical health, and pain. The pool and walking programs increased aerobic capacity. The AIMS scores based on physical activity anxiety and depression subscales improved from 5% to 12% in both groups, significantly more than for the OA than the RA group. There were no significant differences between the groups for the pain subscale. Both groups decreased their times to walk 50 feet by 12%.

The use of aquatic exercise (hydrotherapy) in the treatment of patients with OA of the hip has been assessed.^{13, 32, 55, 57, 71, 80} Aquatic exercise appears to have some beneficial short-term effects for patients with hip and/or knee OA while no long-term effects have been documented.¹³ In a recent study Hinman et al⁸⁰ used a randomized controlled trial and compared a 6-week program of aquatic physical therapy to no intervention. The aquatic physical therapy group demonstrated significantly less pain and improved physical function, strength, and quality of life after the intervention.⁸⁰ However, effect size calculations revealed only small benefits of aquatic physical therapy for pain, stiffness, right hip abductor strength, and quality of life, and doubtful clinical benefits for physical function and left hip abductor strength.⁸⁰ Patients who have an intolerance to land based exercise because of pain or obesity may better tolerate aquatic based exercise.⁸⁰

B

Clinicians should consider the use of flexibility, strengthening, and endurance exercises in patients with hip osteoarthritis.

II

II

CLINICAL GUIDELINES

SUMMARY OF RECOMMENDATIONS

Pathoanatomical Features

B

Clinicians should assess for impairments in mobility of the hip joint and strength of the surrounding muscles, especially the gluteus medius, when a patient presents with hip pain.

Risk Factors

A

Clinicians should consider age, hip developmental disorders, and previous hip joint injury as risk factors for hip osteoarthritis.

Diagnosis/Classification

A

Moderate lateral or anterior hip pain during weight bearing, in adults over the age of 50 years, with morning stiffness less than 1 hour, with limited hip internal rotation and hip flexion by more than 15° when comparing the painful to the non painful side are useful clinical findings to classify a patient with hip pain into the International Statistical Classification of Diseases and Related Health Problems (ICD) category of unilateral coxarthrosis and the associated International Classification of Functioning, Disability, and Health (ICF) impairment-based category of hip pain (b2816 Pain in joints) and mobility deficits (b7100 Mobility of a single joint).

Differential Diagnosis

E

Clinicians should consider diagnostic classifications other than osteoarthritis of the hip when the patient's history, reported activity limitations, or impairments of body function and structure are not consistent with those presented in the diagnosis/classification section of this guideline - or - when the patient's symptoms are not diminishing with interventions aimed at normalization of the patient's impairments of body function.

Examination – Outcome Measures

A

Clinicians should use validated functional outcome measures, such as the Western Ontario and McMaster Universities Osteoarthritis Index, Lower Extremity Functional Scale, and the Harris Hip Score before and after interventions intended to alleviate the impairments of body function and structure, activity limitations, and participation restrictions associated with hip osteoarthritis.

Examination – Activity Limitation and Participation Restriction Measures

A

Clinicians should utilize easily reproducible physical performance measures, such as the 6 Minute Walk, Self-Paced Walk, Stair Measure, and Timed Up and Go to assess activity limitation and participation restrictions associated with their patient's hip pain and to assess the changes in the patient's level of function over the episode of care.

Interventions – Patient Education

B

Clinicians should consider the use of patient education to teach activity modification, exercise, weight reduction when overweight, and methods of unloading the arthritic joints.

Interventions – Functional, Gait, and Balance Training

C

Functional, gait, and balance training, including the use of assistive devices such as canes, crutches, and walkers, can be used in patients with hip osteoarthritis to improve function associated with weight bearing activities.

Interventions – Manual Therapy

B

Clinicians should consider the use of manual therapy procedures to provide short-term pain relief and improve hip mobility and function in patients with mild hip osteoarthritis.

Interventions – Flexibility, Strengthening and Endurance Exercises

B

Clinicians should consider the use of flexibility, strengthening and endurance exercises in patients with hip osteoarthritis

AFFILIATIONS & CONTACTS

Authors:

Michael T. Cibulka, DPT, OCS
Assistant Professor
Physical Therapy Program
Maryville University
St. Louis, MO 63141
mcibulka@maryville.edu

Douglas M. White, DPT
Principal & Consultant
Milton Orthopaedic & Sports Physical
Therapy, PC
101 Blue Hills Parkway
Milton, MA 02186
dr.white@miltonortho.com

Judith Woehrle, PT, PhD
Director, Physical Therapy Program
Maryville University
650 Maryville University Dr
St Louis, MO 63141
woehrle@maryville.edu

Marcie Harris-Hayes, DPT
Assistant Professor of Physical Therapy and
Orthopaedic Surgery
Washington University School of Medicine
St. Louis, MO 63108
harrisma@wustl.edu

Keelan Enseki, PT, MS
Centers for Rehab Services
University of Pittsburgh Medical Center
Center for Sports Medicine
3200 South Water Street
Pittsburgh, PA 15203
ensekikr@upmc.edu

Timothy L. Fagerson, DPT, MS
Director, Spine Orthopaedic Sport Physical
Therapy
148 Linden St., Suite B-8
Wellesley, MA 02482
fagerson@verizon.net

Reviewers

Roy D. Altman, MD
Professor of Medicine
Division of Rheumatology and Immunology
David Geffen School of Medicine at UCLA
Los Angeles, California
journals@royaltman.com

Anthony Delitto, PT, PhD
Professor and Chair
School of Health & Rehabilitation Sciences
University of Pittsburgh
Pittsburgh, Pennsylvania
delitto@upmc.edu

John Dewitt, DPT
Director of Physical Therapy Sports Medicine
Residency
The Ohio State University
Columbus, Ohio
john.dewitt@osumc.edu

Amanda Ferland, PT
Clinic Director
MVP Physical Therapy
Federal Way, Washington
aferland@mvppt.com

Helene Fearon, PT
Principal and Consultant
Rehabilitation Consulting & Resource Institute
Phoenix, Arizona
helenefearon@myrehabconsultants.com

Kathleen Kline Mangione, PT, PhD
Professor, Physical Therapy
Arcadia University
450 S Easton Rd
Glenside, PA 19038
mangione@arcadia.edu

Marian A. Minor, PT, PhD
Professor, Department Chair
Physical Therapy

James Slover, MD, MS
Assistant Professor
NYU Hospital for Joint Diseases
301 East 17th St. Suite 1616
New York, NY 10003
james.slover@nyumc.org

Joseph J. Godges, DPT
ICF Practice Guidelines Coordinator
Orthopaedic Section APTA, Inc.
La Crosse, Wisconsin
icf@orthopt.org

School of Health Professions
University of Missouri-Columbia
106A Lewis Hall
Columbia, MO 65211
minorm@health.missouri.edu

Joy MacDermid, PT, PhD
Associate Professor
School of Rehabilitation Science
McMaster University
Hamilton, Ontario, Canada
macderj@mcmaster.ca

James W. Matheson, DPT
Minnesota Sport and Spine Rehabilitation
Burnsville, Minnesota
jw@eipconsulting.com

Philip McClure, PT, PhD
Professor
Department of Physical Therapy
Arcadia University
Glenside, Pennsylvania
mcclure@arcadia.edu

Paul Shekelle, MD, PhD
Director
Southern California Evidenced-Based Practice
Center
Rand Corporation
Santa Monica, California
shekelle@rand.org

A. Russell Smith, Jr., PT, EdD
Chair
Athletic Training & Physical Therapy
University of North Florida
Jacksonville, Florida
arsmith@unf.edu

Leslie Torburn, DPT
Principal and Consultant
Silhouette Consulting, Inc.
Redwood City, California
torburn@yahoo.com

References:

1. Abraham E, Gonzalez MH, Pratap S, Amirouche F, Atluri P, Simon P. Clinical implications of anatomical wear characteristics in slipped capital femoral epiphysis and primary osteoarthritis. *J Pediatr Orthop*. 2007;27:788-795.
2. Altman R, Alarcon G, Appelrouth D, et al. The American College of Rheumatology criteria for the classification and reporting of osteoarthritis of the hip. *Arthritis Rheum*. 1991;34:505-514.
3. Altman RD, Abadie E, Avouac B, et al. Total joint replacement of hip or knee as an outcome measure for structure modifying trials in osteoarthritis. *Osteoarthritis Cartilage*. 2005;13:13-19.
4. Altman RD, Bloch DA, Dougados M, et al. Measurement of structural progression in osteoarthritis of the hip: the Barcelona consensus group. *Osteoarthritis Cartilage*. 2004;12:515-524.
5. Altman RD, Fries JF, Bloch DA, et al. Radiographic assessment of progression in osteoarthritis. *Arthritis Rheum*. 1987;30:1214-1225.
6. Altman RD, Hochberg M, Murphy WA, Jr., Wolfe F, Lequesne M. Atlas of individual radiographic features in osteoarthritis. *Osteoarthritis Cartilage*. 1995;3 Suppl A:3-70.
7. Andrianakos AA, Kontelis LK, Karamitsos DG, et al. Prevalence of symptomatic knee, hand, and hip osteoarthritis in Greece. The ESORDIG study. *J Rheumatol*. 2006;33:2507-2513.
8. Angst F, Aeschlimann A, Michel BA, Stucki G. Minimal clinically important rehabilitation effects in patients with osteoarthritis of the lower extremities. *J Rheumatol*. 2002;29:131-138.
9. Angst F, Aeschlimann A, Steiner W, Stucki G. Responsiveness of the WOMAC osteoarthritis index as compared with the SF-36 in patients with osteoarthritis of the legs undergoing a comprehensive rehabilitation intervention. *Ann Rheum Dis*. 2001;60:834-840.
10. Arokoski MH, Arokoski JP, Haara M, et al. Hip muscle strength and muscle cross sectional area in men with and without hip osteoarthritis. *J Rheumatol*. 2002;29:2185-2195.
11. Atamaz F, Hegguler S, Oncu J. Translation and validation of the Turkish version of the arthritis impact measurement scales 2 in patients with knee osteoarthritis. *J Rheumatol*. 2005;32:1331-1336.
12. Barr S, Bellamy N, Buchanan WW, et al. A comparative study of signal versus aggregate methods of outcome measurement based on the WOMAC Osteoarthritis Index. Western Ontario and McMaster Universities Osteoarthritis Index. *J Rheumatol*. 1994;21:2106-2112.
13. Bartels EM, Lund H, Hagen KB, Dagfinrud H, Christensen R, Danneskiold-Samsoe B. Aquatic exercise for the treatment of knee and hip osteoarthritis. *Cochrane Database Syst Rev*. 2007;CD005523.
14. Bellamy N, Buchanan WW, Goldsmith CH, Campbell J, Stitt LW. Validation study of WOMAC: a health status instrument for measuring clinically important patient relevant outcomes to antirheumatic drug therapy in patients with osteoarthritis of the hip or knee. *J Rheumatol*. 1988;15:1833-1840.
15. Bierma-Zeinstra SM, Koes BW. Risk factors and prognostic factors of hip and knee osteoarthritis. *Nat Clin Pract Rheumatol*. 2007;3:78-85.

16. Bierma-Zeinstra SM, Oster JD, Bernsen RM, Verhaar JA, Ginai AZ, Bohnen AM. Joint space narrowing and relationship with symptoms and signs in adults consulting for hip pain in primary care. *J Rheumatol*. 2002;29:1713-1718.
17. Binkley JM, Stratford PW, Lott SA, Riddle DL. The Lower Extremity Functional Scale (LEFS): scale development, measurement properties, and clinical application. North American Orthopaedic Rehabilitation Research Network. *Phys Ther*. 1999;79:371-383.
18. Birrell F, Croft P, Cooper C, Hosie G, Macfarlane G, Silman A. Predicting radiographic hip osteoarthritis from range of movement. *Rheumatology (Oxford)*. 2001;40:506-512.
19. Birrell F, Croft P, Cooper C, Hosie G, Macfarlane GJ, Silman A. Radiographic change is common in new presenters in primary care with hip pain. PCR Hip Study Group. *Rheumatology (Oxford)*. 2000;39:772-775.
20. Birrell F, Silman A, Croft P, Cooper C, Hosie G, Macfarlane G. Syndrome of symptomatic adult acetabular dysplasia (SAAD syndrome). *Ann Rheum Dis*. 2003;62:356-358.
21. Boutry N, Paul C, Leroy X, Fredoux D, Migaud H, Cotten A. Rapidly destructive osteoarthritis of the hip: MR imaging findings. *AJR Am J Roentgenol*. 2002;179:657-663.
22. Brandt KD. Effects of nonsteroidal anti-inflammatory drugs on chondrocyte metabolism in vitro and in vivo. *Am J Med*. 1987;83:29-34.
23. Buckland-Wright JC. Quantitative radiography of osteoarthritis. *Ann Rheum Dis*. 1994;53:268-275.
24. Buckwalter JA, Lane NE. Athletics and osteoarthritis. *Am J Sports Med*. 1997;25:873-881.
25. Buckwalter JA, Martin JA. Sports and osteoarthritis. *Curr Opin Rheumatol*. 2004;16:634-639.
26. Callahan LF, Mielenz T, Freburger J, et al. A randomized controlled trial of the people with arthritis can exercise program: symptoms, function, physical activity, and psychosocial outcomes. *Arthritis Rheum*. 2008;59:92-101.
27. Childs JD, Piva SR, Fritz JM. Responsiveness of the numeric pain rating scale in patients with low back pain. *Spine*. 2005;30:1331-1334.
28. Cibulka MT. Determination and significance of femoral neck anteversion. *Phys Ther*. 2004;84:550-558.
29. Cichy B, Wilk M. Gait analysis in osteoarthritis of the hip. *Med Sci Monit*. 2006;12:CR507-513.
30. Cliborne AV, Wainner RS, Rhon DI, et al. Clinical hip tests and a functional squat test in patients with knee osteoarthritis: reliability, prevalence of positive test findings, and short-term response to hip mobilization. *J Orthop Sports Phys Ther*. 2004;34:676-685.
31. Click Fenter P, Bellew JW, Pitts TA, Kay RE. Reliability of stabilised commercial dynamometers for measuring hip abduction strength: a pilot study. *Br J Sports Med*. 2003;37:331-334.
32. Cochrane T, Davey RC, Matthes Edwards SM. Randomised controlled trial of the cost-effectiveness of water-based therapy for lower limb osteoarthritis. *Health Technol Assess*. 2005;9:iii-iv, ix-xi, 1-114.

33. Conrozier T, Bertin P, Bailleul F, et al. Clinical response to intra-articular injections of hylan G-F 20 in symptomatic hip osteoarthritis: the OMERACT-OARSI criteria applied to the results of a pilot study. *Joint Bone Spine*. 2006;73:705-709.
34. Conrozier T, Vignon E. Is there evidence to support the inclusion of viscosupplementation in the treatment paradigm for patients with hip osteoarthritis? *Clin Exp Rheumatol*. 2005;23:711-716.
35. Cooper C, Inskip H, Croft P, et al. Individual risk factors for hip osteoarthritis: obesity, hip injury, and physical activity. *Am J Epidemiol*. 1998;147:516-522.
36. Croft P, Coggon D, Cruddas M, Cooper C. Osteoarthritis of the hip: an occupational disease in farmers. *Bmj*. 1992;304:1269-1272.
37. Croft P, Cooper C, Wickham C, Coggon D. Defining osteoarthritis of the hip for epidemiologic studies. *Am J Epidemiol*. 1990;132:514-522.
38. Croft PR, Nahit ES, Macfarlane GJ, Silman AJ. Interobserver reliability in measuring flexion, internal rotation, and external rotation of the hip using a plurimeter. *Ann Rheum Dis*. 1996;55:320-323.
39. D'Ambrosia RD. Epidemiology of osteoarthritis. *Orthopedics*. 2005;28:s201-205.
40. Dagenais S. Intra-articular hyaluronic acid (viscosupplementation) for hip osteoarthritis. *Issues Emerg Health Technol*. 2007;1-4.
41. Dagenais S, Garbedian S, Wai EK. Systematic Review of the Prevalence of Radiographic Primary Hip Osteoarthritis. *Clin Orthop Relat Res*. 2008;
42. Dougados M, Gueguen A, Nguyen M, et al. Radiographic features predictive of radiographic progression of hip osteoarthritis. *Rev Rhum Engl Ed*. 1997;64:795-803.
43. Dougados M, Gueguen A, Nguyen M, et al. Radiological progression of hip osteoarthritis: definition, risk factors and correlations with clinical status. *Ann Rheum Dis*. 1996;55:356-362.
44. Dougados M, Gueguen A, Nguyen M, et al. Requirement for total hip arthroplasty: an outcome measure of hip osteoarthritis? *J Rheumatol*. 1999;26:855-861.
45. Dougados M, Villers C, Amor B. Sensitivity to change of various roentgenological severity scoring systems for osteoarthritis of the hip. *Rev Rhum Engl Ed*. 1995;62:169-173.
46. Ehrich EW, Davies GM, Watson DJ, Bolognese JA, Seidenberg BC, Bellamy N. Minimal perceptible clinical improvement with the Western Ontario and McMaster Universities osteoarthritis index questionnaire and global assessments in patients with osteoarthritis. *J Rheumatol*. 2000;27:2635-2641.
47. Ellison JB, Rose SJ, Sahrman SA. Patterns of hip rotation range of motion: a comparison between healthy subjects and patients with low back pain. *Phys Ther*. 1990;70:537-541.
48. Enright PL. The six-minute walk test. *Respir Care*. 2003;48:783-785.
49. Ezoe M, Naito M, Inoue T. The prevalence of acetabular retroversion among various disorders of the hip. *J Bone Joint Surg Am*. 2006;88:372-379.
50. Fagerson T. *The Hip Handbook*. Philadelphia, PA.: W.B. Saunders Company; 1987.
51. Farrar JT, Young JP, Jr., LaMoreaux L, Werth JL, Poole RM. Clinical importance of changes in chronic pain intensity measured on an 11-point numerical pain rating scale. *Pain*. 2001;94:149-158.
52. Felson DT, Lawrence RC, Dieppe PA, et al. Osteoarthritis: new insights. Part 1: the disease and its risk factors. *Ann Intern Med*. 2000;133:635-646.

53. Fife RS. Imaging, arthroscopy, and markers in osteoarthritis. *Curr Opin Rheumatol*. 1992;4:560-565.
54. Fitzgerald RH, Jr. Acetabular labrum tears. Diagnosis and treatment. *Clin Orthop*. 1995;60-68.
55. Foley A, Halbert J, Hewitt T, Crotty M. Does hydrotherapy improve strength and physical function in patients with osteoarthritis--a randomised controlled trial comparing a gym based and a hydrotherapy based strengthening programme. *Ann Rheum Dis*. 2003;62:1162-1167.
56. Fransen M, McConnell S, Bell M. Exercise for osteoarthritis of the hip or knee. *Cochrane Database Syst Rev*. 2003;CD004286.
57. Fransen M, Nairn L, Winstanley J, Lam P, Edmonds J. Physical activity for osteoarthritis management: a randomized controlled clinical trial evaluating hydrotherapy or Tai Chi classes. *Arthritis Rheum*. 2007;57:407-414.
58. Fries JF, Carey C, McShane DJ. Patient education in arthritis: randomized controlled trial of a mail-delivered program. *J Rheumatol*. 1997;24:1378-1383.
59. Gelber AC, Hochberg MC, Mead LA, Wang NY, Wigley FM, Klag MJ. Body mass index in young men and the risk of subsequent knee and hip osteoarthritis. *Am J Med*. 1999;107:542-548.
60. Gelber AC, Hochberg MC, Mead LA, Wang NY, Wigley FM, Klag MJ. Joint injury in young adults and risk for subsequent knee and hip osteoarthritis. *Ann Intern Med*. 2000;133:321-328.
61. Gelberman RH, Cohen MS, Shaw BA, Kasser JR, Griffin PP, Wilkinson RH. The association of femoral retroversion with slipped capital femoral epiphysis. *J Bone Joint Surg Am*. 1986;68:1000-1007.
62. Gershuni DH. Preliminary evaluation and prognosis in Legg-Calve-Perthes disease. *Clin Orthop Relat Res*. 1980;16-22.
63. Gofton JP. Studies in osteoarthritis of the hip. IV. Biomechanics and clinical considerations. *Can Med Assoc J*. 1971;104:1007-1011.
64. Gofton JP, Trueman GE. Studies in osteoarthritis of the hip. II. Osteoarthritis of the hip and leg-length disparity. *Can Med Assoc J*. 1971;104:791-799.
65. Gofton JP, Trueman GE. Unilateral idiopathic osteoarthritis of the hip. *Can Med Assoc J*. 1967;97:1129-1132.
66. Goker B, Doughan AM, Schnitzer TJ, Block JA. Quantification of progressive joint space narrowing in osteoarthritis of the hip: longitudinal analysis of the contralateral hip after total hip arthroplasty. *Arthritis Rheum*. 2000;43:988-994.
67. Goldkind L, Simon LS. Patients, their doctors, nonsteroidal anti-inflammatory drugs and the perception of risk. *Arthritis Res Ther*. 2006;8:105.
68. Golightly YM, Allen KD, Renner JB, Helmick CG, Salazar A, Jordan JM. Relationship of limb length inequality with radiographic knee and hip osteoarthritis. *Osteoarthritis Cartilage*. 2007;15:824-829.
69. Gossec L, Hawker G, Davis AM, et al. OMERACT/OARSI initiative to define states of severity and indication for joint replacement in hip and knee osteoarthritis. *J Rheumatol*. 2007;34:1432-1435.
70. Gossec L, Tubach F, Baron G, Ravaud P, Logeart I, Dougados M. Predictive factors of total hip replacement due to primary osteoarthritis: a prospective 2 year study of 505 patients. *Ann Rheum Dis*. 2005;64:1028-1032.

71. Green J, McKenna F, Redfern EJ, Chamberlain MA. Home exercises are as effective as outpatient hydrotherapy for osteoarthritis of the hip. *Br J Rheumatol*. 1993;32:812-815.
72. Gregory JS, Waarsing JH, Day J, et al. Early identification of radiographic osteoarthritis of the hip using an active shape model to quantify changes in bone morphometric features: can hip shape tell us anything about the progression of osteoarthritis? *Arthritis Rheum*. 2007;56:3634-3643.
73. Grotle M, Hagen KB, Natvig B, Dahl FA, Kvien TK. Obesity and osteoarthritis in knee, hip and/or hand: an epidemiological study in the general population with 10 years follow-up. *BMC Musculoskelet Disord*. 2008;9:132.
74. Gunther KP, Puhl W, Brenner H, Sturmer T. [Clinical epidemiology of hip and knee joint arthroses: an overview of the results of the "Ulm Osteoarthrosis Study"]. *Z Rheumatol*. 2002;61:244-249.
75. Guyatt G, Sackett D, Sinclair J, Hayward R, Cook D, Cook R. Users' guides to the medical literature. IX. A method for grading health care recommendations. Evidence-Based Medicine Working Group. *JAMA*. 1995;274:1800-1804.
76. Harding L, Barbe M, Shepard K, et al. Posterior-anterior glide of the femoral head in the acetabulum: a cadaver study. *J Orthop Sports Phys Ther*. 2003;33:118-125.
77. Harris WH. Traumatic arthritis of the hip after dislocation and acetabular fractures: treatment by mold arthroplasty. An end-result study using a new method of result evaluation. *J Bone Joint Surg Am*. 1969;51:737-755.
78. Harris-Hayes M, Wendl PM, Sahrman SA, Van Dillen LR. Does stabilization of the tibiofemoral joint affect passive prone hip rotation range of motion measures in unimpaired individuals? A preliminary report. *Physiother Theory Pract*. 2007;23:315-323.
79. Hernandez-Molina G, Reichenbach S, Zhang B, Lavalley M, Felson DT. Effect of therapeutic exercise for hip osteoarthritis pain: Results of a meta-analysis. *Arthritis Rheum*. 2008;59:1221-1228.
80. Hinman RS, Heywood SE, Day AR. Aquatic physical therapy for hip and knee osteoarthritis: results of a single-blind randomized controlled trial. *Phys Ther*. 2007;87:32-43.
81. Hoaglund FT, Steinbach LS. Primary osteoarthritis of the hip: etiology and epidemiology. *J Am Acad Orthop Surg*. 2001;9:320-327.
82. Hoeksma HL, Dekker J, Ronday HK, et al. Comparison of manual therapy and exercise therapy in osteoarthritis of the hip: a randomized clinical trial. *Arthritis Rheum*. 2004;51:722-729.
83. Holm I, Bolstad B, Lutken T, Ervik A, Rokkum M, Steen H. Reliability of goniometric measurements and visual estimates of hip ROM in patients with osteoarthrosis. *Physiother Res Int*. 2000;5:241-248.
84. Holmberg S, Stiernstrom EL, Thelin A, Svardsudd K. Musculoskeletal symptoms among farmers and non-farmers: a population-based study. *Int J Occup Environ Health*. 2002;8:339-345.
85. Hopman-Rock M, Westhoff MH. The effects of a health educational and exercise program for older adults with osteoarthritis for the hip or knee. *J Rheumatol*. 2000;27:1947-1954.

86. Hugenberg ST, Brandt KD, Cole CA. Effect of sodium salicylate, aspirin, and ibuprofen on enzymes required by the chondrocyte for synthesis of chondroitin sulfate. *J Rheumatol*. 1993;20:2128-2133.
87. Hughes SL, Seymour RB, Campbell R, Pollak N, Huber G, Sharma L. Impact of the fit and strong intervention on older adults with osteoarthritis. *Gerontologist*. 2004;44:217-228.
88. Hughes SL, Seymour RB, Campbell RT, et al. Long-term impact of Fit and Strong! on older adults with osteoarthritis. *Gerontologist*. 2006;46:801-814.
89. Ingvarsson T, Hagglund G, Lohmander LS. Prevalence of hip osteoarthritis in Iceland. *Ann Rheum Dis*. 1999;58:201-207.
90. Inoue K, Wicart P, Kawasaki T, et al. Prevalence of hip osteoarthritis and acetabular dysplasia in french and japanese adults. *Rheumatology (Oxford)*. 2000;39:745-748.
91. *International. (The) ICD-10*. . Geneva, Switzerland: World Health Organization; 2005.
92. *International Classification of Functioning DaHI*. World Health Organization. Geneva, Switzerland: 2001.
93. Ippolito E, Tudisco C, Farsetti P. Long-term prognosis of Legg-Calve-Perthes disease developing during adolescence. *J Pediatr Orthop*. 1985;5:652-656.
94. Jacobsen S. Adult hip dysplasia and osteoarthritis. Studies in radiology and clinical epidemiology. *Acta Orthop Suppl*. 2006;77:1-37.
95. Jacobsen S, Romer L, Soballe K. Degeneration in dysplastic hips. A computer tomography study. *Skeletal Radiol*. 2005;34:778-784.
96. Jacobsen S, Sonne-Holm S. Hip dysplasia: a significant risk factor for the development of hip osteoarthritis. A cross-sectional survey. *Rheumatology (Oxford)*. 2005;44:211-218.
97. Jacobsen S, Sonne-Holm S, Soballe K, Gebuhr P, Lund B. Factors influencing hip joint space in asymptomatic subjects. A survey of 4151 subjects of the Copenhagen City Heart Study: the Osteoarthritis Substudy. *Osteoarthritis Cartilage*. 2004;12:698-703.
98. Jarvholm B, Lundstrom R, Malchau H, Rehn B, Vingard E. Osteoarthritis in the hip and whole-body vibration in heavy vehicles. *Int Arch Occup Environ Health*. 2004;77:424-426.
99. Jordan JM, Linder GF, Renner JB, Fryer JG. The impact of arthritis in rural populations. *Arthritis Care Res*. 1995;8:242-250.
100. Karachalios T, Karantanas AH, Malizos K. Hip osteoarthritis: what the radiologist wants to know. *Eur J Radiol*. 2007;63:36-48.
101. Kellgren J, Lawrence J. Radiological assesment of osteoarthritis. *Ann Rheum Dis*. 1957;16:494-501.
102. Kellgren JH, Lawrence JS, Bier F. Genetic Factors In Generalized Osteo-Arthrosis. *Ann Rheum Dis*. 1963;22:237-255.
103. Kendall F, McCreary E, Provance P, Rodgers M, Romani W. *Muscles, Testing and Function, with Posture and Pain*. 5th Edition. Philadelphia, PA: Lippincott Williams & Wilkins; 2005.
104. Kennedy DM, Stratford PW, Wessel J, Gollish JD, Penney D. Assessing stability and change of four performance measures: a longitudinal study evaluating outcome following total hip and knee arthroplasty. *BMC Musculoskelet Disord*. 2005;6:3.

105. Kim YJ, Jaramillo D, Millis MB, Gray ML, Burstein D. Assessment of early osteoarthritis in hip dysplasia with delayed gadolinium-enhanced magnetic resonance imaging of cartilage. *J Bone Joint Surg Am.* 2003;85-A:1987-1992.
106. Klassbo M, Larsson G, Harms-Ringdahl K. Promising outcome of a hip school for patients with hip dysfunction. *Arthritis Rheum.* 2003;49:321-327.
107. Lambert RG, Hutchings EJ, Grace MG, Jhangri GS, Conner-Spady B, Maksymowych WP. Steroid injection for osteoarthritis of the hip: a randomized, double-blind, placebo-controlled trial. *Arthritis Rheum.* 2007;56:2278-2287.
108. Lane NE, Lian K, Nevitt MC, et al. Frizzled-related protein variants are risk factors for hip osteoarthritis. *Arthritis Rheum.* 2006;54:1246-1254.
109. Lane NE, Michel B, Bjorkengren A, et al. The risk of osteoarthritis with running and aging: a 5-year longitudinal study. *J Rheumatol.* 1993;20:461-468.
110. Lane NE, Nevitt MC, Cooper C, Pressman A, Gore R, Hochberg M. Acetabular dysplasia and osteoarthritis of the hip in elderly white women. *Ann Rheum Dis.* 1997;56:627-630.
111. Lanyon P, Muir K, Doherty S, Doherty M. Assessment of a genetic contribution to osteoarthritis of the hip: sibling study. *BMJ.* 2000;321:1179-1183.
112. Laslett M, Aprill CN, McDonald B. Provocation sacroiliac joint tests have validity in the diagnosis of sacroiliac joint pain. *Arch Phys Med Rehabil.* 2006;87:874; author reply 874-875.
113. Ledingham J, Dawson S, Preston B, Milligan G, Doherty M. Radiographic progression of hospital referred osteoarthritis of the hip. *Ann Rheum Dis.* 1993;52:263-267.
114. Leibold MR, Huijbregts PA, Jensen R. Concurrent criterion-related validity of physical examination tests for hip labral lesions: a systematic review. *J Man Manip Ther.* 2008;16:E24-41.
115. Lian K, Zmuda JM, Nevitt MC, et al. Type I collagen alpha1 Sp1 transcription factor binding site polymorphism is associated with reduced risk of hip osteoarthritis defined by severe joint space narrowing in elderly women. *Arthritis Rheum.* 2005;52:1431-1436.
116. Lievens A, Bierma-Zeinstra S, Verhagen A, Verhaar J, Koes B. Influence of work on the development of osteoarthritis of the hip: a systematic review. *J Rheumatol.* 2001;28:2520-2528.
117. Lievens AM, Bierma-Zeinstra SM, Verhagen AP, Bernsen RM, Verhaar JA, Koes BW. Influence of sporting activities on the development of osteoarthritis of the hip: a systematic review. *Arthritis Rheum.* 2003;49:228-236.
118. Lievens AM, Bierma-Zeinstra SM, Verhagen AP, van Baar ME, Verhaar JA, Koes BW. Influence of obesity on the development of osteoarthritis of the hip: a systematic review. *Rheumatology (Oxford).* 2002;41:1155-1162.
119. Lievens AM, Bierma-Zeinstra SM, Verhagen AP, Verhaar JA, Koes BW. Prognostic factors of progress of hip osteoarthritis: a systematic review. *Arthritis Rheum.* 2002;47:556-562.
120. Lievens AM, Koes BW, Verhaar JA, Bohnen AM, Bierma-Zeinstra SM. Prognosis of hip pain in general practice: A prospective followup study. *Arthritis Rheum.* 2007;57:1368-1374.
121. Lincoln TL, Suen PW. Common rotational variations in children. *J Am Acad Orthop Surg.* 2003;11:312-320.

122. Lloyd-Roberts GC. The role of capsular changes in osteoarthritis of the hip joint. *J Bone Joint Surg Br.* 1953;35-B:627-642.
123. Loughlin J. The genetic epidemiology of human primary osteoarthritis: current status. *Expert Rev Mol Med.* 2005;7:1-12.
124. Loughlin J, Dowling B, Chapman K, et al. Functional variants within the secreted frizzled-related protein 3 gene are associated with hip osteoarthritis in females. *Proc Natl Acad Sci U S A.* 2004;101:9757-9762.
125. MacDonald CW, Whitman JM, Cleland JA, Smith M, Hoeksma HL. Clinical outcomes following manual physical therapy and exercise for hip osteoarthritis: A case series. *J Orthop Sports Phys Ther.* 2006;36:588-599.
126. MacGregor AJ, Antoniades L, Matson M, Andrew T, Spector TD. The genetic contribution to radiographic hip osteoarthritis in women: results of a classic twin study. *Arthritis Rheum.* 2000;43:2410-2416.
127. Maetzel A, Makela M, Hawker G, Bombardier C. Osteoarthritis of the hip and knee and mechanical occupational exposure--a systematic overview of the evidence. *J Rheumatol.* 1997;24:1599-1607.
128. Mannoni A, Briganti MP, Di Bari M, et al. Epidemiological profile of symptomatic osteoarthritis in older adults: a population based study in Dicomano, Italy. *Ann Rheum Dis.* 2003;62:576-578.
129. Marks R, Allegrante JP. Body mass indices in patients with disabling hip osteoarthritis. *Arthritis Res.* 2002;4:112-116.
130. Mathias S, Nayak US, Isaacs B. Balance in elderly patients: the "get-up and go" test. *Arch Phys Med Rehabil.* 1986;67:387-389.
131. Matles AL. Motion of the hip joint. *Bull Hosp Joint Dis.* 1975;36:170-176.
132. McAlindon T. Why are clinical trials of glucosamine no longer uniformly positive? *Rheum Dis Clin North Am.* 2003;29:789-801.
133. McAlindon T, Formica M, LaValley M, Lehmer M, Kabbara K. Effectiveness of glucosamine for symptoms of knee osteoarthritis: results from an internet-based randomized double-blind controlled trial. *Am J Med.* 2004;117:643-649.
134. McAlindon TE, LaValley MP, Gulin JP, Felson DT. Glucosamine and chondroitin for treatment of osteoarthritis: a systematic quality assessment and meta-analysis. *Jama.* 2000;283:1469-1475.
135. McGibbon CA, Krebs DE, Mann RW. In vivo hip pressures during cane and load-carrying gait. *Arthritis Care Res.* 1997;10:300-307.
136. McGibbon CA, Krebs DE, Scarborough DM. Rehabilitation effects on compensatory gait mechanics in people with arthritis and strength impairment. *Arthritis Rheum.* 2003;49:248-254.
137. Minor MA, Hewett JE, Webel RR, Anderson SK, Kay DR. Efficacy of physical conditioning exercise in patients with rheumatoid arthritis and osteoarthritis. *Arthritis Rheum.* 1989;32:1396-1405.
138. Minor MA, Hewett JE, Webel RR, Dreisinger TE, Kay DR. Exercise tolerance and disease related measures in patients with rheumatoid arthritis and osteoarthritis. *J Rheumatol.* 1988;15:905-911.
139. Mitchell B, McCrory P, Brukner P, O'Donnell J, Colson E, Howells R. Hip joint pathology: clinical presentation and correlation between magnetic resonance arthrography, ultrasound, and arthroscopic findings in 25 consecutive cases. *Clin J Sport Med.* 2003;13:152-156.

140. Murphy WA, Jr., Altman RD. Updated osteoarthritis reference standard. *J Rheumatol Suppl.* 1995;43:56-59.
141. Nagasawa F, Miyake Y, Akazawa H, et al. Predictability of the progress of secondary osteoarthritis after developmental dislocation of the hip, utilizing inferior edge (of the teardrop)--center (of the femoral head) distance. *J Orthop Sci.* 2000;5:10-17.
142. Nahoda J. [Difference of lower limb length in coxarthrosis (author's transl)]. *Acta Chir Orthop Traumatol Cech.* 1977;44:218-220.
143. Nahoda J. [Significance of timely correction of leg length inequality in the prevention of osteoarthroses]. *Beitr Rheumatol.* 1972;18:146-148.
144. Neumann DA. Biomechanical analysis of selected principles of hip joint protection. *Arthritis Care Res.* 1989;2:146-155.
145. Neumann DA, Cook TM. Effect of load and carrying position on the electromyographic activity of the gluteus medius muscle during walking. *Phys Ther.* 1985;65:305-311.
146. Nevitt MC. Definition of hip osteoarthritis for epidemiological studies. *Ann Rheum Dis.* 1996;55:652-655.
147. O'Connor MI. Sex differences in osteoarthritis of the hip and knee. *J Am Acad Orthop Surg.* 2007;15 Suppl 1:S22-25.
148. Oostendorp RA, vanden Heuvel JH, Dekker J, van Baar ME. *Exercise therapy in patients with osteoarthritis of knee or hip: a protocol.* Amersfoort/Utrecht, The Netherlands: NPI/NIVEL; 1998.
149. Palmoski MJ, Brandt KD. Proteoglycan depletion, rather than fibrillation, determines the effects of salicylate and indomethacin on osteoarthritic cartilage. *Arthritis Rheum.* 1985;28:548-553.
150. Pearson JR, Riddell DM. Idiopathic osteo-arthritis of the hip. *Ann Rheum Dis.* 1962;21:31-39.
151. Perera S, Mody SH, Woodman RC, Studenski SA. Meaningful change and responsiveness in common physical performance measures in older adults. *J Am Geriatr Soc.* 2006;54:743-749.
152. Pisters MF, Veenhof C, van Meeteren NL, et al. Long-term effectiveness of exercise therapy in patients with osteoarthritis of the hip or knee: a systematic review. *Arthritis Rheum.* 2007;57:1245-1253.
153. Podsiadlo D, Richardson S. The timed "Up & Go": a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc.* 1991;39:142-148.
154. Pua YH, Wrigley TV, Cowan SM, Bennell KL. Intrarater test-retest reliability of hip range of motion and hip muscle strength measurements in persons with hip osteoarthritis. *Arch Phys Med Rehabil.* 2008;89:1146-1154.
155. Puett DW, Griffin MR. Published trials of nonmedicinal and noninvasive therapies for hip and knee osteoarthritis. *Ann Intern Med.* 1994;121:133-140.
156. Quintana JM, Arostegui I, Escobar A, Azkarate J, Goenaga JI, Lafuente I. Prevalence of knee and hip osteoarthritis and the appropriateness of joint replacement in an older population. *Arch Intern Med.* 2008;168:1576-1584.
157. Qvistgaard E, Christensen R, Torp-Pedersen S, Bliddal H. Intra-articular treatment of hip osteoarthritis: a randomized trial of hyaluronic acid, corticosteroid, and isotonic saline. *Osteoarthritis Cartilage.* 2006;14:163-170.

158. Rasch A, Bystrom AH, Dalen N, Berg HE. Reduced muscle radiological density, cross-sectional area, and strength of major hip and knee muscles in 22 patients with hip osteoarthritis. *Acta Orthop*. 2007;78:505-510.
159. Ravaut P, Giraudeau B, Logeart I, et al. Management of osteoarthritis (OA) with an unsupervised home based exercise programme and/or patient administered assessment tools. A cluster randomised controlled trial with a 2x2 factorial design. *Ann Rheum Dis*. 2004;63:703-708.
160. Reichenbach S, Sterchi R, Scherer M, et al. Meta-analysis: chondroitin for osteoarthritis of the knee or hip. *Ann Intern Med*. 2007;146:580-590.
161. Reijman M, Bierma-Zeinstra SM, Pols HA, Koes BW, Stricker BH, Hazes JM. Is there an association between the use of different types of nonsteroidal antiinflammatory drugs and radiologic progression of osteoarthritis? The Rotterdam Study. *Arthritis Rheum*. 2005;52:3137-3142.
162. Reijman M, Hazes JM, Pols HA, Bernsen RM, Koes BW, Bierma-Zeinstra SM. Role of radiography in predicting progression of osteoarthritis of the hip: prospective cohort study. *BMJ*. 2005;330:1183.
163. Reijman M, Pols HA, Bergink AP, et al. Body mass index associated with onset and progression of osteoarthritis of the knee but not of the hip: the Rotterdam Study. *Ann Rheum Dis*. 2007;66:158-162.
164. Reikeras O, Bjerkreim I, Kolbenstvedt A. Anteversion of the acetabulum and femoral neck in normals and in patients with osteoarthritis of the hip. *Acta Orthop Scand*. 1983;54:18-23.
165. Robinson P, Keenan AM, Conaghan PG. Clinical effectiveness and dose response of image-guided intra-articular corticosteroid injection for hip osteoarthritis. *Rheumatology (Oxford)*. 2007;46:285-291.
166. Roddy E, Zhang W, Doherty M, et al. Evidence-based recommendations for the role of exercise in the management of osteoarthritis of the hip or knee--the MOVE consensus. *Rheumatology (Oxford)*. 2005;44:67-73.
167. Rooks DS, Huang J, Bierbaum BE, et al. Effect of preoperative exercise on measures of functional status in men and women undergoing total hip and knee arthroplasty. *Arthritis Rheum*. 2006;55:700-708.
168. Roos EM, Klassbo M, Lohmander LS. WOMAC osteoarthritis index. Reliability, validity, and responsiveness in patients with arthroscopically assessed osteoarthritis. Western Ontario and MacMaster Universities. *Scand J Rheumatol*. 1999;28:210-215.
169. Salaffi F, Carotti M, Stancati A, Grassi W. Radiographic assessment of osteoarthritis: analysis of disease progression. *Aging Clin Exp Res*. 2003;15:391-404.
170. Singleton MC, LeVeau BF. The hip joint: structure, stability, and stress; a review. *Phys Ther*. 1975;55:957-973.
171. Soderman P, Malchau H. Validity and reliability of Swedish WOMAC osteoarthritis index: a self-administered disease-specific questionnaire (WOMAC) versus generic instruments (SF-36 and NHP). *Acta Orthop Scand*. 2000;71:39-46.
172. Steffen TM, Hacker TA, Mollinger L. Age- and gender-related test performance in community-dwelling elderly people: Six-Minute Walk Test, Berg Balance Scale, Timed Up & Go Test, and gait speeds. *Phys Ther*. 2002;82:128-137.
173. Steultjens MP, Dekker J, van Baar ME, Oostendorp RA, Bijlsma JW. Muscle strength, pain and disability in patients with osteoarthritis. *Clin Rehabil*. 2001;15:331-341.

174. Steultjens MP, Dekker J, van Baar ME, Oostendorp RA, Bijlsma JW. Range of joint motion and disability in patients with osteoarthritis of the knee or hip. *Rheumatology (Oxford)*. 2000;39:955-961.
175. Sturnieks DL, Tiedemann A, Chapman K, Munro B, Murray SM, Lord SR. Physiological risk factors for falls in older people with lower limb arthritis. *J Rheumatol*. 2004;31:2272-2279.
176. Superio-Cabuslay E, Ward MM, Lorig KR. Patient education interventions in osteoarthritis and rheumatoid arthritis: a meta-analytic comparison with nonsteroidal antiinflammatory drug treatment. *Arthritis Care Res*. 1996;9:292-301.
177. Surgeons AAoO. *Osteoarthritis of the Hip, A Compendium of Evidence-based Information and Resources*. 2003.
178. Sutlive TG, Lopez HP, Schnitker DE, et al. Development of a clinical prediction rule for diagnosing hip osteoarthritis in individuals with unilateral hip pain. *J Orthop Sports Phys Ther*. 2008;38:542-550.
179. Tak E, Staats P, Van Hespén A, Hopman-Rock M. The effects of an exercise program for older adults with osteoarthritis of the hip. *J Rheumatol*. 2005;32:1106-1113.
180. Tepper S, Hochberg MC. Factors associated with hip osteoarthritis: data from the First National Health and Nutrition Examination Survey (NHANES-I). *Am J Epidemiol*. 1993;137:1081-1088.
181. Terjesen T, Benum P, Anda S, Svenningsen S. Increased femoral anteversion and osteoarthritis of the hip joint. *Acta Orthop Scand*. 1982;53:571-575.
182. Thelin A. Hip joint arthrosis: an occupational disorder among farmers. *Am J Ind Med*. 1990;18:339-343.
183. Thelin A, Holmberg S. Hip osteoarthritis in a rural male population: A prospective population-based register study. *Am J Ind Med*. 2007;50:604-607.
184. Thelin A, Jansson B, Jacobsson B, Strom H. Coxarthrosis and farm work: a case-referent study. *Am J Ind Med*. 1997;32:497-501.
185. Thelin A, Vingard E, Holmberg S. Osteoarthritis of the hip joint and farm work. *Am J Ind Med*. 2004;45:202-209.
186. Thumboo J, Chew LH, Soh CH. Validation of the Western Ontario and McMaster University osteoarthritis index in Asians with osteoarthritis in Singapore. *Osteoarthritis Cartilage*. 2001;9:440-446.
187. Tokuhashi Y, Matsuzaki H, Uematsu Y, Oda H. Symptoms of thoracolumbar junction disc herniation. *Spine*. 2001;26:E512-518.
188. Tonnis D, Heinecke A. Acetabular and femoral anteversion: relationship with osteoarthritis of the hip. *J Bone Joint Surg Am*. 1999;81:1747-1770.
189. Tonnis D, Heinecke A. [Decreased acetabular anteversion and femur neck antetorsion cause pain and arthrosis. 1: Statistics and clinical sequelae]. *Z Orthop Ihre Grenzgeb*. 1999;137:153-159.
190. Tonnis D, Heinecke A. [Decreased acetabular anteversion and femur neck antetorsion cause pain and arthrosis. 2: Etiology, diagnosis and therapy]. *Z Orthop Ihre Grenzgeb*. 1999;137:160-167.
191. Tonnis D, Heinecke A. Diminished femoral antetorsion syndrome: a cause of pain and osteoarthritis. *J Pediatr Orthop*. 1991;11:419-431.
192. Towheed TE, Hochberg MC. A systematic review of randomized controlled trials of pharmacological therapy in osteoarthritis of the hip. *J Rheumatol*. 1997;24:349-357.

193. van Baar ME, Assendelft WJ, Dekker J, Oostendorp RA, Bijlsma JW. Effectiveness of exercise therapy in patients with osteoarthritis of the hip or knee: a systematic review of randomized clinical trials. *Arthritis Rheum.* 1999;42:1361-1369.
194. van Baar ME, Dekker J, Lemmens JA, Oostendorp RA, Bijlsma JW. Pain and disability in patients with osteoarthritis of hip or knee: the relationship with articular, kinesiological, and psychological characteristics. *J Rheumatol.* 1998;25:125-133.
195. van Baar ME, Dekker J, Oostendorp RA, Bijl D, Voorn TB, Bijlsma JW. Effectiveness of exercise in patients with osteoarthritis of hip or knee: nine months' follow up. *Ann Rheum Dis.* 2001;60:1123-1130.
196. van Baar ME, Dekker J, Oostendorp RA, et al. The effectiveness of exercise therapy in patients with osteoarthritis of the hip or knee: a randomized clinical trial. *J Rheumatol.* 1998;25:2432-2439.
197. van den Bekerom MP, Lamme B, Sermon A, Mulier M. What is the evidence for viscosupplementation in the treatment of patients with hip osteoarthritis? Systematic review of the literature. *Arch Orthop Trauma Surg.* 2008;128:815-823.
198. van den Bekerom MP, Rys B, Mulier M. Viscosupplementation in the hip: evaluation of hyaluronic acid formulations. *Arch Orthop Trauma Surg.* 2008;128:275-280.
199. Van der Esch M, Heijmans M, Dekker J. Factors contributing to possession and use of walking aids among persons with rheumatoid arthritis and osteoarthritis. *Arthritis Rheum.* 2003;49:838-842.
200. van Dijk GM, Veenhof C, Schellevis F, et al. Comorbidity, limitations in activities and pain in patients with osteoarthritis of the hip or knee. *BMC Musculoskeletal Disord.* 2008;9:95.
201. Villanueva I, del Mar Guzman M, Javier Toyos F, Ariza-Ariza R, Navarro F. Relative efficiency and validity properties of a visual analogue vs a categorical scaled version of the Western Ontario and McMaster Universities Osteoarthritis (WOMAC) Index: Spanish versions. *Osteoarthritis Cartilage.* 2004;12:225-231.
202. Vinciguerra C, Gueguen A, Revel M, Heuleu JN, Amor B, Dougados M. Predictors of the need for total hip replacement in patients with osteoarthritis of the hip. *Rev Rhum Engl Ed.* 1995;62:563-570.
203. Vingard E. Overweight predisposes to coxarthrosis. Body-mass index studied in 239 males with hip arthroplasty. *Acta Orthop Scand.* 1991;62:106-109.
204. Vingard E, Alfredsson L, Goldie I, Hogstedt C. Occupation and osteoarthritis of the hip and knee: a register-based cohort study. *Int J Epidemiol.* 1991;20:1025-1031.
205. Vingard E, Alfredsson L, Malchau H. Lifestyle factors and hip arthrosis. A case referent study of body mass index, smoking and hormone therapy in 503 Swedish women. *Acta Orthop Scand.* 1997;68:216-220.
206. Vingard E, Alfredsson L, Malchau H. Osteoarthritis of the hip in women and its relation to physical load at work and in the home. *Ann Rheum Dis.* 1997;56:293-298.
207. Vingard E, Hogstedt C, Alfredsson L, Fellenius E, Goldie I, Koster M. Coxarthrosis and physical work load. *Scand J Work Environ Health.* 1991;17:104-109.
208. Vlad SC, LaValley MP, McAlindon TE, Felson DT. Glucosamine for pain in osteoarthritis: why do trial results differ? *Arthritis Rheum.* 2007;56:2267-2277.
209. Vossinakis IC, Georgiades G, Kafidas D, Hartofilakidis G. Unilateral hip osteoarthritis: can we predict the outcome of the other hip? *Skeletal Radiol.* 2008;

210. Vossinakis LC, Karnezis LA, Parry K, Learmonth ID. Radiographic associations for "primary" hip osteoarthritis: a retrospective cohort study of 47 patients. *Acta Orthop Scand*. 2001;72:600-608.
211. Walker-Bone K, Palmer KT. Musculoskeletal disorders in farmers and farm workers. *Occup Med (Lond)*. 2002;52:441-450.
212. Weigl M, Angst F, Aeschlimann A, Lehmann S, Stucki G. Predictors for response to rehabilitation in patients with hip or knee osteoarthritis: a comparison of logistic regression models with three different definitions of responder. *Osteoarthritis Cartilage*. 2006;14:641-651.
213. Xu L, Nevitt MC, Zhang Y, Yu W, Alibadi P, Felson DT. [High prevalence of knee, but not hip or hand osteoarthritis in Beijing elders: comparison with data of Caucasian in United States]. *Zhonghua Yi Xue Za Zhi*. 2003;83:1206-1209.
214. Yang KG, Raijmakers NJ, Verbout AJ, Dhert WJ, Saris DB. Validation of the short-form WOMAC function scale for the evaluation of osteoarthritis of the knee. *J Bone Joint Surg Br*. 2007;89:50-56.
215. Youdas JW, Mraz ST, Norstad BJ, Schinke JJ, Hollman JH. Determining meaningful changes in hip abductor muscle strength obtained by handheld dynamometry. *Physiother Theory Pract*. 2008;24:215-220.
216. Yrjonen T. Long-term prognosis of Legg-Calve-Perthes disease: a meta-analysis. *J Pediatr Orthop B*. 1999;8:169-172.
217. Zhang W, Doherty M, Arden N, et al. EULAR evidence based recommendations for the management of hip osteoarthritis: report of a task force of the EULAR Standing Committee for International Clinical Studies Including Therapeutics (ESCISIT). *Ann Rheum Dis*. 2005;64:669-681.
218. Zhang W, Moskowitz RW, Nuki G, et al. OARSI recommendations for the management of hip and knee osteoarthritis, part I: critical appraisal of existing treatment guidelines and systematic review of current research evidence. *Osteoarthritis Cartilage*. 2007;15:981-1000.