


Alternative Special Topics: Innovations in Practice

Independent Study Course 25.3.5

Orthopaedic Management of the Patient with Obesity

Christopher Lavallee, DPT
Denver Physical Therapy
Castle Rock, Colorado



REFERENCES

1. Ogden CL, Carroll MD, Kit BK, Flegal KM. *Prevalence of Obesity in the United States 2009-2010*. <http://www.cdc.gov/nchs/data/databriefs/db82.pdf>: Centers for Disease Control; January 2012.
2. Go AS, Mozaffarian D, Roger VL, et al. Executive summary: heart disease and stroke statistics--2013 update: a report from the American Heart Association. *Circulation*. 2013;127(1):143-152. doi: 10.1161/CIR.0b013e318282ab8f.
3. Khan LK, Sobush K, Keener D, et al. Recommended community strategies and measurements to prevent obesity in the United States. *MMWR*. 2009;58(RR-7):1-26.
4. Geliebter A, Atalayer D, Flancbaum L, Gibson CD. Comparison of body adiposity index (BAI) and BMI with estimations of % body fat in clinically severe obese women. *Obesity (Silver Spring)*. 2013;21(3):493-498. doi: 10.1002/oby.20264.
5. Obesity: preventing and managing the global epidemic. Report of a WHO consultation. *World Health Organ Tech Rep Ser*. 2000;894:i-xii.
6. Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults—The Evidence Report. National Institutes of Health. *Obes Res*. 1998;6 Suppl 2:51S-209S.
7. American Physical Therapy Association House of Delegates. Health Promotion And Wellness By Physical Therapists And Physical Therapist Assistants HOD P06-93-25-50 [Position] Accessed January 5, 2014. http://www.apta.org/uploaded-Files/APTAorg/About_Us/Policies/HOD/Health/HealthPromotion.pdf.pdf
8. American College of Sports Medicine. *ACSM's Exercise Management for Persons with Chronic Diseases and Disabilities*. 2nd ed. Champaign, IL: Human Kinetics; 2003.
9. World Health Organization. Waist Circumference and Waist-Hip Ratio: Report of a WHO Expert Consultation; 2011.
10. Mooney S, J Baecker A, Rundle AG. Comparison of anthropometric and body composition measures as predictors of components of the metabolic syndrome in a clinical setting. *Obes Res Clin Pract*. 2013;7(1):e55-e66. doi: 10.1016/j.orcp.2012.10.004.
11. Haftenberger M, Lahmann P, Panico S, et al. Overweight, obesity and fat distribution in 50- to 64-year-old participants in the European Prospective Investigation into Cancer and Nutrition (EPIC). *Public Health Nutr*. 2002;5(6B):1147-1162.
12. Hu FB. *Obesity Epidemiology*. New York, NY: Oxford University Press; 2008.
13. Krachler B, Volgyi E, Savonen K, Tylavsky FA, Alen M, Cheng S. BMI and an anthropometry-based estimate of fat mass percentage are both valid discriminators of cardiometabolic risk: a comparison with DXA and bioimpedance. *J Obes*. 2013;2013:862514. doi: 10.1155/2013/862514.
14. Sweeting HN. Measurement and definitions of

- obesity in childhood and adolescence: a field guide for the uninitiated. *Nutr J.* 2007;6(32):1-8.
15. Mignardot JB, Olivier I, Promayon E, Nougier V. Obesity impact on the attentional cost for controlling posture. *PLoS One.* 2010;5(12):e14387. doi: 10.1371/journal.pone.0014387.
 16. Mignardot JB, Olivier I, Promayon E, Nougier V. Origins of balance disorders during a daily living movement in obese: can biomechanical factors explain everything? *PLoS One.* 2013;8(4):e60491. doi: 10.1371/journal.pone.0060491.
 17. Ku PX, Abu Osman NA, Yusof A, Wan Abas WA. Biomechanical evaluation of the relationship between postural control and body mass index. *J Biomech.* 2012;45(9):1638-1642. doi: 10.1016/j.jbiomech.2012.03.029.
 18. Greve J, Alonso A, Bordini AC, Camanho, GL. Correlation between body mass index and postural balance. *Clin (Sao Paulo).* 2007;62(6):717-720.
 19. Freitas SM, Wieczorek SA, Marchetti PH, Duarte M. Age-related changes in human postural control of prolonged standing. *Gait Posture.* 2005;22(4):322-330.
 20. Lepers D, Bigard AX, Diard JP, Gouteyron JF, Guezennec C. Posture control after exercise. *Eur J Appl Physiol Occup Physiol.* 1997;76:55-61.
 21. LaFortuna CL, Agosti F, Galli C, Busti C, Lazzer S, Sartorio A. The energetic and cardiovascular response to treadmill walking and cycle ergometer exercise in obese women. *Eur J Appl Physiol Occup Physiol.* 2008;103(6):707-717. doi: 10.1007/s00421-008-0758-y.
 22. Singh D, Park W, Levy MS, Jung ES. The effects of obesity and standing time on postural sway during prolonged quiet standing. *Ergonomics.* 2009; 52(8):977-986. doi: 10.1080/00140130902777636.
 23. Blaszczyk JW, Cieslinska-Swider J, Plewa M, Zahorska-Markiewicz B, Markiewicz A. Effects of excessive body weight on postural control. *J Biomech.* 2009;42(9):1295-1300. doi: 10.1016/j.jbiomech.2009.03.006.
 24. Wearing SC, Hennig EM, Byrne NM, Steele JR, Hills AP. The biomechanics of restricted movement in adult obesity. *Obes Rev.* 2006;2006;7:13-24.
 25. Gilleard W, Smith T. Effect of obesity on posture and hip joint moments during a standing task, and trunk forward flexion motion. *Int J Obes (Lond).* 2007;31(2):267-271.
 26. Hue O, Simoneau M, Marcotte J, et al. Body weight is a strong predictor of postural stability. *Gait Posture.* 2007;26(1):32-38.
 27. Shumway-Cook A, Woollacott MH. *Motor Control: Translating Research Into Clinical Practice.* Vol 3: Lippincott Williams & Wilkins; 2007.
 28. Buschbacher RM. Body mass index effect on common nerve conduction study measurements. *Muscle Nerve.* 1998;21(11):1398-1404.
 29. Bensmaia SJ, Leung YY, Hsaio S, Johnson K. Vibratory adaptation of cutaneous mechanoreceptor afferents. *Neurophysiol.* 2005;94(5):3023-3036.
 30. Nantel J, Mathieu ME, Prince F. Physical activity and obesity: biomechanical and physiological key concepts. *J Obes.* 2011;2011:650230. doi: 10.1155/2011/650230.
 31. Colne P, Frelut ML, Peres G, Thoumie P. Postural control in obese adolescents assessed by limits of stability and gait initiation. *Gait Posture.* 2008;28(1):164-169. doi: 10.1016/j.gaitpost.2007.11.006.
 32. Browning RC, McGowan CP, Kram R. Obesity does not increase external mechanical work per kilogram body mass during walking. *J Biomech.* 2009;42(14):2273-2278. doi: 10.1016/j.jbiomech.2009.06.046.
 33. Runhaar J, Koes BW, Clockaerts S, Bierma-Zeinstra SM. A systematic review on changed biomechanics of lower extremities in obese individuals: a possible role in development of osteoarthritis. *Obes Rev.* 2011;12(12):1071-1082. doi: 10.1111/j.1467-789X.2011.00916.x.
 34. Yamakawa K, Tsai CK, Haig AJ, Miner JA, Harris MJ. Relationship between ambulation and obesity in older persons with and without low back pain. *Int J Obes Relat Metab Disord.* 2004;28(1):137-143.
 35. Sheehan K, Gormley J. Gait and increased body weight (potential implications for musculoskeletal disease). *Phys Ther Rev.* 2012;17(2):91-98.
 36. Hortobagyi T, Herring C, Pories WJ, Rider P, Devita P. Massive weight loss-induced mechanical plasticity in obese gait. *J Appl Physiol.* 2011;111(5):1391-1399. doi: 10.1152/japplphysiol.00291.2011.
 37. De Vita P, Hortobagyi T. Obesity is not associated with increased knee joint torque and power during walking. *J Biomech.* 2003;36:1355-1362.
 38. Browning RC, Kram R. Effects of obesity on biomechanics of walking at different speeds. *Med Sci Sports Exerc.* 2007;39:1632-1641
 39. Ohtake PJ. The impact of obesity on walking: implications for fitness assessment and exercise prescription (select presentations from 2008 Combined Sections Meeting). *Cardiopulm Phys Ther J.* 2008;19(2):52-53.

40. Westlake CG, Milner CE, Zhang S, Fitzhugh CE. Do thigh circumference and mass changes alter knee biomechanics during walking? *Gait Posture*. 2013;37(3):359-362. doi: 10.1016/j.gaitpost.2012.07.031.
41. Freedman Silvernail J, Milner CE, Thompson D, Zhang S, Zhao X. The influence of body mass index and velocity on knee biomechanics during walking. *Gait Posture*. 2013;37(4):575-579. doi: 10.1016/j.gaitpost.2012.09.016.
42. Ling C, Kelechi T, Mueller M, Brotherton S, Smith S. Gait and Function in Class III Obesity. *J Obes*. 2012;2012:257468. doi: 10.1155/2012/257468.
43. Wearing SC, Hennig EM, Byrne NM, Steele JR, Hills AP. The impact of childhood obesity on musculoskeletal form. *Obes Rev*. 2006;7:209-218.
44. Hulens M, Vansant G, Claessens AL, Lysens R, Muls E. Predictors of 6-minute walk test results in lean, obese and morbidly obese women. *Scand J Med Sci Sports*. 2003;13(2):98-105.
45. Vincent HK, Raiser SN, Vincent KR. The aging musculoskeletal system and obesity-related considerations with exercise. *Ageing Res Rev*. 2012;11(3):361-373. doi: 10.1016/j.arr.2012.03.002.
46. Janke EA, Collins A, Kozak AT. Overview of the relationship between pain and obesity: What do we know? Where do we go next? *J Rehabil Res Dev*. 2007;44(2):245-262.
47. Nijs J, Van Houdenhove B, Oostendorp RA. Recognition of central sensitization in patients with musculoskeletal pain: application of pain neurophysiology in manual therapy practice. *Man Ther*. 2010;15(2):135-141. doi: 10.1016/j.math.2009.12.001.
48. Vasudevan SV, Potts EE, Mehrotra C. Pain management in arthritis: evidence-based guidelines. *WMJ*. 2003;102(7):14-18.
49. Waddell G, Newton M, Henderson I, Somerville D, Main CJ. A Fear-Avoidance Beliefs Questionnaire (FABQ) and the role of fear-avoidance beliefs in chronic low back pain and disability. *Pain*. 1993;52(2):157-168.
50. Hart DL, Werneke MW, George SZ, et al. Screening for elevated levels of fear-avoidance beliefs regarding work or physical activities in people receiving outpatient therapy. *Phys Ther*. 2009;89(8):770-785. doi: 10.2522/pptj.20080227.
51. Nijs J, Paul van Wilgen C, Van Oosterwijck J, van Ittersum M, Meeus M. How to explain central sensitization to patients with 'unexplained' chronic musculoskeletal pain: practice guidelines. *Man Ther*. 2011;16(5):413-418. doi: 10.1016/j.math.2011.04.005.
52. Leboeuf-Yde C, Kyvik K, Bruun N. Low back pain and life-style. Part II—Obesity. Information from a population-based sample of 29,424 twin subjects. *Spine (Phila Pa 1976)*. 1999;24(8):779-783.
53. Rihn JA, Radcliff K, Hilibrand AS, et al. Does obesity affect outcomes of treatment for lumbar stenosis and degenerative spondylolisthesis? Analysis of the Spine Patient Outcomes Research Trial (SPORT). *Spine (Phila Pa 1976)*. 2012;37(23):1933-1946. doi: 10.1097/BRS.0b013e31825e21b2.
54. Liuke M, Solovieva S, Lamminen A, et al. Disc degeneration of the lumbar spine in relation to overweight. *Int J Obes (Lond)*. 2005;29(8):903-908.
55. Toda Y, Segal N, Toda T, Morimoto T, Ogawa R. Lean body mass and body fat distribution in participants with chronic low back pain. *Arch Intern Med*. 2000;160(21):3265-3269.
56. Shiri R, Solovieva S, Husgafvel-Pursiainen K, et al. The association between obesity and the prevalence of low back pain in young adults: the Cardiovascular Risk in Young Finns Study. *Am J Epidemiol*. 2008;167(9):1110-1119. doi: 10.1093/aje/kwn007.
57. Vismara L, Menegoni F, Zaina F, Galli M, Negrini S, Capodaglio P. Effect of obesity and low back pain on spinal mobility: a cross sectional study in women. *J Neuroeng Rehabil*. 2010;7:3. doi: 10.1186/1743-0003-7-3.
58. Cimolin V, Vismara L, Galli M, Cau N, Capodaglio P. *Omega-3 Fatty Acids in Brain and Neurological Health*. Watson RR, ed. Boston, MA: Elsevier; 2014:417-427.
59. Fanuele JC, Abdu WA Hanscom B, Weinstein JN. Association between obesity and functional status in patients with spine disease. *Spine (Phila Pa 1976)*. 2002;27(3):306-312.
60. Mangwani J, Giles C, Mullins M, Salih T, Natali C. Obesity and recovery from low back pain: a prospective study to investigate the effect of body mass index on recovery from low back pain. *Ann R Coll Surg Engl*. 2010;92(1):23-26. doi: 10.1308/003588410X12518836438967.
61. Cuesta-Vargas AI, Gonzalez-Sanchez M. Obesity effect on a multimodal physiotherapy program for low back pain sufferers: patient reported outcome. *J Occup Med Toxicol*. 2013;8(1):13. doi: 10.1186/1745-6673-8-13.
62. Burns SA, Mintken PE, Austin GP, Cleland J. Short-term response of hip mobilizations and exercise in individuals with chronic low back pain: a case series. *J Man Manip Ther*. 2011;19(2):100-107. doi: 10.1179/2042618610Y.0000000007.

63. Rajgopal V, Bourne RB, Chesworth BM, MacDonald SJ, McCalden RW, Rorabeck CH. The impact of morbid obesity on patient outcomes after total knee arthroplasty. *J Arthroplasty*. 2008;23(6):795-800. doi: 10.1016/j.arth.2007.08.005.
64. Brosseau L, Wells GA, Tugwell P, et al. Ottawa Panel evidence-based clinical practice guidelines for the management of osteoarthritis in adults who are obese or overweight. *Phys Ther*. 2011;91(6):843-861. doi: 10.2522/ptj.20100104.
65. Berry PA, Wluka AE, Davies-Tuck ML, et al. The relationship between body composition and structural changes at the knee. *Rheumatology (Oxford)*. 2010;49(12):2362-2369. doi: 10.1093/rheumatology/keq255.
66. Cimmino MA, Parodi M. Risk factors for osteoarthritis. *Semin Arthritis Rheum*. 2004;34(6):29-34.
67. Stürmer T, Güntherb K-P, Brennera H. Obesity, overweight and patterns of osteoarthritis: the Ulm Osteoarthritis Study. *J Clin Epidemiol*. 2000;53(3):307-313.
68. Vasudevan SV, Potts EE, Mehrotra C. Pain management in arthritis: evidence based guidelines. *WMJ*. 2003;102(7):14-18.
69. Ottawa Panel Evidence-Based Clinical Practice Guidelines for Therapeutic Exercises and Manual Therapy in the Management of Osteoarthritis. *Phys Ther*. 2005;85:907-971.
70. Slemenda C, Brandt KD, Heilman DK, et al. Quadriceps weakness and osteoarthritis of the knee. *Ann Intern Med*. 1997;127(2):97-104.
71. Oliveria SA, Felson DT, Cirillo PA, Reed JI, Walker AM. Body weight, body mass index, and incident symptomatic osteoarthritis of the hand, hip, and knee. *Epidemiology*. 1999;10(2):161-166.
72. Grotle M, Hagen KB, Natvig B, Dahl FA, Kvien TK. Obesity and osteoarthritis in knee, hip and/or hand: an epidemiological study in the general population with 10 years follow-up. *BMC Musculoskeletal Disord*. 2008;9:132. doi: 10.1186/1471-2474-9-132.
73. Tukker A, Visscher TL, Picavet HS. Overweight and health problems of the lower extremities: osteoarthritis, pain and disability. *Public Health Nutr*. 2009;12(3):359-368. doi: 10.1017/S1368980008002103.
74. Foster NA, Segal NA, Clearfield JS, et al. Central versus lower body obesity distribution and the association with lower limb physical function and disability. *PM R*. 2010;2(12):1119-1126. doi: 10.1016/j.pmrj.2010.09.002.
75. Jiang L, Tian W, Wang Y, et al. Body mass index and susceptibility to knee osteoarthritis: a systematic review and meta-analysis. *Joint Bone Spine*. 2012;79(3):291-297. doi: 10.1016/j.jbspin.2011.05.015.
76. Hall M, Boyer ER, Gillette JC, Mirka GA. Medial knee joint loading during stair ambulation and walking while carrying loads. *Gait Posture*. 2013;37(3):460-462. doi: 10.1016/j.gaitpost.2012.08.008.
77. Workgroup of the American Association of Hip and Knee Surgeons Evidence Based Committee. Obesity and total joint arthroplasty: a literature based review. *J Arthroplasty*. 2013;28(5):714-721. doi: 10.1016/j.arth.2013.02.011.
78. Sharma L LC, Cahue S, Dunlop DD. The mechanism of the effect of obesity in knee osteoarthritis: the mediating role of malalignment. *Arthritis Rheum*. 2000;43:568-575.
79. Jadelis K, Miller ME, Ettinger WH Jr, Messier SP. Strength, balance, and the modifying effects of obesity and knee pain: results from the Observational Arthritis Study in Seniors (oasis). *J Am Geriatr Soc*. 2001;49(7):884-891.
80. Powers CM, Bolgla LA, Callaghan MJ, Collins N, Sheehan FT. Patellofemoral Pain: Proximal, Distal, and Local Factors – 2nd International Research Retreat, August 31-September 2, 2011, Ghent Belgium. *J Orthop Sports Phys Ther*. 2012;42(6):A1-A54. doi: 10.2519/jospt.2012.0301.
81. Anandacoomarasamy A, Caterson I, Sambrook P, Fransen M, March L. The impact of obesity on the musculoskeletal system. *Int J Obes (Lond)*. 2008;32(2):211-222.
82. Tragard BS, Gill NW, Silvernail JL, Teyhen DS, Allison SC. Joint mobilization forces and therapist reliability in subjects with knee osteoarthritis. *J Man Manip Ther*. 2013;21(4):196-206. doi: 10.1179/2042618613Y.0000000033.
83. Lievense AM, Bierma-Zeinstra SMA, Verhagen AP, van Baar ME, Verhaar JA, Koes BW. Influence of obesity on the development of osteoarthritis of the hip: a systematic review. *Rheumatology (Oxford)*. 2002;41(10):1155-1162.
84. Robertsson O, Wingstrand H, Onnerfalt R. Intracapsular pressure and pain in coxarthrosis. *J Arthroplasty*. 1995;10(5):632-635.
85. Villafane JH, Cleland JA, Fernandez-de-Las-Penas C. The effectiveness of a manual therapy and exercise protocol in patients with thumb carpometacarpal osteoarthritis: a randomized controlled trial. *J Orthop Sports Phys Ther*. 2013;43(4):204-213. doi: 10.2519/jospt.2013.4524.

86. Kerkhoffs GM, Servien E, Dunn W, Dahm D, Bramer JA, Haverkamp D. The influence of obesity on the complication rate and outcome of total knee arthroplasty: a meta-analysis and systematic literature review. *J Bone Joint Surg Am.* 2012;94(20):1839-1844. doi: 10.2106/JBJS.K.00820.
87. Meier W, Mizner RL, Marcus R, Dibble L, Peters C, Lastayo PC. Total knee arthroplasty: muscle impairments, functional limitations, and recommended rehabilitation approaches. *J Orthop Sports Phys Ther.* 2008;38(4):246-256. doi: 10.2519/jospt.2008.2715.
88. Vissers D, Hens W, Taeymans J, Baeyens JP, Poortmans J, Van Gaal L. The effect of exercise on visceral adipose tissue in overweight adults: a systematic review and meta-analysis. *PLoS one.* 2013;8(2):e56415. doi: 10.1371/journal.pone.0056415.
89. Paans N, Stevens M, Wagenmakers R, et al. Changes in body weight after total hip arthroplasty: short-term and long-term effects. *Phys Ther.* 2012;92(5):680-687. doi: 10.2522/ptj.20110176.
90. Vaidya R, Carp J, Bartol S, Ouellette N, Lee S, Sethi A. Lumbar spine fusion in obese and morbidly obese patients. *Spine (Phila Pa 1976).* 2009;34(5):495-500. doi: 10.1097/BRS.0b013e318198c5f2.
91. Dijurasovic M, Bratcher KR, Glassman SD, Dimar JR, Carreon LY. The effect of obesity on clinical outcomes after lumbar fusion. *Spine (Phila Pa 1976).* 2008;33(16):1789-1792. doi: 10.1097/BRS.0b013e31817b8f6f.
92. Hills AP, Hennig EM, McDonald M, Bar-Or O. Plantar pressure differences between obese and non-obese adults: a biomechanical analysis. *Int J Obes Relat Metab Disord.* 2001;25(11):1674-1679.
93. Hills AP, Hennig EM, Byrne NM, Steele JR. The biomechanics of adiposity--structural and functional limitations of obesity and implications for movement. *Obes Rev.* 2002;3(1):35-43.
94. Butterworth PA, Landorf KB, Smith SE, Menz HB. The association between body mass index and musculoskeletal foot disorders: a systematic review. *Obes Rev.* 2012;13(7):630-642. doi: 10.1111/j.1467-789X.2012.00996.x.
95. Irving DB, Cook JL, Young MA, Menz HB. Obesity and pronated foot type may increase the risk of chronic plantar heel pain: a matched case-control study. *BMC Musculoskelet Disord.* 2007;8:41.
96. Wearing SC, Hennig EM, Byrne NM, Steele JR, Hills AP. Musculoskeletal disorders associated with obesity: a biomechanical perspective. *Obes Rev.* 2006;7(3):239-250.
97. Riddle DL, Pulisic M, Pidcoe P, Johnson RE. Risk factors for Plantar fasciitis: a matched case-control study. *J Bone Joint Surg.* 2003;85-A(5):872-877.
98. Chen MM, Lear SA, Gao M, Frohlich JJ, Birmingham CL. Intraobserver and interobserver reliability of waist circumference and the waist-to-hip ratio. *Obes Res.* 2001;9(10):651.
99. Drake M, Bittenbender C, Boyles RE. The short-term effects of treating plantar fasciitis with a temporary custom foot orthosis and stretching. *J Orthop Sports Phys Ther.* 2011;41(4):221-231. doi: 10.2519/jospt.2011.3348.
100. Renan-Ordine R, Alburquerque-Sendin F, de Souza DP, Cleland JA, Fernandez-de-Las-Penas C. Effectiveness of myofascial trigger point manual therapy combined with a self-stretching protocol for the management of plantar heel pain: a randomized controlled trial. *J Orthop Sports Phys Ther.* 2011;41(2):43-50. doi: 10.2519/jospt.2011.3504.
101. Cleland JA, Abbott JH, Kidd MO, et al. Manual physical therapy and exercise versus electrophysical agents and exercise in the management of plantar heel pain: a multicenter randomized clinical trial. *J Orthop Sports Phys Ther.* 2009;39(8):573-585. doi: 10.2519/jospt.2009.3036.
102. Alshami AM, Souvlis T, Coppeters MW. A review of plantar heel pain of neural origin: differential diagnosis and management. *Man Ther.* 2008;13(2):103-111.
103. Woodley SJ, Nicholson HD, Livingstone V, et al. Lateral hip pain: findings from magnetic resonance imaging and clinical examination. *J Orthop Sports Phys Ther.* 2008;38(6):313-328. doi: 10.2519/jospt.2008.2685.
104. Segal NA, Felson DT, Torner JC, et al. Greater trochanteric pain syndrome: epidemiology and associated factors. *Arch Phys Med Rehabil.* 2007;88(8):988-992.
105. Williams BS, Cohen SP. Greater trochanteric pain syndrome: a review of anatomy, diagnosis and treatment. *Anesth Analg.* 2009;108(5):1662-1670. doi: 10.1213/ane.0b013e31819d6562.
106. Bewyer DC, Bewyer KJ. Rationale for treatment of hip abductor pain syndrome. *Iowa Orthop J.* 2003;23:57-60.
107. Selkowitz DM, Beneck GJ, Powers CM. Which exercises target the gluteal muscles while minimizing activation of the tensor fascia lata? Electromyographic assessment using fine-

- wire electrodes. *J Orthop Sports Phys Ther.* 2013;43(2):54-64. doi: 10.2519/jospt.2013.4116.
108. Moghtaderi A, Izadi S, Sharafadinzadeh N. An evaluation of gender, body mass index, wrist circumference and wrist ratio as independent risk factors for carpal tunnel syndrome. *Acta Neurol Scand.* 2005;112(6):375-379.
109. Fernandez-de-Las-Penas C, Cleland JA, Plaza-Manzano G, et al. Clinical, physical, and neuro-physiological impairments associated with decreased function in women with carpal tunnel syndrome. *J Orthop Sports Phys Ther.* 2013;43(9):641-649. doi: 10.2519/jospt.2013.4830.
110. De-la-Llave-Rincón AI, Fernández-de-las-Peñas C, Palacios-Ceña D, Cleland JA. Increased forward head posture and restricted cervical range of motion in patients with carpal tunnel syndrome. *J Orthop Sports Phys Ther.* 2009;38(9):658-664. doi: 10.2519/jospt.2009.3058.
111. Bialosky JE, Bishop MD, Price DD, Robinson ME, Vincent KR, George SZ. A randomized sham-controlled trial of a neurodynamic technique in the treatment of carpal tunnel syndrome. *J Orthop Sports Phys Ther.* 2009;39(10):709-723. doi: 10.2519/jospt.2009.3117.
112. Ursini F, Naty S, Grembiale RD. Fibromyalgia and obesity: the hidden link. *Rheumatol Int.* 2011;31(11):1403-1408. doi: 10.1007/s00296-011-1885-z.
113. Brosseau L, Wells GA, Tugwell P, et al. Ottawa Panel evidence-based clinical practice guidelines for strengthening exercises in the management of fibromyalgia: part 2. *Phys Ther.* 2008;88(7):873-886. doi: 10.2522/ptj.20070115.
114. Mork PJ, Vasseljen O, Nilsen TI. Association between physical exercise, body mass index, and risk of fibromyalgia: longitudinal data from the Norwegian Nord-Trøndelag Health Study. *Arthritis Care Res.* 2010;62(5):611-617. doi: 10.1002/acr.20118.
115. Gaida JE, Ashe MC, Bass SL, Cook JL. Is adiposity an under-recognized risk factor for tendinopathy? A systematic review. *Arthritis Rheum.* 2009;61(6):840-849. doi: 10.1002/art.24518.
116. Rechardt M, Shiri R, Karppinen J, Jula A, Heliovaara M, Viikari-Juntura E. Lifestyle and metabolic factors in relation to shoulder pain and rotator cuff tendinitis: a population-based study. *BMC Musculoskelet Disord.* 2010;11:165. doi: 10.1186/1471-2474-11-165.
117. Werner RA, Franzblau A, Gell N, Ulin SS, Armstrong TJ. A longitudinal study of industrial and clerical workers: predictors of upper extremity tendonitis. *J Occup Rehabil.* 2005;15(1):37-46.
118. Wendelboe AM, Hegmann KT, Gren LH, Alder SC, White GL Jr, Lyon JL. Associations between body-mass index and surgery for rotator cuff tendinitis. *J Bone Joint Surg.* 2004;86-A(4):743-747.
119. Greer WJ, Richter HE, Bartolucci AA, Burgio KL. Obesity and pelvic floor disorders: a systematic review. *Obstet Gynecol.* 2008;112(2 Pt 1):341-349. doi: 10.1097/AOG.0b013e31817cfddc.
120. Wasserberg N, Haney M, Petrone P, et al. Morbid obesity adversely impacts pelvic floor function in females seeking attention for weight loss surgery. *Dis Colon Rectum.* 2007;50(12):2096-2103.
121. Ottawa Panel. Ottawa Panel evidence-based clinical practice guidelines for therapeutic exercise in the management of rheumatoid arthritis in adults. *Phys Ther.* 2004;84(10):934-972.
122. Focht BC. Affective responses to 10-minute and 30-minute walks in sedentary, overweight women: Relationships with theory-based correlates of walking for exercise. *Psychol Sport Exerc.* 2013;14(5):759-766.
123. George SZ, Zeppieri G. Physical therapy utilization of graded exposure for patients with low back pain. *J Orthop Sports Phys Ther.* 2009;39(7):496-505. doi: 10.2519/jospt.2009.2983.
124. Rosenbaum M, Leibel R. Effects of weight change on skeletal muscle work efficiency in human subjects. *Am J Physiol.* 2003;R183-R192. doi: 10.1152/ajpregu.00397.2011.
125. American College of Sports Medicine. Appropriate physical activity intervention strategies for weight loss and prevention of weight regain in adults. *Med Sci Sports Exerc.* 2009;41(1):459-471. doi: 10.1249/MSS.0b013e3181949333.
126. Obesity Action Coalition. Understanding Obesity Brochure. 2015. <http://www.obesityaction.org/educational-resources/brochures-and-guides/understanding-obesity-brochure>. Accessed January 18, 2014.
127. Lai P, Leung A, Li A, Zhang M. Three-dimensional motion analysis of obese adults. *Clin Biomech.* 2008;23(1S2-6).
128. De Souza S, Faintuch J, Valezi A, et al. Gait cinematic analysis in morbidly obese patients. *Obes Surg.* 2005;15(9):1238-1242.
129. Spyropoulos P, Pisciotta JC, Pavlou KN, Cairns MA, Simon SR. Biomechanical gait analysis in obese men. *Arch Phys Med Rehabil.* 1991;72:1065-1070.

130. Russell E, Braun B, Hamill J. Does stride length influence metabolic cost and biomechanical risk factors for knee osteoarthritis in women? *Clin Biomech (Bristol, Avon)*. 2010;25(5):438-443. doi: 10.1016/j.clinbiomech.2010.01.016.
131. Ling C, Kelechi T, Mueller M, Brotherton S, Smith S. Gait and function in class III obesity. *J Obes*. 2012;1-8. doi: 10.1155/2012/257468.
132. George S, Wittmer V, Fillingim R, Robinson M. Comparison of graded exercise and graded exposure clinical outcomes for patients with chronic low back pain. *J Orthop Sports Phys Ther*. 2010;40(11):694-704. doi: 10.2519/jospt.2010.3396.
133. Rundell SD, Davenport TE. Patient education based on principles of cognitive behavioral therapy for a patient with persistent low back pain: a case report. *J Orthop Sports Phys Ther*. 2010;40(8):494-501. doi: 10.2519/jospt.2010.3264.
134. Nielsen M, Keefe FJ, Bennell K, Jull GA. Physical therapist-delivered cognitive-behavioral therapy: a qualitative study of physical therapists' perceptions and experiences. *Phys Ther*. 2014;94(2):197-209. doi: 10.2522/ptj.20130047.
135. Sutlive TG, Lopez HP, Schnitker DE, et al. Development of a clinical prediction rule for diagnosing hip osteoarthritis in individuals with unilateral hip pain. *J Orthop Sports Phys Ther*. 2008;38(9):542-550. doi: 10.2519/jospt.2008.2753.

NOTES