


Outcomes in Orthopaedic Physical Therapy Practice

Independent Study Course 26.1.2

Performance-rated Outcome Measures—Framework, Psychometrics, and Uses

Charles Sheets, PT, OCS, SCS, Dip MDT
Department of Physical and Occupational Therapy
Doctor of Physical Therapy Program
Duke University
Durham, North Carolina

Orthopaedic Section, APTA, Inc.
Downloaded from www.orthopteam.org at the Orthopaedic Section on September 26, 2018. For personal use only. No other uses without permission.
Copyright © 2016 Orthopaedic Section, APTA, Inc. All rights reserved.



REFERENCES

1. American Physical Therapy Association. Vision Statement for the Physical Therapy Profession and Guiding Principles to Achieve the Vision. 2015; <http://www.apta.org/Vision/>. Accessed June 31, 2015.
2. Terwee C, Mokkink L, Steultjens M, Dekker J. Performance-based methods for measuring the physical function of patients with osteoarthritis of the hip or knee: a systematic review of measurement properties. *Rheumatology*. 2006;45(7):890-902.
3. Stucki G, Cieza A, Ewert T, Kostanjsek N, Chatterji S, ÜstÜN TB. Application of the International Classification of Functioning, Disability and Health (ICF) in clinical practice. *Disabil Rehabil*. 2002;24(5):281-282.
4. Stratford PW, Kennedy DM, Woodhouse LJ. Performance measures provide assessments of pain and function in people with advanced osteoarthritis of the hip or knee. *Phys Ther*. 2006;86(11):1489-1496.
5. Dobson F, Hinman R, Hall M, Terwee C, Roos EM, Bennell K. Measurement properties of performance-based measures to assess physical function in hip and knee osteoarthritis: a systematic review. *Osteoarthritis Cartilage*. 2012;20(12):1548-1562. doi: 10.1016/j.joca.2012.08.015. Epub 2012 Aug 31.
6. Fitzpatrick R, Davey C, Buxton MJ, Jones DR. Evaluating patient-based outcome measures for use in clinical trials. *Health Technol Assess*. 1998;2(14):i-iv, 1-74.
7. Dawson J, Doll H, Fitzpatrick R, Jenkinson C, Carr AJ. The routine use of patient reported outcome measures in healthcare settings. *BMJ*. 2010;340:c186. doi: 10.1136/bmj.c186.
8. Bellamy N, Kirwan J, Boers M, et al. Recommendations for a core set of outcome measures for future phase III clinical trials in knee, hip, and hand osteoarthritis. Consensus development at OMERACT III. *J Rheumatol*. 1997;24(4):799-802.
9. Guralnik JM, Branch LG, Cummings SR, Curb JD. Physical performance measures in aging research. *J Gerontol*. 1989;44(5):M141-146.
10. Reuben DB, Siu AL. An objective measure of physical function of elderly outpatients. The Physical Performance Test. *J Am Geriatr Soc*. 1990;38(10):1105-1112.
11. Mahurin RK, DeBettignies BH, Pirozzolo FJ. Structured assessment of independent living skills: preliminary report of a performance measure of functional abilities in dementia. *J Gerontol*. 1991;46(2):P58-66.
12. Myers AM, Holliday PJ, Harvey KA, Hutchinson KS. Functional performance measures: are they superior to self-assessments? *J Gerontol*. 1993;48(5):M196-206.
13. Stevens-Lapsley JE, Schenkman ML, Dayton MR. Comparison of self-reported knee injury and osteoarthritis outcome score to performance measures in patients after total knee arthroplasty. *PM R*. 2011;3(6):541-549; quiz 549. doi: 10.1016/j.pmrj.2011.03.002.
14. Mizner RL, Petterson SC, Clements KE, Zeni JA, Jr, Irrgang JJ, Snyder-Mackler L. Measuring functional improvement after total knee arthroplasty requires both performance-based and patient-report assessments: a longitudinal analysis of outcomes. *J Arthroplasty*. 2011;26(5):728-737. doi: 10.1016/j.arth.2010.06.004. Epub 2010 Sep 20.
15. Stratford PW, Kennedy DM, Riddle DL. New study design evaluated the validity of measures to

- assess change after hip or knee arthroplasty. *J Clin Epidemiol.* 2009;62(3):347-352. doi: 10.1016/j.jclinepi.2008.06.008. Epub 2008 Oct 1.
16. Jordan KP, Wilkie R, Muller S, Myers H, Nicholls E, Arthritis Research Campaign National Primary Care Centre. Measurement of change in function and disability in osteoarthritis: current approaches and future challenges. *Curr Opin Rheumatol.* 2009;21(5):525-530. doi: 10.1097/BOR.0b013e32832e45fc.
 17. Wright AA, Hegedus EJ, Baxter GD, Abbott JH. Measurement of function in hip osteoarthritis: developing a standardized approach for physical performance measures. *Physiother Theory Practice.* 2011;27(4):253-262. doi: 10.3109/09593985.2010.491150. Epub 2010 Jul 22.
 18. Keenan AM, Redmond AC, Horton M, Conaghan PG, Tennant A. The Foot Posture Index: Rasch analysis of a novel, foot-specific outcome measure. *Arch Phys Med Rehabil.* 2007;88(1):88-93.
 19. Agresti A. *Analysis of Ordinal Categorical Data.* Vol 656. Hoboken, NJ: John Wiley & Sons; 2010.
 20. Hall AM, Maher CG, Latimer J, Ferreira ML, Costa LO. The patient-specific functional scale is more responsive than the Roland Morris disability questionnaire when activity limitation is low. *Eur Spine J.* 2011;20(1):79-86. doi: 10.1007/s00586-010-1521-8. Epub 2010 Jul 14.
 21. Functional Movement. functional.movement.com. FMS Level 1 with Online Certification. 2015; <http://www.functionalmovement.com/certification/FMSLevel1>. Accessed June 30, 2015.
 22. Cusick A, Vasquez M, Knowles L, Wallen M. Effect of rater training on reliability of Melbourne Assessment of Unilateral Upper Limb Function scores. *Dev Med Child Neurol.* 2005;47(01):39-45.
 23. Jette AM. Using health-related quality of life measures in physical therapy outcomes research. *Phys Ther.* 1993;73(8):528-537.
 24. Hughes PC, Taylor NF, Green RA. Most clinical tests cannot accurately diagnose rotator cuff pathology: a systematic review. *Aust J Physiother.* 2008;54(3):159-170.
 25. Malanga GA, Andrus S, Nadler SF, McLean J. Physical examination of the knee: a review of the original test description and scientific validity of common orthopaedic tests. *Arch Phys Med Rehabil.* 2003;84(4):592-603.
 26. Hoogeboom TJ, van Meeteren NL, Kim RH, Stevens-Lapsley JE, Assassi S. Linear and curvilinear relationship between knee range of motion and physical functioning in people with knee osteoarthritis: a cross-sectional study. *PLoS one.* 2013;8(9):e76173. doi: 10.1371/journal.pone.0076173. eCollection 2013.
 27. Manske RC, Smith BS, Rogers ME, Wyatt FB. Closed kinetic chain (linear) isokinetic testing: relationships to functional testing. *Isokinet Exerc Sci.* 2003;11(3):171-179.
 28. Biering-Sorensen F. Physical measurements as risk indicators for low-back trouble over a one-year period. *Spine.* 1984;9(2):106-119.
 29. Gillespie J, Keenum S. A validity and reliability-analysis of the seated shot put as a test of power. *J Human Movement Studies.* 1987;13(2):97-105.
 30. Swank AM, Kachelman JB, Bibeau W, et al. Prehabilitation before total knee arthroplasty increases strength and function in older adults with severe osteoarthritis. *J Strength Cond Res.* 2011;25(2):318-325. doi: 10.1519/JSC.0b013e318202e431.
 31. Choi S, Trang A, McCartney CJ. Reporting functional outcome after knee arthroplasty and regional anesthesia: a methodological primer. *Reg Anesth Pain Med.* 2013;38(4):340-349. doi: 10.1097/AAP.0b013e318295d973.
 32. Gandhi R, Tsvetkov D, Davey JR, Syed KA, Mahomed NN. Relationship between self-reported and performance-based tests in a hip and knee joint replacement population. *Clin Rheumatol.* 2009;28(3):253-257. doi: 10.1007/s10067-008-1021-y. Epub 2008 Oct 8.
 33. Kennedy D, Stratford PW, Pagura SM, Walsh M, Woodhouse LJ. Comparison of gender and group differences in self-report and physical performance measures in total hip and knee arthroplasty candidates. *J Arthroplasty.* 2002;17(1):70-77.
 34. Parent E, Moffet H. Comparative responsiveness of locomotor tests and questionnaires used to follow early recovery after total knee arthroplasty. *Arch Phys Med Rehabil.* 2002;83(1):70-80.
 35. Shulman KI. Clock-drawing: is it the ideal cognitive screening test? *Int J Geriatr Psychiatry.* 2000;15(6):548-561.
 36. Fitzgerald GK, Axe MJ, Snyder-Mackler L. Proposed practice guidelines for nonoperative anterior cruciate ligament rehabilitation of physically active individuals. *J Orthop Sports Phys Ther.* 2000;30(4):194-203.
 37. Dodds TA, Martin DP, Stolov WC, Deyo RA. A validation of the functional independence measurement and its performance among rehabilitation inpatients. *Arch Phys Med Rehabil.* 1993;74(5):531-536.
 38. Bade MJ, Kittelson JM, Kohrt WM, Stevens-Lapsley JE. Predicting functional performance and range of motion outcomes after total knee arthroplasty.

- Am J Phys Med Rehabil.* 2014;93(7):579-585. doi: 10.1097/PHM.0000000000000065.
39. Studenski S, Perera S, Wallace D, et al. Physical performance measures in the clinical setting. *J Am Geriatr Soc.* 2003;51(3):314-322.
 40. Plisky PJ, Rauh MJ, Kaminski TW, Underwood FB. Star Excursion Balance Test as a predictor of lower extremity injury in high school basketball players. *J Orthop Sports Phys Ther.* 2006;36(12):911-919.
 41. Smith CA, Chimera NJ, Warren M. Association of y balance test reach asymmetry and injury in division I athletes. *Med Sci Sports Exerc.* 2015;47(1):136-141. doi: 10.1249/MSS.0000000000000380.
 42. Garrison M, Westrick R, Johnson MR, Benenson J. Association between the functional movement screen and injury development in college athletes. *Int J Sports Phys Ther.* 2015;10(1):21-28.
 43. Lehr ME, Plisky PJ, Butler RJ, Fink ML, Kiesel KB, Underwood FB. Field-expedient screening and injury risk algorithm categories as predictors of noncontact lower extremity injury. *Scand J Med Sci Sports.* 2013;23(4):e225-e232. doi: 10.1111/sms.12062. Epub 2013 Mar 20.
 44. Adams D, Logerstedt D, Hunter-Giordano A, Axe MJ, Snyder-Mackler L. Current concepts for anterior cruciate ligament reconstruction: a criterion-based rehabilitation progression. *J Orthop Sports Phys Ther.* 2012;42(7):601-614. doi: 10.2519/jospt.2012.3871. Epub 2012 Mar 8.
 45. Creighton DW, Shrier I, Shultz R, Meeuwisse WH, Matheson GO. Return-to-play in sport: a decision-based model. *Clin J Sport Med.* 2010;20(5):379-385. doi: 10.1097/JSM.0b013e3181f3c0fe.
 46. Fitzgerald GK, Axe MJ, Snyder-Mackler L. The efficacy of perturbation training in nonoperative anterior cruciate ligament rehabilitation programs for physically active individuals. *Phys Ther.* 2000;80(2):128-140.
 47. Fitzgerald G, Axe M, Snyder-Mackler L. A decision-making scheme for returning patients to high-level activity with nonoperative treatment after anterior cruciate ligament rupture. *Knee Surg Sports Traumatol Arthrosc.* 2000;8(2):76-82.
 48. Shultz R, Bido J, Shrier I, Meeuwisse WH, Garza D, Matheson GO. Team clinician variability in return-to-play decisions. *Clin J Sport Med.* 2013;23(6):456-461. doi: 10.1097/JSM.0b013e318295bb17.
 49. Mayer SW, Queen RM, Taylor D, et al. Functional testing differences in anterior cruciate ligament reconstruction patients released versus not released to return to sport. *Am J Sports Med.* 2015;43(7):1648-1655. doi: 10.1177/0363546515578249. Epub 2015 Apr 13.
 50. Hegedus EJ, Cook CE. Return to play and physical performance tests: evidence-based, rough guess or charade? *Br J Sports Med.* 2015;49(20):1288-1289. doi: 10.1136/bjsports-2015-094796. Epub 2015 Jun 9.
 51. Haines S, Baker T, Donaldson M. Development of a physical performance assessment checklist for athletes who sustained a lower extremity injury in preparation for return to sport: a DELPHI study. *Int J Sports Phys Ther.* 2013;8(1):44-53.
 52. Tegner Y, Lysholm J, Lysholm M, Gillquist J. A performance test to monitor rehabilitation and evaluate anterior cruciate ligament injuries. *Am J Sports Med.* 1986;14(2):156-159.
 53. Myer GD, Schmitt LC, Brent JL, et al. Utilization of modified NFL combine testing to identify functional deficits in athletes following ACL reconstruction. *J Orthop Sports Phys Ther.* 2011;41(6):377-387. doi: 10.2519/jospt.2011.3547. Epub 2011 Feb 2.
 54. Wing Chiu TT, Hung Law EY, Fai Chiu TH. Performance of the craniocervical flexion test in subjects with and without chronic neck pain. *J Orthop Sports Phys Ther.* 2005;35(9):567-571.
 55. Curb JD, Ceria-Ulep CD, Rodriguez BL, et al. Performance-based measures of physical function for high-function populations. *J Am Geriatr Soc.* 2006;54(5):737-742.
 56. Brox JJ, Storheim K, Holm I, Friis A, Reikeras O. Disability, pain, psychological factors and physical performance in healthy controls, patients with sub-acute and chronic low back pain: a case-control study. *J Rehabil Med.* 2005;37(2):95-99.
 57. Falsone SA, Gross MT, Guskiewicz KM, Schneider RA. One-arm hop test: reliability and effects of arm dominance. *J Orthop Sports Phys Ther.* 2002;32(3):98-103.
 58. Hopkins WG. Measures of reliability in sports medicine and science. *Sports Med.* 2000;30(1):1-15.
 59. Valovich TC, Perrin DH, Gansneder BM. Repeat administration elicits a practice effect with the Balance Error Scoring System but not with the Standardized Assessment of Concussion in high school athletes. *J Athl Train.* 2003;38(1):51-56.
 60. McLeod TCV, Perrin DH, Guskiewicz KM, Shultz SJ, Diamond R, Gansneder BM. Serial administration of clinical concussion assessments and learning effects in healthy young athletes. *Clin J Sport Med.* 2004;14(5):287-295.
 61. Smeets RJ, Hijdra HJ, Kester AD, Hitters MW, Knottnerus JA. The usability of six physical performance tasks in a rehabilitation population with chronic low back pain. *Clin Rehabil.* 2006;20(11):989-997.

62. Bolgla LA, Keskula DR. Reliability of lower extremity functional performance tests. *J Orthop Sports Phys Ther.* 1997;26(3):138-142.
63. Broglio SP, Zhu W, Sapienza K, Park Y. Generalizability theory analysis of balance error scoring system reliability in healthy young adults. *J Athl Train.* 2009;44(5):497-502. doi: 10.4085/1062-6050-44.5.497.
64. Almangoush A, Herrington L, Jones R. A preliminary reliability study of a qualitative scoring system of limb alignment during single leg squat. *Phys Ther Rehabil.* 2014;1(1):2. <http://dx.doi.org/10.7243/2055-2386-1-2>. Accessed February 2, 2016.
65. Frost DM, Beach TA, Callaghan JP, McGill S. FMS scores change with performers' knowledge of the grading criteria—Are general whole-body movement screens capturing “dysfunction”? *J Strength Cond Res.* 2015;29(11):3037-3044. doi: 10.1097/JSC.0000000000000211.
66. Lakke SE, Soer R, Krijnen WP, van der Schans CP, Reneman MF, Geertzen JH. Influence of physical therapists' kinesiophobic beliefs on lifting capacity in healthy adults. *Phys Ther.* 2015;95(9):1224-1233. doi: 10.2522/ptj.20130194. Epub 2015 Apr 2.
67. Osteoarthritis Research Society International. Recommended performance-based tests to assess physical function in people diagnosed with hip or knee osteoarthritis. <http://oarsi.org/sites/default/files/docs/2013/manual.pdf>. Accessed Jul 31, 2015.
68. Borg GA. *Physical Performance and Perceived Exertion.* Gleeup Lund; 1962.
69. Gatchel RJ. Psychosocial factors that can influence the self-assessment of function. *J Occup Rehabil.* 2004;14(3):197-206.
70. Wilmore JH. Influence of motivation on physical work capacity and performance. *J Appl Physiol.* 1968;24(4):459-463.
71. Erdal K. Neuropsychological testing for sports-related concussion: how athletes can sandbag their baseline testing without detection. *Archi Clin Neuropsychol.* 2012;acs050. doi: 10.1093/arclin/acs050. Epub 2012 Jun 8.
72. Schatz P, Kelley T, Ott SD, et al. Utility of repeated assessment after invalid baseline neurocognitive test performance. *J Athl Train.* 2014;49(5):659-664. doi: 10.4085/1062-6050-49.3.37. Epub 2014 Aug 27.
73. McNamara JM, Stearne DJ. Flexible nonlinear periodization in a beginner college weight training class. *J Strength Cond Res.* 2010;24(8):2012-2017. doi: 10.1519/JSC.0b013e3181b1b15d.
74. Echemendia RJ, Iverson GL, McCrea M, et al. Advances in neuropsychological assessment of sport-related concussion. *Br J Sports Med.* 2013;47(5):294-298. doi: 10.1136/bjsports-2013-092186.
75. Schmidt JD, Register-Mihalik JK, Mihalik JP, Kerr ZY, Guskiewicz KM. Identifying impairments after concussion: normative data versus individualized baselines. *Med Sci Sports Exerc.* 2012;44(9):1621-1628. doi: 10.1249/MSS.0b013e318258a9fb.
76. Manske RC, Smith BS, Wyatt FB. Test-retest reliability of lower extremity functional tests after a closed kinetic chain isokinetic testing bout. *J Sport Rehabil.* 2003;12:119-132.
77. Marcora SM, Staiano W, Manning V. Mental fatigue impairs physical performance in humans. *J Appl Physiol.* 2009;106(3):857-864. doi: 10.1152/jappphysiol.91324.2008. Epub 2009 Jan 8.
78. Hart DL, Mioduski JE, Werneke MW, Stratford PW. Simulated computerized adaptive test for patients with lumbar spine impairments was efficient and produced valid measures of function. *J Clin Epidemiol.* 2006;59(9):947-956. Epub 2006 Jul 11.
79. Hart DL, Mioduski JE, Stratford PW. Simulated computerized adaptive tests for measuring functional status were efficient with good discriminant validity in patients with hip, knee, or foot/ankle impairments. *J Clin Epidemiol.* 2005;58(6):629-638.
80. Hsueh IP, Chen JH, Wang CH, et al. Development of a computerized adaptive test for assessing balance function in patients with stroke. *Phys Ther.* 2010;90(9):1336-1344. doi: 10.2522/ptj.20090395. Epub 2010 Jun 30.
81. Haley SM, Coster WJ, Dumas HM, et al. Accuracy and precision of the Pediatric Evaluation of Disability Inventory computer adaptive tests (PEDI-CAT). *Dev Med Child Neurol.* 2011;53(12):1100-1106. doi: 10.1111/j.1469-8749.2011.04107.x. Epub 2011 Nov 11.
82. Van Grinsven S, Van Cingel R, Holla C, Van Loon C. Evidence-based rehabilitation following anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2010;18(8):1128-1144. doi: 10.1007/s00167-009-1027-2. Epub 2010 Jan 13.
83. Minick KI, Kiesel KB, Burton L, Taylor A, Plisky P, Butler RJ. Interrater reliability of the functional movement screen. *J Strength Cond Res.* 2010;24(2):479-486. doi: 10.1519/JSC.0b013e3181c09c04.
84. Davies G. The need for critical thinking in rehabilitation. *J Sport Rehabil.* 1995;4(1):1-22.
85. Rudy TE, Lieber SJ, Boston JR, Gourley LM, Baysal E. Psychosocial predictors of physical performance in disabled individuals with chronic pain. *Clin J Pain.* 2003;19(1):18-30.

86. Swinkels-Meewisse IE, Roelofs J, Oostendorp RA, Verbeek AL, Vlaeyen JW. Acute low back pain: pain-related fear and pain catastrophizing influence physical performance and perceived disability. *Pain*. 2006;120(1-2):36-43.
87. de Gier M, Peters ML, Vlaeyen JW. Fear of pain, physical performance, and attentional processes in patients with fibromyalgia. *Pain*. 2003;104(1-2):121-130.
88. Rainville J, Hartigan C, Jouve C, Martinez E. The influence of intense exercise-based physical therapy program on back pain anticipated before and induced by physical activities. *Spine J*. 2004;4(2):176-183.
89. Gross DP. Are functional capacity evaluations affected by the patient's pain? *Curr Pain Headache Rep*. 2006;10(2):107-113.
90. Borg G, Holmgren A, Lindblad I. Quantitative evaluation of chest pain. *Acta Med Scand Suppl*. 1981;644:43-45.
91. Shechtman O, Anton SD, Kanasky WF Jr, Robinson ME. The use of the coefficient of variation in detecting sincerity of effort: a meta-analysis. *Work*. 2006;26(4):335-341.
92. Lechner DE, Bradbury SF, Bradley LA. Detecting sincerity of effort: a summary of methods and approaches. *Phys Ther*. 1998;78(8):867-888.
93. Hegedus EJ, McDonough S, Bleakley C, Cook CE, Baxter GD. Clinician-friendly lower extremity physical performance measures in athletes: a systematic review of measurement properties and correlation with injury, part 1. The tests for knee function including the hop tests. *Br J Sports Med*. 2015;49(10):642-648. doi: 10.1136/bjsports-2014-094094. Epub 2014 Dec 10.
94. Hegedus EJ, McDonough SM, Bleakley C, Baxter D, Cook CE. Clinician-friendly lower extremity physical performance tests in athletes: a systematic review of measurement properties and correlation with injury. Part 2--the tests for the hip, thigh, foot and ankle including the star excursion balance test. *Br J Sports Med*. 2015;49(10):649-656. doi: 10.1136/bjsports-2014-094341. Epub 2015 Jan 22.
95. Dobson F, Hinman RS, Hall M, Terwee CB, Roos EM, Bennell KL. Measurement properties of performance-based measures to assess physical function in hip and knee osteoarthritis: a systematic review. *Osteoarthritis Cartilage*. 2012;20(12):1548-1562. doi: 10.1016/j.joca.2012.08.015. Epub 2012 Aug 31.
96. Negrete RJ, Hanney WJ, Kolber MJ, et al. Reliability, minimal detectable change, and normative values for tests of upper extremity function and power. *J Strength Cond Res*. 2010;24(12):3318-3325. doi: 10.1519/JSC.0b013e3181e7259c.
97. McGill SM, Childs A, Liebenson C. Endurance times for low back stabilization exercises: clinical targets for testing and training from a normal database. *Arch Phys Med Rehabil*. 1999;80(8):941-944.
98. Finnoff JT, Peterson VJ, Hollman JH, Smith J. Intrarater and interrater reliability of the Balance Error Scoring System (BESS). *PM R*. 2009;1(1):50-54. doi: 10.1016/j.pmrj.2008.06.002. Epub 2008 Dec 10.
99. Hunt TN, Ferrara MS, Bornstein RA, Baumgartner TA. The reliability of the modified Balance Error Scoring System. *Clin J Sport Med*. 2009;19(6):471-475. doi: 10.1097/JSM.0b013e3181c12c7b.
100. Munro AG, Herrington LC. Between-session reliability of the star excursion balance test. *Phys Ther Sport*. 2010;11(4):128-132. doi: 10.1016/j.ptsp.2010.07.002. Epub 2010 Aug 17.
101. Kristensen MT, Nielsen AO, Topp UM, et al. Number of test trials needed for performance stability and interrater reliability of the one leg stand test in patients with a major non-traumatic lower limb amputation. *Gait Posture*. 2014;39(1):424-429. doi: 10.1016/j.gaitpost.2013.08.017. Epub 2013 Aug 25.
102. Stratford PW, Riddle DL. When minimal detectable change exceeds a diagnostic test-based threshold change value for an outcome measure: resolving the conflict. *Phys Ther*. 2012;92(10):1338-1347. doi: 10.2522/ptj.20120002. Epub 2012 Jul 5.
103. Mannerkorpi K, Svantesson U, Broberg C. Relationships between performance-based tests and patients' ratings of activity limitations, self-efficacy, and pain in fibromyalgia. *Arch Phys Med Rehabil*. 2006;87(2):259-264.
104. Filho IT, Simmonds MJ, Protas EJ, Jones S. Back pain, physical function, and estimates of aerobic capacity: what are the relationships among methods and measures? *Am J Phys Med Rehabil*. 2002;81(12):913-920.
105. Thomas SG, Pagura SM, Kennedy D. Physical activity and its relationship to physical performance in patients with end stage knee osteoarthritis. *J Orthop Sports Phys Ther*. 2003;33(12):745-754.