

Outcomes in Orthopaedic Physical Therapy Practice


Independent Study Course 26.1.3

Lower Extremity Outcome Measures

Joseph Zeni, Jr. PT, PhD
Assistant Professor
University of Delaware
Department of Physical Therapy
Newark, Delaware

Kathleen Madara, PT, DPT
University of Delaware
Department of Physical Therapy
Biomechanics and Movement Science Program
Newark, Delaware

Orthopaedic Section, APTA, Inc.
Downloaded from www.orthopteam.org at the Orthopaedic Section on September 26, 2018. For personal use only. No other uses without permission.
Copyright © 2016 Orthopaedic Section, APTA, Inc. All rights reserved.



REFERENCES

1. Mizner RL, Petterson SC, Clements KE, Zeni JA, Irrgang JJ, Snyder-Mackler L. Measuring functional improvement after total knee arthroplasty requires both performance-based and patient-report assessments: a longitudinal analysis of outcomes. *J Arthroplasty*. 2011;26(5):728-737. doi: 10.1016/j.arth.2010.06.004. Epub 2010 Sep 20.
2. Zeni J, Abujaber S, Pozzi F, Rasis L. Strength and pain are related to different measures of functional ability in patients with end-stage hip osteoarthritis. *Arthritis Care Res (Hoboken)*. 2014;66(10):1506-12. doi: 10.1002/acr.22329.
3. Stratford PW, Kennedy DM. Performance measures were necessary to obtain a complete picture of osteoarthritic patients. *J Clin Epidemiol*. 2006;59(2):160-167. Epub 2005 Dec 27.
4. Mathias S, Nayak US, Isaacs B. Balance in elderly patients: the "get-up and go" test. *Arch Phys Med Rehabil*. 1986;67(6):387-389.
5. Podsiadlo D, Richardson S. The timed "Up & Go": a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc*. 1991;39(2):142-148.
6. Kennedy DM, Stratford PW, Wessel J, Gollish JD, Penney D. Assessing stability and change of four performance measures: a longitudinal study evaluating outcome following total hip and knee arthroplasty. *BMC Musculoskelet Disord*. 2005;6:3.
7. Dite W, Connor HJ, Curtis HC. Clinical identification of multiple fall risk early after unilateral transtibial amputation. *Arch Phys Med Rehabil*. 2007;88(1):109-114. doi:10.1016/j.apmr.2006.10.015.
8. Zeni JA, Snyder-Mackler L. Clinical outcomes after simultaneous bilateral total knee arthroplasty: comparison to unilateral total knee arthroplasty and healthy controls. *J Arthroplasty*. 2010;25(4):541-546. doi:10.1016/j.arth.2009.02.016. Epub 2009 Apr 7.
9. Dobson F, Hinman RS, Roos EM, et al. OARSI recommended performance-based tests to assess physical function in people diagnosed with hip or knee osteoarthritis. *Osteoarthritis Cartilage*. 2013;21(10):1625-1626. doi: 10.1016/j.joca.2013.07.011. Epub 2013 Jul 21.
10. Freter SH, Fruchter N. Relationship between timed "up and go" and gait time in an elderly orthopaedic rehabilitation population. *Clin Rehabil*. 2000;14(1):96-101.
11. Wright AA, Cook CE, Baxter GD, Dockerty JD, Abbott JH. A comparison of 3 methodological approaches to defining major clinically important improvement of 4 performance measures in patients with hip osteoarthritis. *J Orthop Sports Phys Ther*. 2011;41(5):319-327. doi:10.2519/jospt.2011.3515. Epub 2011 Feb 18.
12. Kristensen MT, Foss NB, Kehlet H. Timed "up & go" test as a predictor of falls within 6 months after hip fracture surgery. *Phys Ther*. 2007;87(1):24-30. doi:10.2522/ptj.20050271. Epub 2006 Dec 1.
13. Arnold CM, Faulkner RA. The history of falls and the association of the timed up and go test to falls and near-falls in older adults with hip osteoarthritis. *BMC Geriatr*. 2007;7:17. doi:10.1186/1471-2318-7-17.
14. Steffen TM, Hacker TA, Mollinger L. Age- and gender-related test performance in community-dwelling elderly people: Six-Minute Walk Test, Berg Balance Scale, Timed Up & Go Test, and gait speeds. *Phys Ther*. 2002;82(2):128-137.
15. McGavin CR, Gupta SP, McHardy GJ. Twelve-minute walking test for assessing disability in chronic bronchitis. *Br Med J*. 1976;1(6013):822-823.
16. Butland RJ, Pang J, Gross ER, Woodcock AA, Geddes DM. Two-, six-, and 12-minute walking tests in respiratory disease. *Br Med J (Clin Res Ed)*. 1982;284(6329):1607-1608.
17. Guyatt GH, Sullivan MJ, Thompson PJ, et al. The 6-minute walk: a new measure of exercise capacity in patients with chronic heart failure. *Can Med Assoc J*. 1985;132(8):919-923.
18. Bade MJ, Wolfe P, Zeni JA, Stevens-Lapsley JE, Snyder-Mackler L. Predicting poor physical performance after total knee arthroplasty. *J Orthop Res*. 2012;30(11):1805-1810. doi: 10.1002/jor.22140. Epub 2012 Apr 26.
19. Bade MJ, Kohrt WM, Stevens-Lapsley JE. Outcomes before and after total knee arthroplasty compared to healthy adults. *J Orthop Sports Phys Ther*. 2010;40(9):559-567. doi: 10.2519/jospt.2010.3317.
20. Marmon AR, Milcarek BI, Snyder-Mackler L. Associations between knee extensor power and functional performance in patients after total knee arthroplasty and normal controls without knee pain. *Int J Sports Phys Ther*. 2014;9(2):168-178.
21. Casanova C, Celli BR, Barria P, et al. The 6-min walk distance in healthy subjects: reference standards from seven countries.

- Eur Respir J.* 2011;37(1):150-156. doi:10.1183/09031936.00194909. Epub 2010 Jun 4.
22. Beekman E, Mesters I, Hendriks EJM, et al. Course length of 30 metres versus 10 metres has a significant influence on six-minute walk distance in patients with COPD: an experimental crossover study. *J Physiother.* 2013;59(3):169-176. doi:10.1016/S1836-9553(13)70181-4.
 23. Kennedy DM, Stratford PW, Wessel J, Gollish JD, Penney D. Assessing stability and change of four performance measures: a longitudinal study evaluating outcome following total hip and knee arthroplasty. *BMC Musculoskelet Disord.* 2005;6:3. doi:10.1186/1471-2474-6-3.
 24. Parent E, Moffet H. Comparative responsiveness of locomotor tests and questionnaires used to follow early recovery after total knee arthroplasty. *Arch Phys Med Rehabil.* 2002;83(1):70-80.
 25. French HP, Fitzpatrick M, FitzGerald O. Responsiveness of physical function outcomes following physiotherapy intervention for osteoarthritis of the knee: an outcome comparison study. *Physiotherapy.* 2011;97(4):302-308. doi:10.1016/j.physio.2010.03.002. Epub 2010 May 4.
 26. Ko V, Naylor JM, Harris IA, Crosbie J, Yeo AE. The six-minute walk test is an excellent predictor of functional ambulation after total knee arthroplasty. *BMC Musculoskelet Disord.* 2013;14:145. doi:10.1186/1471-2474-14-145.
 27. Rikli R, Jones CJ. Development and validation of a functional fitness test for community-residing older adults. *J Aging Phys Act.* 1999;7:129-161.
 28. McCarthy EK, Horvat MA, Holsberg PA, Wisenbaker JM. Repeated chair stands as a measure of lower limb strength in sexagenarian women. *J Gerontol A Biol Sci Med Sci.* 2004;59(11):1207-1212.
 29. Jones CJ, Rikli RE, Beam WC. A 30-s chair-stand test as a measure of lower body strength in community-residing older adults. *Res Q Exerc Sport.* 1999;70(2):113-119. doi:10.1080/02701367.1999.10608028.
 30. Gill S, McBurney H. Reliability of performance-based measures in people awaiting joint replacement surgery of the hip or knee. *Physiother Res Int.* 2008;13(3):141-152. doi:10.1002/pri.411.
 31. Noyes FR, Barber SD, Mangine RE. Abnormal lower limb symmetry determined by function hop tests after anterior cruciate ligament rupture. *Am J Sports Med.* 1991;19(5):513-518.
 32. Barber SD, Noyes FR, Mangine RE, McCloskey JW, Hartman W. Quantitative assessment of functional limitations in normal and anterior cruciate ligament-deficient knees. *Clin Orthop Relat Res.* 1990;(255):204-214.
 33. Adams D, Logerstedt DS, Hunter-Giordano A, Axe MJ, Snyder-Mackler L. Current concepts for anterior cruciate ligament reconstruction: a criterion-based rehabilitation progression. *J Orthop Sports Phys Ther.* 2012;42(7):601-614. doi:10.2519/jospt.2012.3871. Epub 2012 Mar 8.
 34. Gerber JP, Williams GN, Scoville CR, Arciero RA, Taylor DC. Persistent disability associated with ankle sprains: a prospective examination of an athletic population. *Foot Ankle Int.* 1998;19(10):653-660.
 35. Williams GN, Molloy JM, DeBerardino TM, Arciero RA, Taylor DC. Evaluation of the Sports Ankle Rating System in young, athletic individuals with acute lateral ankle sprains. *Foot Ankle Int.* 2003;24(3):274-282.
 36. Johnson MR, Stoneman PD. Comparison of a lateral hop test versus a forward hop test for functional evaluation of lateral ankle sprains. *J Foot Ankle Surg.* 2007;46(3):162-174. doi:10.1053/j.jfas.2006.12.007.
 37. Kea J, Kramer J, Forwell L, Birmingham T. Hip abduction-adduction strength and one-leg hop tests: test-retest reliability and relationship to function in elite ice hockey players. *J Orthop Sports Phys Ther.* 2001;31(8):446-455. doi:10.2519/jospt.2001.31.8.446.
 38. Meylan C, McMaster T, Cronin J, Mohammad NI, Rogers C, Deklerk M. Single-leg lateral, horizontal, and vertical jump assessment: reliability, interrelationships, and ability to predict sprint and change-of-direction performance. *J Strength Cond Res.* 2009;23(4):1140-1147. doi:10.1519/JSC.0b013e318190f9c2.
 39. Miller MG, Herniman JJ, Ricard MD, Cheatham CC, Michael TJ. The effects of a 6-week plyometric training program on agility. *J Sports Sci Med.* 2006;5(3):459-465. eCollection 2006.
 40. Raya MA, Gailey RS, Gaunaud IA, et al. Comparison of three agility tests with male servicemembers: Edgren Side Step Test, T-Test, and Illinois Agility Test. *J Rehabil Res Dev.* 2013;50(7):951-960. doi:10.1682/JRRD.2012.05.0096.
 41. Ramírez-Campillo R, Meylan C, Alvarez C, et al. Effects of in-season low-volume high-intensity plyometric training on explosive actions and endurance of young soccer players. *J Strength Cond Res.* 2014;28(5):1335-42. doi:10.1519/JSC.0000000000000284.
 42. Marmon AR, McClelland JA, Stevens-Lapsley J, Snyder-Mackler L. Single-step test for unilateral limb ability following total knee arthroplasty.

- J Orthop Sports Phys Ther.* 2013;43(2):66-73. doi:10.2519/jospt.2013.4372 Epub 2012 Nov 16.
43. Silbernagel KG, Nilsson-Helander K, Thomeé R, Eriksson BI, Karlsson J. A new measurement of heel-rise endurance with the ability to detect functional deficits in patients with Achilles tendon rupture. *Knee Surg Sports Traumatol Arthrosc.* 2010;18(2):258-64. doi:10.1007/s00167-009-0889-7. Epub 2009 Aug 19.
44. Binkley JM, Stratford PW, Lott SA, Riddle DL. The Lower Extremity Functional Scale (LEFS): scale development, measurement properties, and clinical application. North American Orthopaedic Rehabilitation Research Network. *Phys Ther.* 1999;79(4):371-383.
45. Irrgang JJ, Snyder-Mackler L, Wainner RS, Fu FH, Harner CD. Development of a patient-reported measure of function of the knee. *J Bone Joint Surg Am.* 1998;80(8):1132-1145.
46. Robinson JM, Cook JL, Purdam C, et al. The VISA-A questionnaire: a valid and reliable index of the clinical severity of Achilles tendinopathy. *Br J Sports Med.* 2001;35(5):335-341.
47. Pua Y-H, Cowan SM, Wrigley TV, Bennell KL. The Lower Extremity Functional Scale could be an alternative to the Western Ontario and McMaster Universities Osteoarthritis Index physical function scale. *J Clin Epidemiol.* 2009;62(10):1103-11. doi:10.1016/j.jclinepi.2008.11.011. Epub 2009 Mar 17.
48. Stratford PW, Kennedy DM, Hanna SE. Condition-specific Western Ontario McMaster Osteoarthritis Index was not superior to region-specific Lower Extremity Functional Scale at detecting change. *J Clin Epidemiol.* 2004;57(10):1025-1032. doi:10.1016/j.jclinepi.2004.03.008.
49. Kennedy DM, Stratford PW, Riddle DL, Hanna SE, Gollish JD. Assessing recovery and establishing prognosis following total knee arthroplasty. *Phys Ther.* 2008;88(1):22-32. doi:10.2522/ptj.20070051. Epub 2007 Nov 6.
50. Alcock GK, Werstine MS, Robbins SM, Stratford PW. Longitudinal changes in the lower extremity functional scale after anterior cruciate ligament reconstructive surgery. *Clin J Sport Med.* 2012;22(3):234-239. doi:10.1097/JSM.0b013e31824cb53d.
51. Lin CW, Moseley AM, Refshauge KM, Bundy AC. The lower extremity functional scale has good clinimetric properties in people with ankle fracture. *Phys Ther.* 2009;89(6):580-588. doi:10.2522/ptj.20080290. Epub 2009 May 7.
52. Mohtadi NG, Griffin DR, Pedersen ME, et al. The development and validation of a self-administered quality-of-life outcome measure for young, active patients with symptomatic hip disease: The International Hip Outcome Tool (iHOT-33). *Arthroscopy.* 2012;28(5):595-605; quiz 606-10. e1. doi: 10.1016/j.arthro.2012.03.013.
53. Thorborg K, Tijssen M, Habets B, et al. Patient-Reported Outcome (PRO) questionnaires for young to middle-aged adults with hip and groin disability: a systematic review of the clinimetric evidence. *Br J Sports Med.* 2015; 49(12):812. doi:10.1136/bjsports-2014-094224. Epub 2015 Jan 13.
54. Ramisetty N, Kwon Y, Mohtadi N. Patient-reported outcome measures for hip preservation surgery--a systematic review of the literature. *J Hip Preserv Surg.* 2015;2(1):15-27. doi:10.1093/jhps/hnv002.
55. Kemp JL, Collins NJ, Roos EM, Crossley KM. Psychometric properties of patient-reported outcome measures for hip arthroscopic surgery. *Am J Sports Med.* 2013;41(9):2065-2073. doi:10.1177/0363546513494173. Epub 2013 Jul 8.
56. Jónasson P, Baranto A, Karlsson J, et al. A standardised outcome measure of pain, symptoms and physical function in patients with hip and groin disability due to femoro-acetabular impingement: Cross-cultural adaptation and validation of the international Hip Outcome Tool (iHOT12) in Swedish. *Knee Surg Sport Traumatol Arthrosc.* 2014;22(4):826-834. doi:10.1007/s00167-013-2710-x. Epub 2013 Oct 18.
57. Ruiz-Ibán MA, Seijas R, Sallent A, et al. The international Hip Outcome Tool-33 (iHOT-33): multicenter validation and translation to Spanish. *Health Qual Life Outcomes.* 2015;13:62. doi:10.1186/s12955-015-0255-z.
58. Griffin DR, Parsons N, Mohtadi NGH, Safran MR. A short version of the International Hip Outcome Tool (iHOT-12) for use in routine clinical practice. *Arthroscopy.* 2012;28(5):611-618. doi:10.1016/j.arthro.2012.02.027.
59. Thorborg K, Hölmich P, Christensen R, Petersen J, Roos EM. The Copenhagen Hip and Groin Outcome Score (HAGOS): development and validation according to the COSMIN checklist. *Br J Sports Med.* 2011;45(6):478-491. doi:10.1136/bjsm.2010.080937.
60. Klässbo M, Larsson E, Mannevik E. Hip disability and osteoarthritis outcome score. An extension of the Western Ontario and McMaster Universities Osteoarthritis Index. *Scand J Rheumatol.* 2003;32(1):46-51. doi:10.1080/03009740310000409.
61. Davis AM, Perruccio AV, Canizares M, et al. The development of a short measure of physical function for hip OA HOOS-Physical Function

- Shortform (HOOS-PS): an OARSI/OMERACT initiative. *Osteoarthritis Cartilage*. 2008;16(5):551-559. doi:10.1016/j.joca.2007.12.016. Epub 2008 Mar 4.
62. Davis AM, Perruccio AV, Canizares M, et al. Comparative, validity and responsiveness of the HOOS-PS and KOOS-PS to the WOMAC physical function subscale in total joint replacement for Osteoarthritis. *Osteoarthritis Cartilage*. 2009;17(7):843-847. doi:10.1016/j.joca.2009.03.025. Epub 2009 Jan 31.
 63. Nilsson AK, Lohmander LS, Klässbo M, Roos EM. Hip disability and osteoarthritis outcome score (HOOS)--validity and responsiveness in total hip replacement. *BMC Musculoskeletal Disord*. 2003;4:10. doi:10.1186/1471-2474-4-10. Epub 2003 May 30.
 64. Nilsson A, Bremander A. Measures of hip function and symptoms: Harris Hip Score (HHS), Hip Disability and Osteoarthritis Outcome Score (HOOS), Oxford Hip Score (OHS), Lequesne Index of Severity for Osteoarthritis of the Hip (LISOH), and American Academy of Orthopedic Surgeons (AAOS) Hip and Knee Questionnaire. *Arthritis Care Res*. 2011;63(SUPPL. 11):200-207. doi:10.1002/acr.20549.
 65. de Groot IB, Reijman M, Terwee CB, et al. Validation of the Dutch version of the Hip Disability and Osteoarthritis Outcome Score. *Osteoarthritis Cartilage*. 2007;15(1):104-109. doi:10.1016/j.joca.2006.06.014. Epub 2006 Aug 4.
 66. Martin RL, Philippon MJ. Evidence of reliability and responsiveness for the hip outcome score. *Arthroscopy*. 2008;24(6):676-682. doi:10.1016/j.arthro.2007.12.011. Epub 2008 Mar 12.
 67. Martin RL, Philippon MJ. Evidence of reliability and responsiveness for the hip outcome score. *Arthroscopy*. 2006;22(12):1304-1311. doi:10.1016/j.arthro.2007.12.011.
 68. Martin RL. Hip arthroscopy and outcome assessment. *Oper Tech Orthop*. 2005;15(3):290-296. doi:10.1053/j.oto.2005.06.008.
 69. Martin RL, Philippon MJ. Evidence of validity for the hip outcome score in hip arthroscopy. *Arthroscopy*. 2007;23(8):822-826. doi:10.1016/j.arthro.2007.02.004.
 70. Martin RL, Kelly BT, Philippon MJ. Evidence of validity for the hip outcome score. *Arthroscopy*. 2006;22(12):1304-1311. doi:10.1016/j.arthro.2006.07.027.
 71. Chahal J, Van Thiel GS, Mather RC, Lee S, Salata MJ, Nho SJ. The Minimal Clinically Important Difference (MCID) And Patient Acceptable Symptomatic State (PASS) For The Modified Harris Hip Score And Hip Outcome Score Among Patients Undergoing Surgical Treatment For Femoroacetabular Impingement. *Orthop J Sport Med*. 2014;2(2 Suppl): 2325967114S00105. doi:10.1177/2325967114S00105.
 72. Harris WH. Traumatic arthritis of the hip after dislocation and acetabular fractures: treatment by mold arthroplasty. An end-result study using a new method of result evaluation. *J Bone Joint Surg Am*. 1969;51(4):737-755.
 73. Ravi B, Croxford R, Reichmann WM, Losina E, Katz JN, Hawker GA. The changing demographics of total joint arthroplasty recipients in the United States and Ontario from 2001 to 2007. *Best Pract Res Clin Rheumatol*. 2012;26(5):637-647. doi:10.1016/j.berh.2012.07.014.
 74. Roos EM, Roos HP, Lohmander LS, Ekdahl C, Beynon BD. Knee Injury and Osteoarthritis Outcome Score (KOOS)--development of a self-administered outcome measure. *J Orthop Sports Phys Ther*. 1998;28(2):88-96.
 75. Roos EM, Toksvig-Larsen S. Knee injury and Osteoarthritis Outcome Score (KOOS) - validation and comparison to the WOMAC in total knee replacement. *Health Qual Life Outcomes*. 2003;1:17. doi:10.1186/1477-7525-1-17.
 76. Roos EM, Roos HP, Lohmander LS. WOMAC Osteoarthritis Index--additional dimensions for use in subjects with post-traumatic osteoarthritis of the knee. Western Ontario and McMaster Universities. *Osteoarthritis Cartilage*. 1999;7(2):216-221. doi:10.1053/joca.1998.0153.
 77. Bekkers JEJ, de Windt TS, Raijmakers NJH, Dhert WJA, Saris DBF. Validation of the Knee Injury and Osteoarthritis Outcome Score (KOOS) for the treatment of focal cartilage lesions. *Osteoarthritis Cartilage*. 2009;17(11):1434-1439. doi:10.1016/j.joca.2009.04.019.
 78. Paradowski PT, Bergman S, Sundén-Lundius A, Lohmander LS, Roos EM. Knee complaints vary with age and gender in the adult population. Population-based reference data for the Knee Injury and Osteoarthritis Outcome Score (KOOS). *BMC Musculoskeletal Disord*. 2006;7(1):38. doi:10.1186/1471-2474-7-38.
 79. Collins NJ, Misra D, Felson DT, Crossley KM, Roos EM. Measures of Knee Function: International Knee Documentation Committee (IKDC) Subjective Knee Evaluation Form, Knee Injury and Osteoarthritis Outcome Score (KOOS), Knee Injury and Osteoarthritis Outcome Score Physical Function Short Form (KOOS-PS), Knee Outcome Survey Activities of Daily Living Scale (KOS-ADL), Lysholm Knee Scoring Scale, Oxford Knee Score (OKS), Western Ontario and McMaster

- Universities Osteoarthritis Index (WOMAC), Activity Rating Scale (ARS), and Tegner Activity Score (TAS). *Arthritis Care Res (Hoboken)*. 2011;63 Suppl 1:S208-228. doi:10.1002/Acr.20632.
80. Irrgang JJ, Anderson AF, Boland AL, et al. Development and validation of the international knee documentation committee subjective knee form. *Am J Sports Med*. 2001;29(5):600-613.
81. Hefti F, Müller W, Jakob RP, Stäubli HU. Evaluation of knee ligament injuries with the IKDC form. *Knee Surg Sports Traumatol Arthrosc*. 1993;1(3-4):226-234.
82. Padua R, Bondi R, Ceccarelli E, et al. Italian version of the International Knee Documentation Committee Subjective Knee Form: cross-cultural adaptation and validation. *Arthroscopy*. 2004;20(8):819-823. doi:10.1016/j.arthro.2004.06.011.
83. Crawford K, Briggs KK, Rodkey WG, Steadman JR. Reliability, validity, and responsiveness of the IKDC score for meniscus injuries of the knee. *Arthroscopy*. 2007;23(8):839-44. doi:10.1016/j.arthro.2007.02.005.
84. Greco NJ, Anderson AF, Mann BJ, et al. Responsiveness of the International Knee Documentation Committee Subjective Knee Form in comparison to the Western Ontario and McMaster Universities Osteoarthritis Index, modified Cincinnati Knee Rating System, and Short Form 36 in patients with focal art. *Am J Sports Med*. 2010;38(5):891-902. doi:10.1177/0363546509354163.
85. van de Graaf VA, Wolterbeek N, Scholtes VA, Mutsaerts EL, Poolman RW. Reliability and validity of the IKDC, KOOS, and WOMAC for patients with meniscal injuries. *Am J Sports Med*. 2014;42(6):1408-1416. doi:10.1177/0363546514524698. Epub 2014 Mar 11.
86. van Meer BL, Meuffels DE, Vissers MM, et al. Knee injury and Osteoarthritis Outcome Score or International Knee Documentation Committee Subjective Knee Form: which questionnaire is most useful to monitor patients with an anterior cruciate ligament rupture in the short term? *Arthroscopy*. 2013;29(4):701-715. doi:10.1016/j.arthro.2012.12.015. Epub 2013 Feb 8.
87. Logerstedt D, Di Stasi S, Grindem H, et al. Self-reported knee function can identify athletes who fail return-to-activity criteria up to 1 year after anterior cruciate ligament reconstruction: a Delaware-Oslo ACL cohort study. *J Orthop Sports Phys Ther*. 2014;44(12):914-923. doi:10.2519/jospt.2014.4852. Epub 2014 Oct 27.
88. Lentz TA, Zeppieri G, Tillman SM, et al. Return to preinjury sports participation following anterior cruciate ligament reconstruction: contributions of demographic, knee impairment, and self-report measures. *J Orthop Sports Phys Ther*. 2012;42(11):893-901. doi:10.2519/jospt.2012.4077. Epub 2012 Aug 2.
89. Anderson AF. The International Knee Documentation Committee Subjective Knee Evaluation Form: normative data. *Am J Sports Med*. 2005;34(1):128-135. doi:10.1177/0363546505280214. Epub 2005 Oct 11.
90. Grindem H, Logerstedt D, Eitzen I, et al. Single-legged hop tests as predictors of self-reported knee function in nonoperatively treated individuals with anterior cruciate ligament injury. *Am J Sports Med*. 2011;39(11):2347-2354. doi:10.1177/0363546511417085. Epub 2011 Aug 9.
91. Martin RL, Irrgang JJ, Burdett RG, Conti SF, Van Swearingen JM. Evidence of validity for the Foot and Ankle Ability Measure (FAAM). *Foot Ankle Int*. 2005;26(11):968-983.
92. Kivlan BR, Martin RL, Wukich DK. Responsiveness of the foot and ankle ability measure (FAAM) in individuals with diabetes. *Foot (Edinb)*. 2011;21(2):84-87. doi:10.1016/j.foot.2011.04.004. Epub 2011 May 7.
93. Roos EM, Brandsson S, Karlsson J. Validation of the foot and ankle outcome score for ankle ligament reconstruction. *Foot Ankle Int*. 2001;22(10):788-794.
94. Chen L, Lyman S, Do H, et al. Validation of Foot and Ankle Outcome Score for Hallux Valgus. *Foot Ankle Int*. 2012;33(12):1145-1155. doi:10.3113/FAI.2012.1145.
95. Golightly YM, Devellis RF, Nelson AE, et al. Psychometric properties of the foot and ankle outcome score in a community-based study of adults with and without osteoarthritis. *Arthritis Care Res (Hoboken)*. 2014;66(3):395-403. doi:10.1002/acr.22162.
96. Mani SB, Do H, Vulcano E, et al. Evaluation of the foot and ankle outcome score in patients with osteoarthritis of the ankle. *Bone Joint J*. 2015;97-B(5):662-667. doi:10.1302/0301-620X.97B5.33940.
97. Mani SB, Brown HC, Nair P, et al. Validation of the Foot and Ankle Outcome Score in adult acquired flatfoot deformity. *Foot Ankle Int*. 2013;34(8):1140-6. doi:10.1177/1071100713483117. Epub 2013 Mar 19.
98. Sierevelt IN, Beimers L, van Bergen CJA, Haverkamp D, Terwee CB, Kerkhoffs GM.

- Validation of the Dutch language version of the Foot and Ankle Outcome Score. *Knee Surg Sports Traumatol Arthrosc.* 2015;23(8):2413-2419. doi:10.1007/s00167-014-3017-2. Epub 2014 May 4.
99. van Bergen CJA, Sierevelt IN, Hoogervorst P, Waizy H, van Dijk CN, Becher C. Translation and validation of the German version of the foot and ankle outcome score. *Arch Orthop Trauma Surg.* 2014;134(7):897-901. doi:10.1007/s00402-014-1994-8. Epub 2014 Apr 19.
 100. Dawson J, Fitzpatrick R, Murray D, Carr A. Questionnaire on the perceptions of patients about total knee replacement. *J Bone Joint Surg Br.* 1998;80(1):63-69.
 101. Murray DW, Fitzpatrick R, Rogers K, et al. The use of the Oxford hip and knee scores. *J Bone Joint Surg Br.* 2007;89(8):1010-1014. doi:10.1302/0301-620X.89B8.19424.
 102. Impellizzeri FM, Mannion AF, Leunig M, Bizzini M, Naal FD. Comparison of the reliability, responsiveness, and construct validity of 4 different questionnaires for evaluating outcomes after total knee arthroplasty. *J Arthroplasty.* 2011;26(6):861-869. doi:10.1016/j.arth.2010.07.027. Epub 2010 Nov 12.
 103. Jenny J-Y, Diesinger Y. The Oxford Knee Score: compared performance before and after knee replacement. *Orthop Traumatol Surg Res.* 2012;98(4):409-412. doi:10.1016/j.otsr.2012.03.004. Epub 2012 May 18.
 104. Harris KK, Dawson J, Jones LD, Beard DJ, Price AJ. Extending the use of PROMs in the NHS--using the Oxford Knee Score in patients undergoing non-operative management for knee osteoarthritis: a validation study. *BMJ Open.* 2013;3(8):e003365. doi:10.1136/bmjopen-2013-003365.
 105. Harris K, Dawson J, Doll H, et al. Can pain and function be distinguished in the Oxford Knee Score in a meaningful way? An exploratory and confirmatory factor analysis. *Qual Life Res.* 2013;22(9):2561-2568. doi:10.1007/s11136-013-0393-x.
 106. Talbot S, Hooper G, Stokes A, Zordan R. Use of a new high-activity arthroplasty score to assess function of young patients with total hip or knee arthroplasty. *J Arthroplasty.* 2010;25(2):268-73. doi:10.1016/j.arth.2008.09.019.
 107. Jenny JY, Louis P, Diesinger Y. High Activity Arthroplasty Score has a lower ceiling effect than standard scores after knee arthroplasty. *J Arthroplasty.* 2014;29(4):719-721. doi:10.1016/j.arth.2013.07.015. Epub 2013 Aug 13.
 108. Iversen JV, Bartels EM, Langberg H. The victorian institute of sports assessment - achilles questionnaire (visa-a) - a reliable tool for measuring achilles tendinopathy. *Int J Sports Phys Ther.* 2012;7(1):76-84.
 109. Tumilty S, Munn J, Abbott JH, McDonough S, Hurley DA, Baxter GD. Laser therapy in the treatment of achilles tendinopathy: a pilot study. *Photomed Laser Surg.* 2008;26(1):25-30. doi:10.1089/pho.2007.2126.
 110. de Vos RJ, Weir A, van Schie HTM, et al. Platelet-rich plasma injection for chronic Achilles tendinopathy: a randomized controlled trial. *JAMA.* 2010;303(2):144-149. doi:10.1001/jama.2009.1986.
 111. Nilsson-Helander K, Thomeé R, Silbernagel KG, et al. The Achilles tendon Total Rupture Score (ATRS): development and validation. *Am J Sports Med.* 2007;35(3):421-426. doi:10.1177/0363546506294856.
 112. Maffulli N, Kenward MG, Testa V, Capasso G, Regine R, King JB. Clinical diagnosis of Achilles tendinopathy with tendinosis. *Clin J Sport Med.* 2003;13:11-15. doi:10.1097/00042752-200301000-00003.
 113. Silbernagel KG, Gustavsson A, Thomeé R, Karlsson J. Evaluation of lower leg function in patients with Achilles tendinopathy. *Knee Surg Sport Traumatol Arthrosc.* 2006;14:1207-1217. doi:10.1007/s00167-006-0150-6.
 114. Silbernagel KG, Thomeé R, Thomeé P, Karlsson J. Eccentric overload training for patients with chronic Achilles tendon pain--a randomised controlled study with reliability testing of the evaluation methods. *Scand J Med Sci Sports.* 2001;11(4):197-206.