

Current Concepts of **Orthopaedic Physical Therapy**

Independent Study Course 26.2.12



4th Edition

The Foot and Ankle: Physical Therapy Patient Management Using Current Evidence

Jeff Houck, PT, PhD
George Fox University
Newberg, Oregon

Christopher Neville, PT, PhD
SUNY Upstate Medical University
Syracuse, New York

Ruth Chimenti, DPT, PhD
University of Iowa
Iowa City, Iowa



REFERENCES

1. Westblad P, Hashimoto T, Winson I, Lundberg A, Arndt A. Differences in ankle-joint complex motion during the stance phase of walking as measured by superficial and bone-anchored markers. *Foot Ankle Int.* 2002;23(9):856-863.
2. Tochigi Y, Amendola A, Rudert MJ, et al. The role of the interosseous talocalcaneal ligament in subtalar joint stability. *Foot Ankle Int.* 2004;25(8):588-596.
3. Budny A. Subtalar joint instability: current clinical concepts. *Clin Podiatr Med Surg.* 2004;21(3):449-460.
4. Boss AP, Hintermann B. Anatomical study of the medial ankle ligament complex. *Foot Ankle Int.* 2002;23(6):547-553.
5. Deland JT, de Asla RJ, Sung IH, Ernberg LA, Potter HG. Posterior tibial tendon insufficiency: which ligaments are involved? *Foot Ankle Int.* 2005;26(6):427-435.
6. Dinucci KR, Christensen JC, Dinucci KA. Biomechanical consequences of lateral column lengthening of the calcaneus: Part I. Long plantar ligament strain. *J Foot Ankle Surg.* 2004;43(1):10-15.
7. Deland JT. The adult acquired flatfoot and spring ligament complex. Pathology and implications for treatment. *Foot Ankle Clin.* 2001;6(1):129-135.
8. Ridola CG, Cappello F, Marciano V, et al. The synovial joints of the human foot. *Ital J Anat Embryol.* 2007;112(2):61-80.
9. Solan MC, Moorman CT 3rd, Miyamoto RG, Jasper LE, Belkoff SM. Ligamentous restraints of the second tarsometatarsal joint: a biomechanical evaluation. *Foot Ankle Int.* 2001;22(8):637-641.
10. Neptune RR, Kautz SA, Zajac FE. Contributions of the individual ankle plantar flexors to support, forward progression and swing initiation during walking. *J Biomech.* 2001;34(11):1387-1398.
11. Neptune RR, Zajac FE, Kautz SA. Muscle mechanical work requirements during normal walking: the energetic cost of raising the body's center-of-mass is significant. *J Biomech.* 2004;37(6):817-825.
12. Perry J. *Gait Analysis. Normal and Pathologic Function.* Thorofare, NJ: SLACK Incorporated; 1992.
13. Sutherland DH. An electromyographic study of the plantar flexors of the ankle in normal walking on the level. *J Bone Joint Surg Am.* 1966;48(1):66-71.
14. MacWilliams BA, Cowley M, Nicholson DE. Foot kinematics and kinetics during adolescent gait. *Gait Posture.* 2003;17(3):214-224.
15. Klein P, Mattys S, Rooze M. Moment arm length variations of selected muscles acting on talocrural and subtalar joints during movement: an in vitro study. *J Biomech.* 1996;29(1):21-30.
16. Heckman DS, Reddy S, Pedowitz D, Wapner KL, Parekh SG. Operative treatment for peroneal tendon disorders. *J Bone Joint Surg Am.* 2008;90(2):404-418. doi:10.2106/JBJS.G.00965.
17. Philbin TM, Landis GS, Smith B.. Peroneal tendon injuries. *J Am Acad of Orthop Surg.* 2009;17(5):306-317.
18. Mizel MS, Temple HT, Scranton PE Jr, et al. Role of the peroneal tendons in the production of the deformed foot with posterior tibial tendon deficiency. *Foot Ankle Int.* 1999;20(5):285-289.

19. Bloome DM, Marymont JV, Varner KE. Variations on the insertion of the posterior tibialis tendon: a cadaveric study. *Foot Ankle Int.* 2003;24(10):780-783.
20. Flemister AS, Neville CG, Houck J. The relationship between ankle, hindfoot, and forefoot position and posterior tibial muscle excursion. *Foot Ankle Int.* 2007;28(4):448-455.
21. Otis JC, Gage T. Function of the posterior tibial tendon muscle. *Foot Ankle Clin.* 2001;6(1):1-14, v.
22. Houck JR, Nomides C, Neville CG, Samuel Flemister A. The effect of Stage II posterior tibial tendon dysfunction on deep compartment muscle strength: a new strength test. *Foot Ankle Int.* 2008;29(9):895-902. doi:10.3113/FAI.2008.0895
23. Niki H, Ching RP, Kiser P, Sangeorzan BJ. The effect of posterior tibial tendon dysfunction on hindfoot kinematics. *Foot Ankle Int.* 2001;22(4):292-300.
24. Sharkey NA, Ferris L, Donahue SW. Biomechanical consequences of plantar fascial release or rupture during gait: part I—disruptions in longitudinal arch conformation. *Foot Ankle Int.* 1998;19(12):812-820.
25. Thordarson DB, Schmotzer H, Choi J, Peters J. Dynamic support of the human longitudinal arch. A biomechanical evaluation. *Clin Orthop Relat Res.* 1995;(316):165-172.
26. Kulig K, Burnfield JM, Reischl S, Requejo SM, Blanco CE, Thordarson DB. Effect of foot orthoses on tibialis posterior activation in persons with pes planus. *Med Sci Sports Exerc.* 2005;37(1):24-29.
27. Kulig K, Burnfield JM, Requejo SM, Sperry M, Terk M. Selective activation of tibialis posterior: evaluation by magnetic resonance imaging. *Med Sci Sports Exerc.* 2004;36(5):862-867.
28. Allen RH, Gross MT. Toe flexors strength and passive extension range of motion of the first metatarsophalangeal joint in individuals with plantar fasciitis. *J Orthop Sports Phys Ther.* 2003;33(8):468-478.
29. Lee SS, Piazza SJ. Inversion-eversion moment arms of gastrocnemius and tibialis anterior measured in vivo. *J Biomech.* 2008;41(16):3366-3370. doi:10.1016/j.jbiomech.2008.09.029.
30. Basmajian JV, Stecko G. The role of muscles in archy support of the foot. *J Bone Joint Surg Am.* 1963;45:1184-1190.
31. Mann R, Inman VT. Phasic activity of intrinsic muscles of the foot. *J Bone Joint Surg Am.* 1964;46:469-481.
32. Fiolkowski P, Brunt D, Bishop M, Woo R, Horodyski M. Intrinsic pedal musculature support of the medial longitudinal arch: an electromyography study. *J Foot Ankle Surg.* 2003;42(6):327-333.
33. Headlee DL, Leonard JL, Hart JM, Ingersoll CD, Hertel J. Fatigue of the plantar intrinsic foot muscles increases navicular drop. *J Electromyogr Kinesiol.* 2008;18(3):420-425.
34. Jung DY, Koh EK, Kwon OY, Yi CH, Oh JS, Weon JH. Effect of medial arch support on displacement of the myotendinous junction of the gastrocnemius during standing wall stretching. *J Orthop Sports Phys Ther.* 2009;39(12):867-874. doi:10.2519/jospt.2009.3158.
35. Kurihara T, Yamauchi J, Otsuka M, Tottori N, Hashimoto T, Isaka T. Maximum toe flexor muscle strength and quantitative analysis of human plantar intrinsic and extrinsic muscles by a magnetic resonance imaging technique. *J Foot Ankle Res.* 2014;7:26. doi:10.1186/1757-1146-7-26.
36. Kura H, Luo ZP, Kitaoka HB, An KN. Quantitative analysis of the intrinsic muscles of the foot. *Anat Rec.* 1997;249(1):143-151.
37. Arinci Incel N, Genc H, Erdem HR, Yorgancioglu ZR. Muscle imbalance in hallux valgus: an electromyographic study. *Am J Phys Med Rehabil.* 2003;82(5):345-349.
38. Arndt A, Westblad P, Winson I, Hashimoto T, Lundberg A. Ankle and subtalar kinematics measured with intracortical pins during the stance phase of walking. *Foot Ankle Int.* 2004;25(5):357-364.
39. Lundberg A, Svensson OK, Bylund C, Goldie I, Selvik G. Kinematics of the ankle/foot complex--Part 2: Pronation and supination. *Foot Ankle.* 1989;9(5):248-253.
40. Imai K, Tokunaga D, Takatori R, et al. In vivo three-dimensional analysis of hindfoot kinematics. *Foot Ankle Int.* 2009;30(11):1094-1100. doi:10.3113/FAI.2009.1094.
41. Beumer A, Valstar ER, Garling EH, et al. Kinematics of the distal tibiofibular syndesmosis: radiostereometry in 11 normal ankles. *Acta Orthop Scand.* 2003;74(3):337-343.
42. Inman VT. Human locomotion. 1966. *Clin Orthop Relat Res.* 1993;(288):3-9.
43. Lewis GS, Cohen TL, Seisler AR, Kirby KA, Sheehan FT, Piazza SJ. In vivo tests of an improved method for functional location of the subtalar joint axis. *J Biomech.* 2009;42(2):146-151. doi:10.1016/j.jbiomech.2008.10.010.
44. Kirby KA. Subtalar joint axis location and rotational equilibrium theory of foot function. *J Am Podiatr Med Assoc.* 2001;91(9):465-487.
45. Piazza SJ. Mechanics of the subtalar joint and its function during walking. *Foot Ankle Clin.* 2005;10(3):425-442, v.
46. Ledoux WR, Sangeorzan BJ. Clinical biomechanics of the peritalar joint. *Foot Ankle Clin.* 2004;9(4):663-683.
47. Blackwood CB, Yuen TJ, Sangeorzan BJ, Ledoux WR. The midtarsal joint locking mechanism. *Foot Ankle Int.* 2005;26(12):1074-1080.
48. Manter JT. Movements of the subtalar and transverse tarsal joints. *Anat Rec.* 1941;80(4):397-410.
49. Okita N, Meyers SA, Challis JH, Sharkey NA. Middatarsal joint locking: New perspectives on an old paradigm. *J Orthop Res.* 2014;32(1):110-115. doi:10.1002/jor.22477.
50. Van Boerum DH, Sangeorzan BJ. Biomechanics and pathophysiology of flat foot. *Foot Ankle Clin.* 2003;8(3):419-430.
51. Carlson RE, Fleming LL, Hutton WC. The biomechanical relationship between the tendoachilles, plantar fascia and metatarsophalangeal joint dorsiflexion angle. *Foot Ankle Int.* 2000;21(1):18-25.
52. Erdemir A, Hamel AJ, Fauth AR, Piazza SJ, Sharkey NA. Dynamic loading of the plantar aponeurosis in walking. *J Bone Joint Surg Am.* 2004;86-A(3):546-552.
53. Kappel-Bargas A, Woolf RD, Cornwall MW, McPoil TG. The windlass mechanism during normal walking and passive first metatarsalphalangeal joint extension. *Clin Biomech (Bristol, Avon).* 1998;13(3):190-194.
54. Nawoczenski DA, Baumhauer JF, Umberger BR. Relationship between clinical measurements and motion of the first metatarsophalangeal joint during gait. *J Bone Joint Surg Am.* 1999;81(3):370-376.
55. Kelly LA, Cresswell AG, Racinais S, Whiteley R, Lichtwark G. Intrinsic foot muscles have the capacity to control deformation of the longitudinal arch. *J R Soc Interface.* 2014;11(93):20131188. doi:10.1098/rsif.2013.1188.
56. Gatens PF, Saeed MA. Electromyographic findings in the intrinsic muscles of normal feet. *Arch Phys Med Rehabil.* 1982;63(7):317-318.
57. Medina McKeon JM, Bush HM, Reed A, Whittington A, Uhl TL, McKeon PO. Return-to-play probabilities following new versus recurrent ankle sprains in high school athletes. *J Sci Med Sport.* 2014;17(1):23-28. doi:10.1016/j.jsmams.2013.04.006.
58. Blackman AJ, Blevins JJ, Sangeorzan BJ, Ledoux WR. Cadaveric flatfoot model: ligament attenuation and Achilles tendon overpull. *J Orthop Res.* 2009;27(12):1547-1554. doi:10.1002/jor.20930.
59. Lichtwark GA, Wilson AM. In vivo mechanical properties of the human Achilles tendon during one-legged hopping. *J Exp Biol.* 2005;208(Pt 24):4715-4725.
60. Nawoczenski DA, Saltzman CL, Cook TM. The effect of foot structure on the three-dimensional kinematic coupling behavior of the leg and rear foot. *Phys Ther.* 1998;78(4):404-416.
61. Tiberio D. Relationship between foot pronation and rotation of the tibia and femur during walking. *Foot Ankle Int.* 2000;21(12):1057-1060.
62. Reischl SF, Powers CM, Rao S, Perry J. Relationship between foot pronation and rotation of the tibia and femur during walking. *Foot Ankle Int.* 1999;20(8):513-520.
63. Gross MT. Lower quarter screening for skeletal malalignment--suggestions for orthotics and shoewear. *J Orthop Sports Phys Ther.* 1995;21(6):389-405.
64. Heiderscheit B, Hamill J, Tiberio D. A Biomechanical Perspective: do foot orthoses work? *Br J Sports Med.* 2001;35(1):4-5.
65. Bellchamber TL, van den Bogert AJ. Contributions of proximal and distal moments to axial tibial rotation during walking and running. *J Biomed.* 2000;33(11):1397-1403.
66. Snyder KR, Earl JE, O'Connor KM, Ebersole KT. Resistance training is accompanied by increases in hip strength and changes in lower extremity biomechanics during running. *Clin Biomech (Bristol, Avon).* 2009;24(1):26-34. doi:10.1016/j.clinbiomech.2008.09.009.
67. Shaffer MA, Okereke E, Esterhai JL Jr, et al. Effects of immobilization on plantar-flexion torque, fatigue resistance, and functional ability following an ankle fracture. *Phys Ther.* 2000;80(8):769-780.
68. Brodsky JV. Preliminary gait analysis results after posterior tibial tendon reconstruction: a prospective study. *Foot Ankle Int.* 2004;25(2):96-100.
69. Winter DA, Patla AE, Frank JS, Walt SE. Biomechanical walking pattern changes in the fit and healthy elderly. *Phys Ther.* 1990;70(6):340-347.
70. Lewis CL, Ferris DP. Walking with increased ankle pushoff decreases hip muscle moments. *J Biomed.* 2008;41(10):2082-2089. doi:10.1016/j.jbiomech.2008.05.013.
71. Houck JR, Neville CG, Tome J, Flemister AS. Ankle and foot kinematics associated with stage II PTTD during stance. *Foot Ankle Int.* 2009;30(6):530-539. doi:10.3113/FAI.2009.0530.
72. Leardini A, Benedetti MG, Berti L, Bettinelli D, Nativo R, Giannini S. Rearfoot, mid-foot and fore-foot motion during the stance phase of gait. *Gait Posture.* 2007;25(3):453-462.
73. McPoil T, Cornwall MW. Relationship between neutral subtalar joint position and pattern of rearfoot motion during walking. *Foot Ankle Int.* 1994;15(3):141-145.
74. Ringleb SI, Kavros SJ, Kotajarvi BR, Hansen DK, Kitaoka HB, Kaufman KR. Changes in gait associated with acute stage II posterior tibial tendon dysfunction. *Gait Posture.* 2007;25(4):555-564.
75. Chang R, Van Emmerik R, Hamill J. Quantifying rearfoot-forefoot coordination in human walking. *J Biomed.* 2008;41(14):3101-3105. doi:10.1016/j.jbiomech.2008.07.024.
76. Neville C, Flemister AS, Houck JR. Effects of the AirLift PTTD brace on foot kinematics in subjects with stage II posterior tibial tendon dysfunction. *J Orthop Sports Phys Ther.* 2009;39(3):201-209. doi:10.2519/jospt.2009.2908.
77. Neville CG, Houck JR, Flemister AS. Science behind use of orthotic devices to manage posterior tibial tendon dysfunction. *Tech Foot Ankle Surg.* 2008;7(2):125-133. doi:10.1097/BTF.0b013e3181757571
78. Houck JR, Tome JM, Nawoczenski DA. Subtalar neutral position as an offset for a kinematic model of the foot during walking. *Gait Posture.* 2008;28(1):29-37.
79. Tome J, Flemister A, Nawoczenski DA, Houck J. Comparison of foot kinematics between subjects with posterior tibial tendon dysfunction and healthy controls. *J Orthop Sports Phys Ther.* 2006;36(9):635-644.
80. Youberg LD, Cornwall MW, McPoil TG, Hannon PR. The amount of rearfoot motion used during the stance phase of walking. *J Am Podiatr Med Assoc.* 2005;95(4):376-382.
81. Cornwall MW, McPoil TG. Classification of frontal plane rearfoot motion patterns during the stance phase of walking. *J Am Podiatr Med Assoc.* 2009;99(5):399-405.
82. Hansen ML, Otis JC, Kenneally SM, Deland JT. A closed-loop cadaveric foot and ankle loading model. *J Biomed.* 2001;34(4):551-555.
83. Rao S, Baumhauer JF, Tome J, Nawoczenski DA. Comparison of in vivo segmental foot motion during walking and step descent in patients with midfoot arthritis and matched asymptomatic control subjects. *J Biomed.* 2009;42(8):1054-1060. doi:10.1016/j.jbiomech.2009.02.006.
84. Houck JR, Neville C, Tome J, Flemister AS. Foot kinematics during a bilateral heel rise test in participants with stage II posterior tibial tendon dysfunction. *J Orthop Sports Phys Ther.* 2009;39(8):593-603. doi:10.2519/jospt.2009.3040.

85. Nawoczenski DA, Ketz J, Baumhauer JF. Dynamic kinematic and plantar pressure changes following cheilectomy for hallux rigidus: a mid-term followup. *Foot Ankle Int.* 2008;29(3):265-272. doi:10.3113/FAI.2008.0265.
86. Stefanyshyn DJ, Nigg BM. Mechanical energy contribution of the metatarsophalangeal joint to running and sprinting. *J Biomech.* 1997;30(11-12):1081-1085.
87. Kirane YM, Michelson JD, Sharkey NA. Evidence of isometric function of the flexor hallucis longus muscle in normal gait. *J Biomech.* 2008;41(9):1919-1928. doi:10.1016/j.jbiomech.2008.03.040.
88. Ledoux WR, Shofer JB, Ahroni JH, Smith DG, Sangeorzan BJ, Boyko EJ. Biomechanical differences among pes cavus, neutrally aligned, and pes planus feet in subjects with diabetes. *Foot Ankle Int.* 2003;24(11):845-850.
89. Donovan A, Rosenberg ZS. Extraarticular lateral hindfoot impingement with posterior tibial tendon tear: MRI correlation. *AJR Am J Roentgenol.* 2009;193(3):672-678. doi:10.2214/AJR.08.2215.
90. Malicky ES, Cray JL, Houghton MJ, Agel J, Hansen ST Jr, Sangeorzan BJ. Talocalcaneal and subtalar impingement in symptomatic flatfoot in adults. *J Bone Joint Surg Am.* 2002;84-A(11):2005-2009.
91. Butler RJ, Hillstrom H, Song J, Richards CJ, Davis IS. Arch height index measurement system: establishment of reliability and normative values. *J Am Podiatr Med Assoc.* 2008;98(2):102-106.
92. Saltzman CL, Brandser EA, Berbaum KS, et al. Reliability of standard foot radiographic measurements. *Foot Ankle Int.* 1994;15(12):661-665.
93. Saltzman CL, Nawoczenski DA, Talbot KD. Measurement of the medial longitudinal arch. *Arch Phys Med Rehabil.* 1995;76(1):45-49.
94. Williams DS, McClay IS. Measurements used to characterize the foot and the medial longitudinal arch: reliability and validity. *Phys Ther.* 2000;80(9):864-871.
95. Younger AS, Sawatzky B, Dryden P. Radiographic assessment of adult flatfoot. *Foot Ankle Int.* 2005;26(10):820-825.
96. Ness ME, Long J, Marks R, Harris G. Foot and ankle kinematics in patients with posterior tibial tendon dysfunction. *Gait Posture.* 2008;27(2):331-339.
97. Neville C, Flemister AS, Houck JR. Deep posterior compartment strength and foot kinematics in subjects with stage II posterior tibial tendon dysfunction. *Foot Ankle Int.* 2010;31(4):320-328. doi:10.3113/FAI.2010.0320.
98. McPoil TG, Cornwall MW. Use of the longitudinal arch angle to predict dynamic foot posture in walking. *J Am Podiatr Med Assoc.* 2005;95(2):114-120.
99. Redmond AC, Crosbie J, Ouvrier RA. Development and validation of a novel rating system for scoring standing foot posture: the Foot Posture Index. *Clin Biomech (Bristol, Avon).* 2006;21(1):89-98.
100. Queen RM, Mall NA, Hardaker WM, Nunley JA 2nd. Describing the medial longitudinal arch using footprint indices and a clinical grading system. *Foot Ankle Int.* 2007;28(4):456-462.
101. Butler RJ, Davis IS, Hamill J. Interaction of arch type and footwear on running mechanics. *Am J Sports Med.* 2006;34(12):1998-2005.
102. Michlitsch MG, Rethlefsen SA, Kay RM. The contributions of anterior and posterior tibialis dysfunction to varus foot deformity in patients with cerebral palsy. *J Bone Joint Surg Am.* 2006;88(8):1764-1768.
103. Daines SB, Rohr ES, Pace AP, Fassbind MJ, Sangeorzan BJ, Ledoux WR. Cadaveric simulation of a pes cavus foot. *Foot Ankle Int.* 2009;30(1):44-50. doi:10.3113/FAI.2009.0044
104. Manoli A 2nd, Graham B. The subtle cavus foot, "the underpronator." *Foot Ankle Int.* 2005;26(3):256-263.
105. Burns J, Landorf KB, Ryan MM, Crosbie J, Ouvrier RA. Interventions for the prevention and treatment of pes cavus. *Cochrane Database Syst Rev.* 2007;(4):CD006154.
106. Williams DS 3rd, McClay IS, Hamill J. Arch structure and injury patterns in runners. *Clin Biomech (Bristol, Avon).* 2001;16(4):341-347.
107. Burns J, Crosbie J, Ouvrier R, Hunt A. Effective orthotic therapy for the painful cavus foot: a randomized controlled trial. *J Am Podiatr Med Assoc.* 2006;96(3):205-211.
108. Badlissi F, Dunn JE, Link CL, Keysor JJ, McKinlay JB, Felson DT. Foot musculoskeletal disorders, pain, and foot-related functional limitation in older persons. *J Am Geriatr Soc.* 2005;53(6):1029-1033.
109. Benvenuti F, Ferrucci L, Guralnik JM, Gangemi S, Baroni A. Foot pain and disability in older persons: an epidemiologic survey. *J Am Geriatr Soc.* 1995;43(5):479-484.
110. Rao S, Ellis SJ, Deland JT, Hillstrom H. Nonmedicinal therapy in the management of ankle arthritis. *Curr Opin Rheumatol.* 2010;22(2):223-228. doi:10.1097/BOR.0b013e328335fc2b.
111. Hung M, Franklin JD, Hon SD, Cheng C, Conrad J, Saltzman CL. Time for a paradigm shift with computerized adaptive testing of general physical function outcomes measurements. *Foot Ankle Int.* 2014;35(1):1-7. doi:10.1177/1071100713507905.
112. Robinson JM, Cook JL, Purdam C, et al. The VISA-A questionnaire: a valid and reliable index of the clinical severity of Achilles tendinopathy. *Br J Sports Med.* 2001;35(5):335-341.
113. Martin RL, Irrgang JJ, Burdett RG, Conti SF, Van Swearingen JM. Evidence of validity for the Foot and Ankle Ability Measure (FAAM). *Foot Ankle Int.* 2005;26(11):968-983.
114. Carcia CR, Martin RL, Houck J, Wukich DK; Orthopaedic Section of the American Physical Therapy Association. Achilles pain, stiffness, and muscle power deficits: achilles tendinitis. *J Orthop Sports Phys Ther.* 2010;40(9):A1-26. doi:10.2519/jospt.2010.0305
115. McPoil TG, Martin RL, Cornwall MW, Wukich DK, Irrgang JJ, Godges JJ. Heel pain-plantar fasciitis: clinical practice guidelines linked to the international classification of function, disability, and health from the orthopaedic section of the American Physical Therapy Association. *J Orthop Sports Phys Ther.* 2008;38(4):A1-A18. doi:10.2519/jospt.2008.0302.
116. Eequeute C, Vaes P, Van Aerschot L, Asman S, Duquest W. The clinimetric qualities of patient-assessed instruments for measuring chronic ankle instability: a systematic review. *BMC Musculoskelet Disord.* 2007;8:6.
117. Martin RL, Irrgang JJ. A survey of self-reported outcome instruments for the foot and ankle. *J Orthop Sports Phys Ther.* 2007;37(2):72-84.
118. Gabel CP, Mellilo M, Burkett B, Michener LA. Lower limb functional index: development and clinimetric properties. *Phys Ther.* 2012;92(1):98-110. doi:10.2522/ptj.20100199.
119. Martin RL, Irrgang JJ, Lalonde KA, Conti S. Current concepts review: foot and ankle outcome instruments. *Foot Ankle Int.* 2006;27(5):383-390.
120. Hiller CE, Reishauge KM, Bundy AC, Herbert RD, Kilbreath SL. The Cumberland ankle instability tool: a report of validity and reliability testing. *Arch Phys Med Rehabil.* 2006;87(9):1235-1241.
121. Docherty CL, Gansneder BM, Arnold BL, Hurwitz SR. Development and reliability of the ankle instability instrument. *J Athl Train.* 2006;41(2):154-158.
122. Keenan AM, Redmond AC, Horton M, Conaghan PG, Tennant A. The Foot Posture Index: Rasch analysis of a novel, foot-specific outcome measure. *Arch Phys Med Rehabil.* 2007;88(1):88-93.
123. McPoil TG, Cornwall MW, Medoff L, Vicenzino B, Forsberg K, Hilz D. Arch height change during sit-to-stand: an alternative for the navicular drop test. *J Foot Ankle Res.* 2008;1(1):3. doi:10.1186/1757-1146-1-3.
124. Teyhen DS, Stoltzenberg BE, Eckard TG, et al. Static foot posture associated with dynamic plantar pressure parameters. *J Orthop Sports Phys Ther.* 2011;41(2):100-107. doi:10.2519/jospt.2011.3412.
125. McPoil TG, Vicenzino B, Cornwall MW, Collins N, Warren M. Reliability and normative values for the foot mobility magnitude: a composite measure of vertical and medial-lateral mobility of the midfoot. *J Foot Ankle Res.* 2009;2:6. doi:10.1186/1757-1146-2-6.
126. Cornwall MW, McPoil TG. Relationship between static foot posture and foot mobility. *J Foot Ankle Res.* 2011;4:4. doi:10.1186/1757-1146-4-4.
127. Brody DM. Techniques in the evaluation and treatment of the injured runner. *Orthop Clin North Am.* 1982;13(3):541-558.
128. Mueller MJ, Host JV, Norton BJ. Navicular drop as a composite measure of excessive pronation. *J Am Podiatr Med Assoc.* 1993;83(4):198-202.
129. Buist I, Bredeweg SW, Lemmink KA, van Mechelen W, Diercks RL. Predictors of running-related injuries in novice runners enrolled in a systematic training program: a prospective cohort study. *Am J Sports Med.* 2010;38(2):273-280. doi:10.1177/0363546509347985.
130. Barton CJ, Bonanno D, Levinger P, Menz HB. Foot and ankle characteristics in patellofemoral pain syndrome: a case control and reliability study. *J Orthop Sports Phys Ther.* 2010;40(5):286-296. doi:10.2519/jospt.2010.3227.
131. Karatsolis K, Nikolopoulos CS, Papadopoulos ES, Vagenas G, Terzis E, Athanasiopoulos S. Eversion and inversion muscle group peak torque in hyperpronated and normal individuals. *Foot (Edinb).* 2009;19(1):29-35. doi:10.1016/j.foo.2008.06.006.
132. Boling MC, Padua DA, Marshall SW, Guskiewicz K, Pyne S, Beutler A. A prospective investigation of biomechanical risk factors for patellofemoral pain syndrome: the Joint Undertaking to Monitor and Prevent ACL Injury (JUMP-ACL) cohort. *Am J Sports Med.* 2009;37(11):2108-2116. doi:10.1177/0363546509337934.
133. Vauhnik R, Morrissey MC, Rutherford OM, Turk Z, Pilih IA, Perme MP. Correlates of knee anterior laxity in sportswomen. *Knee.* 2009;16(6):427-431. doi:10.1016/j.knee.2009.04.001.
134. Reinking MF, Austin TM, Hayes AM. Exercise-related leg pain in collegiate cross-country athletes: extrinsic and intrinsic risk factors. *J Orthop Sports Phys Ther.* 2007;37(11):670-678.
135. Jenkins WL, Killian CB, Williams DS 3rd, Loudon J, Raedeke SG. Anterior cruciate ligament injury in female and male athletes: the relationship between foot structure and injury. *J Am Podiatr Med Assoc.* 2007;97(5):371-376.
136. Riemann BL, Guskiewicz KM. Effects of mild head injury on postural stability as measured through clinical balance testing. *J Athl Train.* 2000;35(1):19-25.
137. Springer BA, Marin R, Cyhan T, Roberts H, Gill NW. Normative values for the unipedal stance test with eyes open and closed. *J Geriatr Phys Ther.* 2007;30(1):8-15.
138. Hertel J, Miller SJ, Denegar CR. Intratester and intertester reliability during the star excursion balance tests. *J Sport Rehabil.* 2000;9:104-116.
139. Hébert-Losier K, Schneiders AG, Newsham-West RJ, Sullivan SJ. Scientific bases and clinical utilisation of the calf-raise test. *Phys Ther Sport.* 2009;10(4):142-149. doi:10.1016/j.ptsp.2009.07.001.
140. Madeley LT, Muntean SE, Bonanno DR. Endurance of the ankle joint plantar flexor muscles in athletes with medial tibial stress syndrome: a case-control study. *J Sci Med Sport.* 2007;10(6):356-362.
141. Olsson N, Karlsson J, Eriksson Bl, Brorsson A, Lundberg M, Silbernagel KG. Ability to perform a single heel-rise is significantly related to patient-reported outcome after Achilles tendon rupture. *Scand J Med Sci Sports.* 2014;24(1):152-158. doi:10.1111/j.1600-0838.2012.01497.x.
142. Silbernagel KG, Brorsson A, Lundberg M. The majority of patients with Achilles tendinopathy recover fully when treated with exercise alone: a 5-year follow-up. *Am J Sports Med.* 2011;39(3):607-613. doi:10.1177/0363546510384789.
143. Chimenti RL, Flemister AS, Tome J, et al. Altered tendon characteristics and mechanical properties associated with insertional Achilles tendinopathy. *J Orthop Sports Phys Ther.* 2014;44(9):680-689. doi:10.2519/jospt.2014.5369.
144. Segura-Ortí E, Martínez-Olmos FJ. Test-retest reliability and minimal detectable change scores for sit-to-stand-to-sit tests, the six-minute walk test, the one-leg heel-rise test, and handgrip strength in people undergoing hemodialysis. *Phys Ther.* 2011;91(8):1244-1252. doi:10.2522/ptj.20100141.
145. Olsson N, Silbernagel KG, Eriksson Bl, et al. Stable surgical repair with accelerated rehabilitation versus nonsurgical treatment for acute Achilles tendon ruptures: a randomized controlled study. *Am J Sports Med.* 2013;41(12):2867-2876. doi:10.1177/0363546513503282.

146. Silbernagel KG, Nilsson-Helander K, Thomeé R, Eriksson BI, Karlsson J. A new measurement of heel-rise endurance with the ability to detect functional deficits in patients with Achilles tendon rupture. *Knee Surg Sports Traumatol Arthrosc.* 2010;18(2):258-264. doi:10.1007/s00167-009-0889-7.
147. Silbernagel KG, Steele R, Manal K. Deficits in heel-rise height and Achilles tendon elongation occur in patients recovering from an Achilles tendon rupture. *Am J Sports Med.* 2012;40(7):1564-1571. doi:10.1177/0363546512447926.
148. Sman AD, Hiller CE, Imer A, Ocsing A, Burns J, Refshauge KM. Design and reliability of a novel heel rise test measuring device for plantarflexion endurance. *BioMed Res Int.* 2014;2014:1-8. doi:10.1155/2014/391646.
149. Brunnekreef JJ, van Uden CJT, van Moorsel S, Kooloos JG. Reliability of videotaped observational gait analysis in patients with orthopedic impairments. *BMC Musculoskelet Disord.* 2005;6:17.
150. Cornwall MW, McPoil TG, Fishco WD, Hunt L, Lane C, O'donnell D. Reliability of visual measurement of forefoot alignment. *Foot Ankle Int.* 2004;25(10):745-748.
151. Rathleff MS, Nielsen RG, Simonsen O, Olesen CG, Kersting UG. Perspectives for clinical measures of dynamic foot function-reference data and methodological considerations. *Gait Posture.* 2010;31(2):191-196. doi:10.1016/j.gaitpost.2009.10.004.
152. Haitz K, Shultz R, Hodgins M, Matheson GO. Test-retest and interrater reliability of the functional lower extremity evaluation. *J Orthop Sports Phys Ther.* 2014;44(12):947-954. doi:10.2519/jospt.2014.4809.
153. Martin RL, Davenport TE, Paulseth S, Wukich DK, Godges JJ; Orthopaedic Section American Physical Therapy Association. Ankle stability and movement coordination impairments: ankle ligament sprains. *J Orthop Sports Phys Ther.* 2013;43(9):A1-A40. doi:10.2519/jospt.2013.0305.
154. Caffrey E, Docherty CL, Schrader J, Klossner J. The ability of 4 single-limb hopping tests to detect functional performance deficits in individuals with functional ankle instability. *J Orthop Sports Phys Ther.* 2009;39(11):799-806. doi:10.2519/jospt.2009.3042.
155. Martin RL, McPoil TG. Reliability of ankle goniometric measurements: a literature review. *J Am Podiatr Med Assoc.* 2005;95(6):564-572.
156. Munteanu SE, Strawhorn AB, Landorf KB, Bird AR, Murley GS. A weightbearing technique for the measurement of ankle joint dorsiflexion with the knee extended is reliable. *J Sci Med Sport.* 2009;12(1):54-59.
157. Denegar CR, Hertel J, Fonseca J. The effect of lateral ankle sprain on dorsiflexion range of motion, posterior talar glide, and joint laxity. *J Orthop Sports Phys Ther.* 2002;32(4):166-173.
158. DiGiovanni CW, Kuo R, Tejwani N, et al. Isolated gastrocnemius tightness. *J Bone Joint Surg Am.* 2002;84-A(6):962-970.
159. DiGiovanni CW, Langer P. The role of isolated gastrocnemius and combined Achilles contractures in the flatfoot. *Foot Ankle Clin.* 2007;12(2):363-379, viii.
160. Root ML, Weed JH, Orien WP. *Normal and Abnormal Function of the Foot.* Vol 2. Los Angeles, CA: Clinical Biomechanics; 1977.
161. Piernowinski MR, Smith SB. Effect of patient position on the consistency of placing the rearfoot at subtalar neutral. *J Am Podiatr Med Assoc.* 1997;87(9):399-406.
162. Piernowinski MR, Smith SB, Mlynarczyk JH. Proficiency of foot care specialists to place the rearfoot at subtalar neutral. *J Am Podiatr Med Assoc.* 1996;86(5):217-223.
163. Sell KE, Verity TM, Worrell TW, Pease BJ, Wigglesworth J. Two measurement techniques for assessing subtalar joint position: a reliability study. *J Orthop Sports Phys Ther.* 1994;19(3):162-167.
164. Elvera RA, Rothstein JM, Lamb RL. Goniometric reliability in a clinical setting. Subtalar and ankle joint measurements. *Phys Ther.* 1988;68(5):672-677.
165. Astrom M, Arvidson T. Alignment and joint motion in the normal foot. *J Orthop Sports Phys Ther.* 1995;22(5):216-222.
166. Kwon OY, Tuttle LJ, Johnson JE, Mueller MJ. Muscle imbalance and reduced ankle joint motion in people with hammer toe deformity. *Clin Biomech (Bristol, Avon).* 2009;24(8):670-675. doi:10.1016/j.clinbiomech.2009.05.010.
167. Kovaleski JE, Norrell PM, Heitman RJ, Hollis JM, Pearsall AW. Knee and ankle position, anterior drawer laxity, and stiffness of the ankle complex. *J Athl Train.* 2008;43(3):242-248. doi:10.4085/1062-6050-43.3.242.
168. Kingwell S, Buckley R, Willis N. The association between subtalar joint motion and outcome satisfaction in patients with displaced intraarticular calcaneal fractures. *Foot Ankle Int.* 2004;25(9):666-673.
169. Lee MS, Vanore JV, Thomas JL, et al. Diagnosis and treatment of adult flatfoot. *J Foot Ankle Surg.* 2005;44(2):78-113.
170. Ogawa BK, Thordarson DB. Current concepts review: peroneal tendon subluxation and dislocation. *Foot Ankle Int.* 2007;28(9):1034-1040.
171. Aydog E, Aydog ST, Cakci A, Doral MN. Reliability of isokinetic ankle inversion- and eversion-strength measurement in neutral foot position, using the Biodes dynamometer. *Knee Surg Sports Traumatol Arthrosc.* 2004;12(5):478-481.
172. Mullaney MJ, McHugh MP, Tyler TF, Nicholas SJ, Lee SJ. Weakness in end-range plantar flexion after Achilles tendon repair. *Am J Sports Med.* 2006;34(7):1120-1125.
173. Webber SC, Porter MM. Effects of ankle power training on movement time in mobility-impaired older women. *Med Sci Sports Exerc.* 2010;42(7):1233-1240. doi:10.1249/MSS.0b013e3181cdd4e9.
174. Cleland JA, Abbott JH, Kidd MO, et al. Manual physical therapy and exercise versus electrophysical agents and exercise in the management of plantar heel pain: a multicenter randomized clinical trial. *J Orthop Sports Phys Ther.* 2009;39(8):573-585. doi:10.2519/jospt.2009.3036.
175. Voorn R. Case report: can sacroiliac joint dysfunction cause chronic Achilles tendinitis? *J Orthop Sports Phys Ther.* 1998;27(6):436-443.
176. Bahr R, Pena F, Shine J, et al. Mechanics of the anterior drawer and talar tilt tests. A cadaveric study of lateral ligament injuries of the ankle. *Acta Orthop Scand.* 1997;68(5):435-441.
177. Tohyama H, Yasuda K, Ohkoshi Y, Beynnon BD, Renstrom PA. Anterior drawer test for acute anterior talofibular ligament injuries of the ankle. How much load should be applied during the test? *Am J Sports Med.* 2003;31(2):226-232.
178. Phisitkul P, Chaichankul C, Sripongsoi R, Prasitdamrong I, Tengtrikalcharoen P, Suarchawaratana S. Accuracy of anterolateral drawer test in lateral ankle instability: a cadaveric study. *Foot Ankle Int.* 2009;30(7):690-695. doi:10.3113/FAI.2009.0690.
179. Fujii T, Luo ZP, Kitaka HB, An KN. The manual stress test may not be sufficient to differentiate ankle ligament injuries. *Clin Biomech (Bristol, Avon).* 2000;15(8):619-623.
180. Hubbard TJ, Hicks-Little CA. Ankle ligament healing after an acute ankle sprain: an evidence-based approach. *J Athl Train.* 2008;43(5):523-529. doi:10.4085/1062-6050-43.5.523.
181. Gaebler C, Kukla C, Breitenseher MJ, et al. Diagnosis of lateral ankle ligament injuries. Comparison between talar tilt, MRI and operative findings in 112 athletes. *Acta Orthop Scand.* 1997;68(3):286-290.
182. Sman AD, Hiller CE, Rae K, et al. Diagnostic accuracy of clinical tests for ankle syndesmosis injury. *Br J Sports Med.* 2015;49(5):323-329. doi:10.1136/bjsports-2013-092787.
183. Alonso A, Khouri L, Adams R. Clinical tests for ankle syndesmosis injury: reliability and prediction of return to function. *J Orthop Sports Phys Ther.* 1998;27(4):276-284.
184. Beumer A, Swierstra BA, Mulder PG. Clinical diagnosis of syndesmotic ankle instability: evaluation of stress tests behind the curtains. *Acta Orthop Scand.* 2002;73(6):667-669.
185. Beumer A, van Hemert WL, Swierstra BA, Jasper LE, Belkoff SM. A biomechanical evaluation of clinical stress tests for syndesmotic ankle instability. *Foot Ankle Int.* 2003;24(4):358-363.
186. Reiman M, Burgi C, Strube E, et al. The utility of clinical measures for the diagnosis of achilles tendon injuries: a systematic review with meta-analysis. *J Athl Train.* 2014;49(6):820-829. doi:10.4085/1062-6050-49.3.36.
187. Thompson TC, Doherty JH. Spontaneous rupture of tendon of Achilles: a new clinical diagnostic test. *J Trauma.* 1962;2:126-129.
188. Maffulli N. The clinical diagnosis of subcutaneous tear of the Achilles tendon. A prospective study in 174 patients. *Am J Sports Med.* 1998;26(2):266-270.
189. Safran MR, O'Malley DJ Jr, Fu FH. Peroneal tendon subluxation in athletes: new exam technique, case reports, and review. *Med Sci Sports Exerc.* 1999;31(7 Suppl):S487-492.
190. Coleman SS, Chesnut WJ. A simple test for hindfoot flexibility in the cavovarus foot. *Clin Orthop Relat Res.* 1977;123(123):60-62.
191. Mulder JD. The causative mechanism in morton's metatarsalgia. *J Bone Joint Surg Br.* 1951;33-B(1):94-95.
192. Torriani M, Kattapuram SV. Technical innovation. Dynamic sonography of the forefoot: The sonographic Mulder sign. *AJR Am J Roentgenol.* 2003;180(4):1121-1123.
193. Tinel J. "Tingling" signs with peripheral nerve injuries. 1915. *J Hand Surg Br.* 2005;30(1):87-89.
194. Gondring WH, Shields B, Wenger S. An outcomes analysis of surgical treatment of tarsal tunnel syndrome. *Foot Ankle Int.* 2003;24(7):545-550.
195. Riddle DL, Wells PS. Diagnosis of lower-extremity deep vein thrombosis in outpatients. *Phys Ther.* 2004;84(8):729-735.
196. Deyle GD. Musculoskeletal imaging in physical therapist practice. *J Orthop Sports Phys Ther.* 2005;35(11):708-721.
197. Campbell WC, Canale ST, Beaty JH. *Campbell's Operative Orthopaedics.* Vol 11th. Philadelphia, PA: Mosby/Elsevier; 2008.
198. Fowler A, Philip JF. Abnormality of the calcaneus as a cause of painful heel its diagnosis and operative treatment. *Br J Surg.* 1945;32(128):494-498.
199. Kang S, Thordarson DB, Charlton TP. Insertional Achilles tendinitis and Haglund's deformity. *Foot Ankle Int.* 2012;33(6):487-491. doi:10.3113/FAI.2012.0487.
200. Sundararajan PP, Wilde TS. Radiographic, clinical, and magnetic resonance imaging analysis of insertional Achilles tendinopathy. *J Foot Ankle Surg.* 2014;53(2):147-151. doi:10.1053/j.jfas.2013.12.009.
201. Whittaker JL, Teyhen DS, Elliott JM, et al. Rehabilitative ultrasound imaging: understanding the technology and its applications. *J Orthop Sports Phys Ther.* 2007;37(8):434-449.
202. Bianchi S, Martinoli C, Gaignot C, et al. Ultrasound of the ankle: anatomy of the tendons, bursae, and ligaments. *Semin Musculoskelet Radiol.* 2005;9(3):243-259.
203. Pfeffer G, Bacchetti P, Deland J, et al. Comparison of custom and prefabricated orthoses in the initial treatment of proximal plantar fasciitis. *Foot Ankle Int.* 1999;20(4):214-221.
204. Staheli LT. Evaluation of planovalgus foot deformities with special reference to the natural history. *J Am Podiatr Med Assoc.* 1987;77(1):2-6.
205. Mickle KJ, Munro BJ, Lord SR, Menz HB, Steele JR. ISB Clinical Biomechanics Award 2009: toe weakness and deformity increase the risk of falls in older people. *Clin Biomech (Bristol, Avon).* 2009;24(10):787-791. doi:10.1016/j.clinbiomech.2009.08.011.
206. Rao UB, Joseph B. The influence of footwear on the prevalence of flat foot. A survey of 2300 children. *J Bone Joint Surg Br.* 1992;74(4):525-527.
207. Sachithanandam V, Joseph B. The influence of footwear on the prevalence of flat foot. A survey of 1846 skeletally mature persons. *J Bone Joint Surg Br.* 1995;77(2):254-257.
208. Ekinici S, Kocak N, Aydin I, et al. Prevalence and distribution of musculoskeletal disorders causing unfitness for military service among young adult men: An epidemiologic study. *Pak J Med Sci.* 2015;31(1):43-48. doi:10.12669/pjms.311.5674.
209. Gumundsson B, Olafsson E, Jakobsson F, Luvigsson P. Prevalence of symptomatic Charcot-Marie-Tooth disease in Iceland: A study of a well-defined population. *Neuroepidemiology.* 2010;34(1):13-17. doi:10.1159/000255461.

210. Khas KS, Pandey PM, Ray AR. Design and development of a device to measure the deformities of clubfoot. *Proc Inst Mech Eng H*. 2015;229(3):194-204. doi:10.1177/0954411915574758.
211. Butler RJ, Hamill J, Davis I. Effect of footwear on high and low arched runners' mechanics during a prolonged run. *Gait Posture*. 2007;26(2):219-225.
212. Lane TJ, Landorf KB, Bonanno DR, Raspovic A, Menz HB. Effects of shoe sole hardness on plantar pressure and comfort in older people with forefoot pain. *Gait Posture*. 2014;39(1):247-251. doi:10.1016/j.gaitpost.2013.07.116.
213. Lucas-Cuevas AG, Perez-Soriano P, Priego-Quesada JI, Llana-Belloch S. Influence of foot orthosis customisation on perceived comfort during running. *Ergonomics*. 2014;57(10):1590-1596. doi:10.1080/00140139.2014.938129.
214. Nix S, Smith M, Vicenzino B. Prevalence of hallux valgus in the general population: a systematic review and meta-analysis. *J Foot Ankle Res*. 2010;3:21. doi:10.1186/1757-1146-3-21.
215. Coughlin MJ, Freund E, Roger A. Mann Award . The reliability of angular measurements in hallux valgus deformities. *Foot Ankle Int*. 2001;22(5):369-379.
216. Coughlin MJ, Shurnas PS. Hallux valgus in men. Part II: First ray mobility after bunionectomy and factors associated with hallux valgus deformity. *Foot Ankle Int*. 2003;24(1):73-78.
217. Hardy RH, Clapham JC. Hallux valgus; predisposing anatomical causes. *Lancet*. 1952;1(6720):1180-1183.
218. Nguyen US, Hillstrom HJ, Li W, et al. Factors associated with hallux valgus in a population-based study of older women and men: the MOBILIZE Boston Study. *Osteoarthritis Cartilage*. 2010;18(1):41-46. doi:10.1016/j.joca.2009.07.008.
219. Menz HB, Lord SR. Gait instability in older people with hallux valgus. *Foot Ankle Int*. 2005;26(6):483-489.
220. Scranton PE Jr, McDermott JE. Prognostic factors in bunion surgery. *Foot Ankle Int*. 1995;16(11):698-704.
221. Sanders AP, Snijders CJ, van Linge B. Medial deviation of the first metatarsal head as a result of flexion forces in hallux valgus. *Foot Ankle*. 1992;13(9):515-522.
222. Hurn SE, Vicenzino B, Smith MD. Functional impairments characterizing mild, moderate, and severe hallux valgus. *Arthritis Care Res*. 2015;67(1):80-88. doi:10.1002/acr.22380.
223. Glasow WM, Allen MK, Saltzman CL. First ray dorsal mobility in relation to hallux valgus deformity and first intermetatarsal angle. *Foot Ankle Int*. 2001;22(2):98-101.
224. Glasow WM, Nuckley DJ, Ludewig PM. Hallux valgus and the first metatarsal arch segment: a theoretical biomechanical perspective. *Phys Ther*. 2010;90(1):110-120. doi:10.2522/ptj.20080298.
225. Torkki M, Malmivaara A, Seitsalo S, Hoikka V, Laippala P, Paavolainen P. Hallux valgus: immediate operation versus 1 year of waiting with or without orthoses: a randomized controlled trial of 209 patients. *Acta Orthop Scand*. 2003;74(2):209-215.
226. Torkki M, Malmivaara A, Seitsalo S, Hoikka V, Laippala P, Paavolainen P. Surgery vs orthosis vs watchful waiting for hallux valgus: a randomized controlled trial. *JAMA*. 2001;285(19):2474-2480.
227. Ferrari J, Malone-Lee J. The shape of the metatarsal head as a cause of hallux abductovalgus. *Foot Ankle Int*. 2002;23(3):236-242.
228. Chong A, Nazarian N, Chandrananth J, Tacey M, Shepherd D, Tran P. Surgery for the correction of hallux valgus: minimum five-year results with a validated patient-reported outcome tool and regression analysis. *Bone Joint J*. 2015;97-B(2):208-214. doi:10.1302/0301-620X.97B2.34891.
229. Yucel I, Tenekecioglu Y, O gut K, Kesmezacar H. Treatment of hallux valgus by modified McBride procedure: a 6-year follow-up. *J Orthop Traumatol*. 2010;11(2):89-97. doi:10.1007/s10195-010-0092-0.
230. Schuh R, Hofstaetter SG, Adams SB Jr, Pichler F, Kristen KH, Trnka HJ. Rehabilitation after hallux valgus surgery: importance of physical therapy to restore weight bearing of the first ray during the stance phase. *Phys Ther*. 2009;89(9):934-945. doi:10.2522/ptj.20080375.
231. Kernozenk TW, Sterriker SA. Chevron (Austin) distal metatarsal osteotomy for hallux valgus: comparison of pre- and post-surgical characteristics. *Foot Ankle Int*. 2002;23(6):503-508.
232. Williams GN, Jones MH, Amendola A. Syndesmotic ankle sprains in athletes. *Am J Sports Med*. 2007;35(7):1197-1207.
233. Lin CF, Gross ML, Weinhold P. Ankle syndesmosis injuries: anatomy, biomechanics, mechanism of injury, and clinical guidelines for diagnosis and intervention. *J Orthop Sports Phys Ther*. 2006;36(6):372-384.
234. Press CM, Gupta A, Hutchinson MR. Management of ankle syndesmosis injuries in the athlete. *Curr Sports Med Rep*. 2009;8(5):228-233. doi:10.1249/JSM.0b013e3181b7ec0c.
235. Han SH, Lee JW, Kim S, Suh JS, Choi YR. Chronic tibiofibular syndesmosis injury: the diagnostic efficiency of magnetic resonance imaging and comparative analysis of operative treatment. *Foot Ankle Int*. 2007;28(3):336-342.
236. Taylor DC, Tenuta JJ, Uhorchak JM, Arciero RA. Aggressive surgical treatment and early return to sports in athletes with grade III syndesmosis sprains. *Am J Sports Med*. 2007;35(11):1833-1838.
237. Miller TL, Skalak T. Evaluation and treatment recommendations for acute injuries to the ankle syndesmosis without associated fracture. *Sports Med*. 2014;44(2):179-188. doi:10.1007/s40279-013-0106-1.
238. Feger MA, Herb CC, Fraser JJ, Glaviano N, Hertel J. Supervised rehabilitation versus home exercise in the treatment of acute ankle sprains: a systematic review. *Clin Sports Med*. 2015;34(2):329-346. doi:10.1016/j.csm.2014.12.001.
239. Hubbard TJ, Cordova M. Mechanical instability after an acute lateral ankle sprain. *Arch Phys Med Rehabil*. 2009;90(7):1142-1146. doi:10.1016/j.apmr.2009.01.020.
240. van Rijn RM, van Os AG, Bernsen RM, Luijsterburg PA, Koes BW, Bierma-Zeinstra SM. What is the clinical course of acute ankle sprains? A systematic literature review. *Am J Med*. 2008;121(4):324-331.e326. doi:10.1016/j.amjmed.2007.11.018.
241. Sugimoto K, Takakura Y, Okahashi K, Samoto N, Kawate K, Iwai M. Chondral injuries of the ankle with recurrent lateral instability: an arthroscopic study. *J Bone Joint Surg Am*. 2009;91(1):99-106. doi:10.2106/JBJS.G.00087.
242. Doherty C, Delahunt E, Caulfield B, Hertel J, Ryan J, Bleakley C. The incidence and prevalence of ankle sprain injury: a systematic review and meta-analysis of prospective epidemiological studies. *Sports Med*. 2014;44(1):123-140. doi:10.1007/s40279-013-0102-5.
243. Swenson DMC, Christy L, Fields SKC, Dawn R. Injuries, ligamentous ankle. *NIH Public Access*. 2014;23(3):190-196.
244. Lambers K, Ootes D, Ring D. Incidence of patients with lower extremity injuries presenting to US emergency departments by anatomic region, disease category, and age. *Clin Orthop Relat Res*. 2012;470(1):284-290. doi:10.1007/s11999-011-1982-z.
245. O'Connor SR, Bleakley CM, Tully MA, McDonough SM. Predicting functional recovery after acute ankle sprain. *PLoS One*. 2013;8(8):e72124. doi:10.1371/journal.pone.0072124.
246. Stiell IG, Greenberg GH, McKnight RD, Nair RC, McDowell I, Worthington JR. A study to develop clinical decision rules for the use of radiography in acute ankle injuries. *Ann Emerg Med*. 1992;21(4):384-390.
247. Alparslan L, Chioldo CP. Lateral ankle instability: MR imaging of associated injuries and surgical treatment procedures. *Semin Musculoskelet Radiol*. 2008;12(4):346-358. doi:10.1055/s-0028-1100641.
248. Judd DB, Kim DH. Foot fractures frequently misdiagnosed as ankle sprains. *Am Fam Physician*. 2002;66(5):785-794.
249. Pires R, Pereira A, Abreu-E-Silva G, et al. Ottawa ankle rules and subjective surgeon perception to evaluate radiograph necessity following foot and ankle sprain. *Ann Med Health Sci Res*. 2014;4(3):432-435. doi:10.4103/2141-9248.133473.
250. Eggl S, Scelabas GM, Eggl S, Zimmermann H, Exadaktylos AK. The Bernese ankle rules: a fast, reliable test after low-energy, supination-type malleolar and midfoot trauma. *J Trauma*. 2005;59(5):1268-1271.
251. Derkken R, Knijnenberg LM, Fransen G, Breederveld RS, Heymans MW, Schipper IB. Diagnostic performance of the Bernese versus Ottawa ankle rules: results of a randomised controlled trial. *Injury*. 2015;46(8):1645-1649. doi:10.1016/j.injury.2015.03.038.
252. Hubbard TJ, Hertel J. Anterior positional fault of the fibula after sub-acute lateral ankle sprains. *Man Ther*. 2008;13(1):63-67.
253. Bonnel F, Toullée E, Mabit C, Tourné Y, Sofcot. Chronic ankle instability: biomechanics and pathomechanics of ligaments injury and associated lesions. *Orthop Traumatol Surg Res*. 2010;96(4):424-432. doi:10.1016/j.otsr.2010.04.003.
254. Lamb SE, Marsh JL, Hutton JL, Nakash R, Cooke MW; Collaborative Ankle Support Trial (CAST Group). Mechanical supports for acute, severe ankle sprain: a pragmatic, multicentre, randomised controlled trial. *Lancet*. 2009;373(9663):575-581. doi:10.1016/S0140-6736(09)60206-3.
255. Cheng Y, Cai Y, Wang Y. Value of ultrasonography for detecting chronic injury of the lateral ligaments of the ankle joint compared with ultrasonography findings. *Br J Radiol*. 2014;87(1033):20130406. doi:10.1259/bjr.20130406.
256. Croy T, Koppenhaver S, Saliba S, Hertel J. Anterior talocrural joint laxity: diagnostic accuracy of the anterior drawer test of the ankle. *J Orthop Sports Phys Ther*. 2013;43(12):911-919. doi:10.2519/jospt.2013.4679.
257. Kanamoto T, Shiozaki Y, Tanaka Y, Yonetani Y, Horibe S. The use of MRI in pre-operative evaluation of anterior talofibular ligament in chronic ankle instability. *Bone Joint Res*. 2014;3(8):241-245. doi:10.1302/2046-3758.38.2000295.
258. Aiken AB, Pelland L, Brison R, Pickett W, Brouwer B. Short-term natural recovery of ankle sprains following discharge from emergency departments. *J Orthop Sports Phys Ther*. 2008;38(9):566-571. doi:10.2519/jospt.2008.2811.
259. Hubbard TJ, Kramer LC, Denegar CR, Hertel J. Correlations among multiple measures of functional and mechanical instability in subjects with chronic ankle instability. *J Athl Train*. 2007;42(3):361-366.
260. Mitchell A, Dyson R, Hale T, Abraham C. Biomechanics of ankle instability. Part 1: Reaction time to simulated ankle sprain. *Med Sci Sports Exerc*. 2008;40(8):1515-1521.
261. Mitchell A, Dyson R, Hale T, Abraham C. Biomechanics of ankle instability. Part 2: Postural sway-reaction time relationship. *Med Sci Sports Exerc*. 2008;40(8):1522-1528. doi:10.1249/MSS.0b013e31817356d6.
262. van Middelkoop M, van Rijn RM, Verhaar JA, Koes BW, Bierma-Zeinstra SM. Re-sprains during the first 3 months after initial ankle sprain are related to incomplete recovery: An observational study. *J Physiother*. 2012;58(3):181-188. doi:10.1016/S1836-9553(12)70109-1.
263. Linens SW, Ross SE, Arnold BL, Gayle R, Pidcoe P. Postural-stability tests that identify individuals with chronic ankle instability. *J Athl Train*. 2014;49(1):15-23. doi:10.4085/1062-6050-48.6.09.
264. Trojian TH, McKeag DB. Single leg balance test to identify risk of ankle sprains. *Br J Sports Med*. 2006;40(7):610-613; discussion 613.
265. Doherty C, Bleakley C, Hertel J, Caulfield B, Ryan J, Delahunt E. Postural control strategies during single limb stance following acute lateral ankle sprain. *Clin Biomech (Bristol, Avon)*. 2014;29(6):643-649. doi:10.1016/j.clinbiomech.2014.04.012.
266. Kaminski TW, Hertel J, Amendola N, et al. National Athletic Trainers' Association position statement: conservative management and prevention of ankle sprains in athletes. *J Athl Train*. 2013;48(4):528-545. doi:10.4085/1062-6050-48.4.02.
267. Beynon BD, Renström PA, Haugh L, Uh BS, Barker H. A prospective, randomized clinical investigation of the treatment of first-time ankle sprains. *Am J Sports Med*. 2006;34(9):1401-1412.
268. Jones MH, Amendola AS. Acute treatment of inversion ankle sprains: immobilization versus functional treatment. *Clin Orthop Relat Res*. 2007;455:169-172.

269. Punt IM, Ziltener JL, Laidet M, Armand S, Allet L. Gait and physical impairments in patients with acute ankle sprains who did not receive physical therapy. *PM R*. 2015;7(1):34-41. doi:10.1016/j.pmrj.2014.06.014.
270. Bleakley CM, O'Connor SR, Tully MA, et al. Effect of accelerated rehabilitation on function after ankle sprain: randomised controlled trial. *BMJ*. 2010;340:c1964. doi:10.1136/bmj.c1964.
271. Bassett SF, Prapavessis H. Home-based physical therapy intervention with adherence-enhancing strategies versus clinic-based management for patients with ankle sprains. *Phys Ther*. 2007;87(9):1132-1143.
272. van Rijn RM, van Heest JA, van der Wees P, Koes BW, Bierma-Zeinstra SM. Some benefit from physiotherapy intervention in the subgroup of patients with severe ankle sprain as determined by the ankle function score: a randomised trial. *Aust J Physiother*. 2009;55(2):107-113.
273. Woitzik E, Jacobs C, Wong JJ, et al. The effectiveness of exercise on recovery and clinical outcomes of soft tissue injuries of the leg, ankle, and foot: A systematic review by the Ontario Protocol for Traffic Injury Management (OPTIMa) Collaboration. *Man Ther*. 2015;20(5):633-645. doi:10.1016/j.math.2015.03.012.
274. Hupperets MD, Verhagen EA, van Mechelen W. Effect of unsupervised home based proprioceptive training on recurrences of ankle sprain: randomised controlled trial. *BMJ*. 2009;339:b2684. doi:10.1136/bmj.b2684.
275. Mailuhu AK, Verhagen EA, van Ochten JM, Bindels PJ, Bierma-Zeinstra SM, van Middelkoop M. The trAPP-study: cost-effectiveness of an unsupervised e-health supported neuromuscular training program for the treatment of acute ankle sprains in general practice: design of a randomized controlled trial. *BMC Musculoskelet Disord*. 2015;16:78. doi:10.1186/s12891-015-0539-9.
276. van Ochten JM, van Middelkoop M, Meuffels D, Bierma-Zeinstra SM. Chronic complaints after ankle sprains: a systematic review on effectiveness of treatments. *J Orthop Sports Phys Ther*. 2014;44(11):862-871, C1-C3. doi:10.2519/jospt.2014.5221.
277. Green T, Refshauge K, Crosbie J, Adams R. A randomized controlled trial of a passive accessory joint mobilization on acute ankle inversion sprains. *Phys Ther*. 2001;81(4):984-994.
278. Whitman JM, Cleland JA, Mintken PE, et al. Predicting short-term response to thrust and nonthrust manipulation and exercise in patients post inversion ankle sprain. *J Orthop Sports Phys Ther*. 2009;39(3):188-200. doi:10.2519/jospt.2009.2940.
279. Cleland JA, Mintken P, McDevitt A, et al. Manual physical therapy and exercise versus supervised home exercise in the management of patients with inversion ankle sprain: a multicenter randomized clinical trial. *J Orthop Sports Phys Ther*. 2013;43(7):443-455. doi:10.2519/jospt.2013.4792.
280. Truyols-Domi Nguez S, Salom-Moreno J, Abian-Vicen J, Cleland JA, Fernández-de-Las-Peñas C. Efficacy of thrust and nonthrust manipulation and exercise with or without the addition of myofascial therapy for the management of acute inversion ankle sprain: a randomized clinical trial. *J Orthop Sports Phys Ther*. 2013;43(5):300-309. doi:10.2519/jospt.2013.4467.
281. Terada M, Pietrosimone BG, Gribble PA. Therapeutic interventions for increasing ankle dorsiflexion after ankle sprain: A systematic review. *J Athl Train*. 2013;48(5):696-709. doi:10.4085/1062-6050-48.4.11.
282. Hubbard TJ, Hertel J. Mechanical contributions to chronic lateral ankle instability. *Sports Med*. 2006;36(3):263-277.
283. Hubbard TJ, Hertel J, Sherbondy P. Fibular position in individuals with self-reported chronic ankle instability. *J Orthop Sports Phys Ther*. 2006;36(1):3-9.
284. Vicenzino B, Branjerdpor M, Teys P, Jordan K. Initial changes in posterior talar glide and dorsiflexion of the ankle after mobilization with movement in individuals with recurrent ankle sprain. *J Orthop Sports Phys Ther*. 2006;36(7):464-471.
285. Delahunt E, Cusack K, Wilson L, Doherty C. Joint mobilization acutely improves landing kinematics in chronic ankle instability. *Med Sci Sports Exerc*. 2013;45(3):514-519. doi:10.1249/MSS.0b013e3182746d0a.
286. Whitman JM, Childs JD, Walker V. The use of manipulation in a patient with an ankle sprain injury not responding to conventional management: a case report. *Man Ther*. 2005;10(3):224-231.
287. Janssen KW, van Mechelen W, Verhagen EA. Bracing superior to neuromuscular training for the prevention of self-reported recurrent ankle sprains: a three-arm randomised controlled trial. *Br J Sports Med*. 2014;48(10):1235-1239. doi:10.1136/bjsports-2013-092947.
288. Janssen KW, van der Zwaard BC, Finch CF, van Mechelen W, Verhagen EA. Interventions preventing ankle sprains; previous injury and high-risk sport participation as predictors of compliance. *J Sci Med Sport*. 2015;19(6):465-469. doi:10.1016/j.jsm.2015.06.005.
289. Lardenoye S, Theunissen E, Cleffken B, Brink PR, de Bie RA, Poeze M. The effect of taping versus semi-rigid bracing on patient outcome and satisfaction in ankle sprains: a prospective, randomized controlled trial. *BMC Musculoskelet Disord*. 2012;13(1):81. doi:10.1186/1471-2474-13-81.
290. Handoll HH, Rowe BH, Quinn KM, de Bie R. Interventions for preventing ankle ligament injuries. *Cochrane Database Syst Rev*. 2001;(3):CD000018.
291. Guillodo Y, Simon T, Le Goff A, Saraux A. Interest of rehabilitation in healing and preventing recurrence of ankle sprains. *Ann Phys Rehabil Med*. 2013;56(7-8):503-514. doi:10.1016/j.rehab.2013.06.007.
292. Guillodo Y, Le Goff A, Saraux A. Adherence and effectiveness of rehabilitation in acute ankle sprain. *Ann Phys Rehabil Med*. 2011;54(4):225-235. doi:10.1016/j.rehab.2011.03.002.
293. Gribble PA, Delahunt E, Bleakley C, et al. Selection criteria for patients with chronic ankle instability in controlled research: a position statement of the International Ankle Consortium. *J Orthop Sports Phys Ther*. 2013;43(8):585-591. doi:10.2519/jospt.2013.0303.
294. DiGiovanni BF, Fraga CJ, Cohen BE, Shereff MJ. Associated injuries found in chronic lateral ankle instability. *Foot Ankle Int*. 2000;21(10):809-815.
295. Ferkel RD, Zanotti RM, Komenda GA, et al. Arthroscopic treatment of chronic osteochondral lesions of the talus: long-term results. *Am J Sports Med*. 2008;36(9):1750-1762. doi:10.1177/0363546508316773.
296. Loudon JK, Santos MJ, Franks L, Liu W. The effectiveness of active exercise as an intervention for functional ankle instability: a systematic review. *Sports Med*. 2008;38(7):553-563.
297. Hirai D, Docherty CL, Schrader J. Severity of functional and mechanical ankle instability in an active population. *Foot Ankle Int*. 2009;30(11):1071-1077. doi:10.3131/FAI.2009.1071.
298. Cha SD, Kim HS, Chung ST, et al. Intra-articular lesions in chronic lateral ankle instability: comparison of arthroscopy with magnetic resonance imaging findings. *Clin Orthop Surg*. 2012;4(4):293-299. doi:10.4055/cios.2012.4.4.293.
299. Houston MN, Van Lunen BL, Hoch MC. Health-related quality of life in individuals with chronic ankle instability. *J Athl Train*. 2014;49(6):758-763. doi:10.4085/1062-6050-49.3.54.
300. Hubbard-Turner T, Turner MJ. Physical activity levels in college students with chronic ankle instability. *J Athl Train*. 2015;50(7):742-747. doi:10.4085/1062-6050-50.3.05.
301. Hiller CE, Kilbreath SL, Refshauge KM. Chronic ankle instability: evolution of the model. *J Athl Train*. 2011;46(2):133-141. doi:10.4085/1062-6050-46.2.133.
302. Wright CJ, Arnold BL, Ross SE, Linens SW. Recalibration and validation of the Cumberland Ankle Instability Tool cutoff score for individuals with chronic ankle instability. *Arch Phys Med Rehabil*. 2014;95(10):1853-1859. doi:10.1016/j.apmr.2014.04.017.
303. Kobayashi T, Suzuki E, Yamazaki N, et al. Fibular malalignment in individuals with chronic ankle instability. *J Orthop Sports Phys Ther*. 2014;44(11):872-878. doi:10.2519/jospt.2014.5217.
304. Croy T, Saliba SA, Saliba E, Anderson MW, Hertel J. Differences in lateral ankle laxity measured via stress ultrasonography in individuals with chronic ankle instability, ankle sprain copers, and healthy individuals. *J Orthop Sports Phys Ther*. 2012;42(7):593-600. doi:10.2519/jospt.2012.3923.
305. Bastien M, Moffet H, Bouyer LJ, Perron M, Hébert LJ, Leblond J. Alteration in global motor strategy following lateral ankle sprain. *BMC Musculoskelet Disord*. 2014;15(1):436. doi:10.1186/1471-2474-15-436.
306. Wikstrom EA, Hubbard-Turner T, McKeon PO. Understanding and treating lateral ankle sprains and their consequences: a constraints-based approach. *Sports Med*. 2013;43(6):385-393. doi:10.1007/s40279-013-0043-z.
307. McLeod MM, Gribble PA, Pietrosimone BG. Chronic ankle instability and neural excitability of the lower extremity. *J Athl Train*. 2015;50(8):847-853. doi:10.4085/1062-6050-50.4.06.
308. Sefton JM, Yarar CA, Hicks-Little CA, Berry JW, Cordova ML. Six weeks of balance training improves sensorimotor function in individuals with chronic ankle instability. *J Orthop Sports Phys Ther*. 2011;41(2):81-89. doi:10.2519/jospt.2011.3365.
309. Hale SA, Fergus A, Axmacher R, Kiser K. Bilateral improvements in lower extremity function after unilateral balance training in individuals with chronic ankle instability. *J Athl Train*. 2014;49(2):181-191. doi:10.4085/1062-6050-49.2.06.
310. Hilgendorf JR, Vela LI, Goertz DV, Harter RA. Influence of vestibular-ocular reflex training on postural stability, dynamic visual acuity, and gaze stabilization in patients with chronic ankle instability. *J Athl Train Sports Health Care*. 2012;4(5):220-229.
311. Ross SE, Linens SW, Wright CJ, Arnold BL. Noise-enhanced eversion force sense in ankles with or without functional instability. *J Athl Train*. 2015;50(8):819-824. doi:10.4085/1062-6050-50.5.06.
312. Sefton JM, Hicks-Little CA, Hubbard TJ, et al. Sensorimotor function as a predictor of chronic ankle instability. *Clin Biomech (Bristol, Avon)*. 2009;24(5):451-458. doi:10.1016/j.clinbiomech.2009.03.003.
313. Hoch MC, Farwell KE, Gaven SL, Weinhandl JT. Weight-bearing dorsiflexion range of motion and landing biomechanics in individuals with chronic ankle instability. *J Athl Train*. 2015;50(8):833-839. doi:10.4085/1062-6050-50.5.07.
314. Gigi R, Haim A, Luger E, et al. Deviations in gait metrics in patients with chronic ankle instability: a case control study. *J Foot Ankle Res*. 2015;8(1):1. doi:10.1186/s13047-014-0058-1.
315. Cruz-Díaz D, Lomas Vega R, Osuna-Pérez MC, Hita-Contreras F, Martínez-Amat A. Effects of joint mobilization on chronic ankle instability: a randomized controlled trial. *Disabil Rehabil*. 2015;37(7):601-610. doi:10.3109/09638288.2014.935877.
316. McKeon PO, Poolin G, Ingersoll CD, et al. Effects of balance training on gait parameters in patients with chronic ankle instability: a randomized controlled trial. *Clin Rehabil*. 2009;23(7):609-621. doi:10.1177/0269215509102954.
317. Huang PY, Chen WL, Lin CF, Lee HJ. Lower extremity biomechanics in athletes with ankle instability after 6-week integrated training program. *J Athl Train*. 2014;49(2):163-172. doi:10.4085/1062-6050-49.2.10.
318. Donovan L, Hart JM, Hertel J. Effects of 2 ankle destabilization devices on electromyography measures during functional exercises in individuals with chronic ankle instability. *J Orthop Sports Phys Ther*. 2015;45(3):220-232. doi:10.2519/jospt.2015.5222.
319. Ross SE, Linens SW, Wright CJ, Arnold BL. Customized noise-stimulation intensity for bipedal stability and unipedal balance deficits associated with functional ankle instability. *J Athl Train*. 2013;48(4):463-470. doi:10.4085/1062-6050-48.3.12.
320. Lee AJ, Lin WH. Twelve-week biomechanical ankle platform system training on postural stability and ankle proprioception in subjects with unilateral functional ankle instability. *Clinical Biomech (Bristol, Avon)*. 2008;23(8):1065-1072. doi:10.1016/j.clinbiomech.2008.04.013.
321. Hoch MC, Mullineaux DR, Andreatta RD, et al. Effect of a 2-week joint mobilization intervention on single-limb balance and ankle arthrokinematics in those with chronic ankle instability. *J Sport Rehabil*. 2014;23(1):18-26. doi:10.1123/jsr.2012-0125.

322. Schaefer JL, Sandrey MA. Effects of a 4-week dynamic-balance-training program supplemented with Graston-instrument assisted soft-tissue mobilization for chronic ankle instability. *J Sport Rehabil.* 2012;21(4):313-326.
323. Ross SE, Linens SW, Wright CJ, Arnold BL. Balance assessments for predicting functional ankle instability and stable ankles. *Gait Posture.* 2011;34(4):539-542. doi:10.1016/j.gaitpost.2011.07.011.
324. Tourné Y, Besse JL, Mabit C; Sofcot. Chronic ankle instability. Which tests to assess the lesions? Which therapeutic options? *Orthop Traumatol Surg Res.* 2010;96(4):433-446. doi:10.1016/j.otsr.2010.04.005.
325. Stehr MT. Lateral ankle ligament injury following inversion ankle sprain. *J Orthop Sports Phys Ther.* 2012;42(2):145. doi:10.2519/jospt.2012.0403.
326. Guillou S, Bauer T, Lee JW, et al. Consensus in chronic ankle instability: aetiology, assessment, surgical indications and place for arthroscopy. *Orthop Traumatol Surg Res.* 2013;99(8 Suppl):S411-S419. doi:10.1016/j.otsr.2013.10.009.
327. Baumhauer JF, O'Brien T. Surgical considerations in the treatment of ankle instability. *J Athl Train.* 2002;37(4):458-462.
328. van den Bekker MP, Raven EE. The distal fascicle of the anterior inferior tibiofibular ligament as a cause of tibiotalar impingement syndrome: A current concepts review. *Knee Surg Sports Traumatol Arthrosc.* 2007;15(4):465-471.
329. Tol JL, Verhagen RA, Krips R, et al. The anterior ankle impingement syndrome: diagnostic value of oblique radiographs. *Foot Ankle Int.* 2004;25(2):63-68.
330. Walsh SJ, Twaddle BC, Rosenfeldt MP, Boyle MJ. Arthroscopic treatment of anterior ankle impingement: a prospective study of 46 patients with 5-year follow-up. *Am J Sports Med.* 2014;42(11):2722-2726. doi:10.1177/0363546514550976.
331. Lee JW, Suh JS, Huh YM, Moon ES, Kim SJ. Soft tissue impingement syndrome of the ankle: diagnostic efficacy of MRI and clinical results after arthroscopic treatment. *Foot Ankle Int.* 2004;25(12):896-902.
332. Takao M, Uchio Y, Naito K, Fukazawa I, Ochi M. Arthroscopic assessment for intra-articular disorders in residual ankle disability after sprain. *Am J Sports Med.* 2005;33(5):686-692.
333. Bauer T, Breda R, Hardy P. Anterior ankle bony impingement with joint motion loss: The arthroscopic resection option. *Orthop Traumatol Surg Res.* 2010;96(4):462-468. doi:10.1016/j.otsr.2010.01.008.
334. O'Neill PJ, Van Aman SE, Guyton GP. Is MRI adequate to detect lesions in patients with ankle instability? *Clin Orthop Relat Res.* 2010;468(4):1115-1119. doi:10.1007/s11999-009-1131-0.
335. Krause FG, Di Silvestro M, Penner MJ, et al. Inter- and intraobserver reliability of the COFAS end-stage ankle arthritis classification system. *Foot Ankle Int.* 2010;31(2):103-108. doi:10.3113/FAI.2010.0103.
336. Lapidus PW, Guidotti FP. Painful heel: report of 323 patients with 364 painful heels. *Clin Orthop Relat Res.* 1965;39:178-186.
337. Osborne HR, Breidahl WH, Allison GT. Critical differences in lateral X-rays with and without a diagnosis of plantar fasciitis. *J Sci Med Sport.* 2006;9(3):231-237.
338. Lemont H, Ammirati KM, Usen N. Plantar fasciitis: a degenerative process (fasciosis) without inflammation. *J Am Podiatr Med Assoc.* 2003;93(3):234-237.
339. Riddle DL, Schappert SM. Volume of ambulatory care visits and patterns of care for patients diagnosed with plantar fasciitis: a national study of medical doctors. *Foot Ankle Int.* 2004;25(5):303-310.
340. Riddle DL, Pulsic M, Pidcoe P, Johnson RE. Risk factors for Plantar fasciitis: a matched case-control study. *J Bone Joint Surg Am.* 2003;85-A(5):872-877.
341. Martin RL, Davenport TE, Reischl SF, et al. Heel pain-plantar fasciitis: revision 2014. *J Orthop Sports Phys Ther.* 2014;44(11):A1-33. doi:10.2519/jospt.2014.0403.
342. Thomas G, Whalley H, Modi C. Early mobilization of operatively fixed ankle fractures: a systematic review. *Foot Ankle Int.* 2009;30(7):666-674. doi:10.3113/FAI.2009.0666.
343. League AC. Current concepts review: plantar fasciitis. *Foot Ankle Int.* 2008;29(3):358-366. doi:10.3113/FAI.2008.0358.
344. Neufeld SK, Cerrato R. Plantar fasciitis: evaluation and treatment. *J Am Acad Orthop Surg.* 2008;16(6):338-346.
345. Beischer AD, Clarke A, de Steiger RN, Donnan L, Ibuki A, Unglik R. The practical application of multimedia technology to facilitate the education and treatment of patients with plantar fasciitis: a pilot study. *Foot Ankle Spec.* 2008;1(1):30-38. doi:10.1177/1938640007312299..
346. Tweed JL, Barnes MR, Allen MJ, Campbell JA. Biomechanical consequences of total plantar fasciotomy: a review of the literature. *J Am Podiatr Med Assoc.* 2009;99(5):422-430.
347. van de Water AT, Speksnijder CM. Efficacy of taping for the treatment of plantar fasciosis: a systematic review of controlled trials. *J Am Podiatr Med Assoc.* 2010;100(1):41-51.
348. Hyland MR, Webber-Gaffney A, Cohen L, Lichtman PT. Randomized controlled trial of calcaneal taping, sham taping, and plantar fascia stretching for the short-term management of plantar heel pain. *J Orthop Sports Phys Ther.* 2006;36(6):364-371.
349. Hawke F, Burns J, Radford JA, du Toit V. Custom-made foot orthoses for the treatment of foot pain. *Cochrane Database Syst Rev.* 2008;(3):CD006801. doi:10.1002/14651858.CD006801.pub2.
350. Landorf KB, Keenan AM, Herbert RD. Effectiveness of foot orthoses to treat plantar fasciitis: a randomized trial. *Arch Intern Med.* 2006;166(12):1305-1310.
351. DiGiovanni BF, Nawoczenski DA, Lintal ME, et al. Tissue-specific plantar fascia-stretching exercise enhances outcomes in patients with chronic heel pain. A prospective, randomized study. *J Bone Joint Surg Am.* 2003;85-A(7):1270-1277.
352. Wearing SC, Smeathers JE, Yates B, Urry SR, Dubois P. Bulk compressive properties of the heel fat pad during walking: a pilot investigation in plantar heel pain. *Clin Biomech (Bristol, Avon).* 2009;24(4):397-402. doi:10.1016/j.clinbiomech.2009.01.002.
353. Landorf KB, Menz HB. Plantar heel pain and fasciitis. *BMJ Clin Evid.* 2008;pii.1111.
354. Sweeting D, Parish B, Hooper L, Chester R. The effectiveness of manual stretching in the treatment of plantar heel pain: a systematic review. *J Foot Ankle Res.* 2011;4:19. doi:10.1186/1757-1146-4-19.
355. Martin RL, Irrgang JJ, Conti SF. Outcome study of subjects with insertional plantar fasciitis. *Foot Ankle Int.* 1998;19(12):803-811.
356. Lee WC, Wong WY, Kung E, Leung AK. Effectiveness of adjustable dorsiflexion night splint in combination with accommodative foot orthosis on plantar fasciitis. *J Rehabil Res Dev.* 2012;49(10):1557-1564.
357. Sheridan L, Lopez A, Perez A, John MM, Willis FB, Shamugam R. Plantar fasciopathy treated with dynamic splinting: a randomized controlled trial. *J Am Podiatr Med Assoc.* 2010;100(3):161-165.
358. Renan-Ordine R, Albuquerque-Sendin F, de Souza DP, Cleland JA, Fernandez-de-Las-Penas C. Effectiveness of myofascial trigger point manual therapy combined with a self-stretching protocol for the management of plantar heel pain: a randomized controlled trial. *J Orthop Sports Phys Ther.* 2011;41(2):43-50. doi:10.2519/jospt.2011.3504.
359. Stratton M, McPoil TG, Cornwell MW, Patrick K. Use of low-frequency electrical stimulation for the treatment of plantar fasciitis. *J Am Podiatr Med Assoc.* 2009;99(6):481-488.
360. DiGiovanni BF, Nawoczenski DA, Malay DP, et al. Plantar fascia-specific stretching exercise improves outcomes in patients with chronic plantar fasciitis. A prospective clinical trial with two-year follow-up. *J Bone Joint Surg Am.* 2006;88(8):1775-1781.
361. Aronow MS, Diaz-Doran V, Sullivan RJ, Adams DJ. The effect of triceps surae contracture force on plantar foot pressure distribution. *Foot Ankle Int.* 2006;27(1):43-52.
362. Maskill JD, Bohay DR, Anderson JG. Gastrocnemius recession to treat isolated foot pain. *Foot Ankle Int.* 2010;31(1):19-23. doi:10.3113/FAI.2010.0019.
363. Monteagudo M, Maceira E, Garcia-Virto V, Canosa R. Chronic plantar fasciitis: plantar fasciotomy versus gastrocnemius recession. *Int Orthop.* 2013;37(9):1845-1850.
364. Chimera NJ, Castro M, Manal K. Function and strength following gastrocnemius recession for isolated gastrocnemius contracture. *Foot Ankle Int.* 2010;31(5):377-384. doi:10.3113/FAI.2010.0377.
365. Liden B, Simmons M, Landsman AS, Liden B, Simmons M, Landsman AS. A retrospective analysis of 22 patients treated with percutaneous radiofrequency nerve ablation for prolonged moderate to severe heel pain associated with plantar fasciitis. *J Foot Ankle Surg.* 2009;48(6):642-647. doi:10.1053/j.jfas.2009.05.013.
366. Cheing GL, Chang H. Extracorporeal shock wave therapy. *J Orthop Sports Phys Ther.* 2003;33(6):337-343.
367. Mosier KM, Asher M. Tarsal coalitions and peroneal spastic flat foot. A review. *J Bone Joint Surgery Am.* 1984;66(7):976-984.
368. Jayakumar S, Cowell HR. Rigid flatfoot. *Clin Orthop Relat Res.* 1977;122:77-84.
369. Nalaboff KM, Schweitzer ME. MRI of tarsal coalition: frequency, distribution, and innovative signs. *Bull NYU Hosp Joint Dis.* 2008;66(1):14-21.
370. Stormont DM, Peterson HA. The relative incidence of tarsal coalition. *Clin Orthop Relat Res.* 1983;181:28-36.
371. Katayama T, Tanaka Y, Kadono K, Taniguchi A, Takakura Y. Talocalcaneal coalition: a case showing the ossification process. *Foot Ankle Int.* 2005;26(6):490-493.
372. Kernbach KJ. Tarsal coalitions: etiology, diagnosis, imaging, and stigmata. *Clin Podiatr Med Surg.* 2010;27(1):105-117. doi:10.1016/j.cpm.2009.08.006.
373. Zaw H, Calder JD. Tarsal coalitions. *Foot Ankle Clin.* 2010;15(2):349-364. doi:10.1016/j.facl.2010.02.003.
374. Hetsonri I, Nyska M, Mann G, Rozenfeld G, Ayalon M. Subtalar kinematics following resection of tarsal coalition. *Foot Ankle Int.* 2008;29(11):1088-1094. doi:10.3113/FAI.2008.1088.
375. Lemley F, Berlet G, Hill K, Philbin T, Isaac B, Lee T. Current concepts review: Tarsal coalition. *Foot Ankle Int.* 2006;27(12):1163-1169.
376. Lahey MD, Zindrick MR, Harris EJ. A comparative study of the clinical presentation of tarsal coalitions. *Clin Podiatr Med Surg.* 1988;5(2):341-357.
377. Leonard MA. The inheritance of tarsal coalition and its relationship to spastic flat foot. *J Bone Joint Surg Br.* 1974;56B(3):520-526.
378. Lovell WW, Winter RB, Morrissey RT, Weinstein SL. *Lovell and Winter's Pediatric Orthopaedics.* Vol 6th. Philadelphia, PA: Lippincott Williams & Wilkins; 2006.
379. Gonzalez P, Kumar SJ. Calcaneonavicular coalition treated by resection and interposition of the extensor digitorum brevis muscle. *J Bone Joint Surg Am.* 1990;72(1):71-77.
380. Wilde PH, Torode IP, Dickens DR, Cole WG. Resection for symptomatic talocalcaneal coalition. *J Bone Joint Surg Br.* 1994;76(5):797-801.
381. Luhmann SJ, Schoenecker PL. Symptomatic talocalcaneal coalition resection: indications and results. *J Pediatr Orthop.* 1998;18(6):748-754.
382. Swionkowski MF, Scranton PE, Hansen S. Tarsal coalitions: long-term results of surgical treatment. *J Pediatr Orthop.* 1983;3(3):287-292.
383. Takakura Y, Sugimoto K, Tanaka Y, Tamai S. Symptomatic talocalcaneal coalition. Its clinical significance and treatment. *Clin Orthop Relat Res.* 1991;269:249-256.
384. Hetsonri I, Ayalon M, Mann G, Meyer G, Nyska M. Walking and running plantar pressure analysis before and after resection of tarsal coalition. *Foot Ankle Int.* 2007;28(5):575-580.
385. Nilesh KM. (iii) Tarsometatarsal injuries—Lisfranc injuries. *2005;19(2):108-118.*
386. Khan W, Oragui E, Akagha E. Common fractures and injuries of the ankle and foot: functional anatomy, imaging, classification and management. *J Peripher Pract.* 2010;20(7):249-258.

387. Kaar S, Femino J, Morag Y. Lisfranc joint displacement following sequential ligament sectioning. *J Bone Joint Surg Am.* 2007;89(10):2225-2232.
388. Khan RJ, Fick D, Keogh A, Crawford J, Brammar T, Parker M. Treatment of acute achilles tendon ruptures: A meta-analysis of randomized, controlled trials. *J Bone Joint Surg Am.* 2005;87(10):2202-2210.
389. Reid JJ, Pinney SJ. Midfoot injuries in athletes: fundamentals of examination and treatment. *Oper Tech Sports Med.* 2010;18(1):46-49. doi:10.1053/j.otsm.2009.11.006
390. Nunley JA, Vertullo CJ. Classification, investigation, and management of midfoot sprains: Lisfranc injuries in the athlete. *Am J Sports Med.* 2002;30(6):871-878.
391. Sherief TI, Mucci B, Greiss M. Lisfranc injury: how frequently does it get missed? And how can we improve? *Injury.* 2007;38(7):856-860.
392. Scharfbillig RW, Jones S, Scutter SD. Sever's disease: what does the literature really tell us? *J Am Podiatr Med Assoc.* 2008;98(3):212-223.
393. Cassas KJ, Cassetta-Wayhs A. Childhood and adolescent sports-related overuse injuries. *Am Fam Physician.* 2006;73(6):1014-1022.
394. James AM, Williams CM, Luscombe M, Hunter R, Haines TP. Factors associated with pain severity in children with calcaneal apophysitis (Sever disease). *J Pediatr.* 2015;167(2):455-459. doi:10.1016/j.jpeds.2015.04.053.
395. Wiegerinck JI, Yntema C, Brouwer HJ, Struijs PA. Incidence of calcaneal apophysitis in the general population. *Eur J Pediatr.* 2014;173(5):677-679. doi:10.1007/s00431-013-2219-9.
396. Manusov EG, Lillegard WA, Raspa RF, Epperly TD. Evaluation of pediatric foot problems: Part II. The hindfoot and the ankle. *Am Fam Physician.* 1996;54(3):1012-1026, 1031.
397. Madden CC, Mellon MB. Sever's disease and other causes of heel pain in adolescents. *Am Fam Physician.* 1996;54(6):1995-2000.
398. Szames SE, Forman WM, Oster J, Eleff JC, Woodward P. Sever's disease and its relationship to equinus: a statistical analysis. *Clin Podiatr Med Surg.* 1990;7(2):377-384.
399. Scharfbillig RW, Jones S, Scutter S. Sever's disease--does it effect quality of life? *Foot (Edinb).* 2009;19(1):36-43. doi:10.1016/j.foot.2008.07.004.
400. Perhamre S, Lazowska D, Papageorgiou S, Lundin F, Klassbo M, Norlin R. Sever's injury: a clinical diagnosis. *J Am Podiatr Med Assoc.* 2013;103(5):361-368.
401. Kose O, Celiktaç M, Yigit S, Kisim B. Can we make a diagnosis with radiographic examination alone in calcaneal apophysitis (Sever's disease)? *J Pediatr Orthop B.* 2010;19(5):396-398. doi:10.1097/BPB.0b013e32833a5fa7.
402. Volpon JB, de Carvalho Filho G. Calcaneal apophysitis: a quantitative radiographic evaluation of the secondary ossification center. *Arch Orthop Trauma Surg.* 2002;122(6):338-341.
403. James AM, Williams CM, Haines TP. "Effectiveness of interventions in reducing pain and maintaining physical activity in children and adolescents with calcaneal apophysitis (Sever's disease): a systematic review." *J Foot Ankle Res.* 2013;6(1):16. doi:10.1186/1757-1146-6-16.
404. Khoury J, Jerushalmi J, Loberant N, Shtarker H, Militianu D, Keidar Z. Kohler disease: diagnoses and assessment by bone scintigraphy. *Clin Nucl Med.* 2007;32(3):179-181.
405. Ippolito E, Ricciardi Pollini PT, Falez F. Kohler's disease of the tarsal navicular: long-term follow-up of 12 cases. *J Pediatr Orthop.* 1984;4(4):416-417.
406. Tsirikos AI, Riddle EC, Kruse R. Bilateral Kohler's disease in identical twins. *Clin Orthop Relat Res.* 2003;409:195-198.
407. Borges JL, Guille JT, Bowen JR. Kohler's bone disease of the tarsal navicular. *J Pediatr Orthop.* 1995;15(5):596-598.
408. Williams GA, Cowell HR. Kohler's disease of the tarsal navicular. *Clin Orthop Relat Res.* 1981;118(158):53-58.
409. Coughlin MJ, Shurnas PS. Hallux rigidus: Grading and long-term results of operative treatment. *J Bone Joint Surg Am.* 2003;85-A(11):2072-2088.
410. Meyer JO, Nishon LR, Weiss L, Docks G. Metatarsus primus elevatus and the etiology of hallux rigidus. *J Foot Surg.* 1987;26(3):237-241.
411. Goodfellow J. Aetiology of hallux rigidus. *Proc R Soc Med.* 1966;59(9):821-824.
412. McMaster MJ. The pathogenesis of hallux rigidus. *J Bone Joint Surg Br.* 1978;60(1):82-87.
413. Shurnas PS. Hallux rigidus: etiology, biomechanics, and nonoperative treatment. *Foot Ankle Clin.* 2009;14(1):1-8. doi:10.1016/j.facl.2008.11.001.
414. Solan MC, Calder JD, Bendall SP. Manipulation and injection for hallux rigidus. Is it worthwhile? *J Bone Joint Surg Br.* 2001;83(5):706-708.
415. Thompson JA, Jennings MB, Hodge W. Orthotic therapy in the management of osteoarthritis. *J Am Podiatr Med Assoc.* 1992;82(3):136-139.
416. Smith RW, Katchis SD, Ayson LC. Outcomes in hallux rigidus patients treated nonoperatively: a long-term follow-up study. *Foot Ankle Int.* 2000;21(11):906-913.
417. Fanous RN, Ridgers S, Sott AH. Minimally invasive arthrodesis of the first metatarsophalangeal joint for hallux rigidus. *Foot Ankle Surg.* 2014;20(3):170-173. doi:10.1016/j.fas.2014.03.004.
418. Herrera-Perez M, Andarcia-Banuelos C, de Bergua-Domingo J, Paul J, Barg A, Valderabano V. Proposed global treatment algorithm for hallux rigidus according to evidence-based medicine. *Rev Esp Cir Ortop Traumatol.* 2014;58(6):377-386. doi:10.1016/j.recot.2014.05.004.
419. Hunt KJ. Hallux metatarsophalangeal (MTP) joint arthroscopy for hallux rigidus. *Foot Ankle Int.* 2015;36(1):113-119. doi:10.1177/1071100714559728.
420. Meric G, Erduran M, Atik A, Kose O, Ulusal AE, Akseki D. Short-term clinical outcomes after first metatarsal head resurfacing hemiarthroplasty for late stage hallux rigidus. *J Foot Ankle Surg.* 2015;54(2):173-178. doi:10.1053/j.jfas.2014.10.016.
421. Hattrup SJ, Johnson KA. Subjective results of hallux rigidus following treatment with cheilectomy. *Clin Orthop Relat Res.* 1988;(226):182-191.
422. Coughlin MJ, Shurnas PS. Hallux rigidus. *J Bone Joint Surg Am.* 2004;86-A Suppl 1(Pt 2):119-130.
423. Mann RA, Clanton TO. Hallux rigidus: treatment by cheilectomy. *J Bone Joint Surg Am.* 1988;70(3):400-406.
424. Moberg E. A simple operation for hallux rigidus. *Clin Orthop Relat Res.* 1979;142(7):55-56.
425. Curtis MJ, Myerson M, Jinnah RH, Cox QG, Alexander I. Arthrodesis of the first metatarsophalangeal joint: a biomechanical study of internal fixation techniques. *Foot Ankle.* 1993;14(7):395-399.
426. Chalmers AC, Busby C, Goyert J, Porter B, Schulzer M. Metatarsalgia and rheumatoid arthritis—a randomized, single blind, sequential trial comparing 2 types of foot orthoses and supportive shoes. *J Rheumatol.* 2000;27(7):1643-1647.
427. Cychoz CC, Phisitkul P, Belatti DA, Glazebrook MA, DiGiovanni CW. Gastrocnemius recession for foot and ankle conditions in adults: Evidence-based recommendations. *Foot Ankle Surg.* 2015;21(2):77-85. doi:10.1016/j.fas.2015.02.001.
428. Aper RL, Saltzman CL, Brown TD. The effect of hallux sesamoid excision on the flexor hallucis longus moment arm. *Clin Orthop Relat Res.* 1996;(325):209-217.
429. Lee S, James WC, Cohen BE, Davis WH, Anderson RB. Evaluation of hallux alignment and functional outcome after isolated tibial sesamoidectomy. *Foot Ankle Int.* 2005;26(10):803-809.
430. Bourke G, Owen J, Machet D. Histological comparison of the third interdigital nerve in patients with Morton's metatarsalgia and control patients. *Australian N Z J Surg.* 1994;64(6):421-424.
431. Bignotti B, Signori A, Sormani MP, Moffetta L, Martinoli C, Tagliafico A. Ultrasound versus magnetic resonance imaging for Morton neuroma: systematic review and meta-analysis. *Eur Radiol.* 2015;25(8):2254-2264. doi:10.1007/s00330-015-3633-3.
432. Pardal-Fernandez JM, Palazon-Garcia E, Hernandez-Fernandez F, de Cabo C. Contribution of a new electrophysiologic test to Morton's neuroma diagnosis. *Foot Ankle Surg.* 2014;20(2):109-114. doi:10.1016/j.fas.2013.12.006.
433. Greenfield J, Rea J Jr, Ilfeld FW. Morton's interdigital neuroma. Indications for treatment by local injections versus surgery. *Clin Orthop Relat Res.* 1984;185(142-144).
434. Saygi B, Yildirim Y, Saygi EK, Kara H, Esemenli T. Morton neuroma: comparative results of two conservative methods. *Foot Ankle Int.* 2005;26(7):556-559.
435. Bennett GL, Graham CE, Mauldin DM. Morton's interdigital neuroma: a comprehensive treatment protocol. *Foot Ankle Int.* 1995;16(12):760-763.
436. Thomson CE, Beggs I, Martin DJ, et al. Methylprednisolone injections for the treatment of Morton neuroma: a patient-blinded randomized trial. *J Bone Joint Surg Am.* 2013;95(9):790-798. doi:10.2106/JBJS.I.01780.
437. Rasmussen MR, Kitaoka HB, Patzer GL. Nonoperative treatment of plantar interdigital neuroma with a single corticosteroid injection. *Clin Orthop Relat Res.* 1996;326(188-193).
438. Gaynor R, Hake D, Spinner SM, Tomczak RL. A comparative analysis of conservative versus surgical treatment of Morton's neuroma. *J Am Podiatr Med Assoc.* 1989;79(1):27-30.
439. Nunan PJ, Giese BD. Management of Morton's neuroma in athletes. *Clin Podiatr Med Surg.* 1997;14(3):489-501.
440. Hirschberg GG. A simple cure for Morton's neuralgia. *J Am Podiatr Med Assoc.* 2000;90(2):100-101.
441. Kasperek M, Schneider W. Surgical treatment of Morton's neuroma: clinical results after open excision. *Int Orthop.* 2013;37(9):1857-1861. doi:10.1007/s00264-013-2002-6.
442. Okafor B, Shergill G, Angel J. Treatment of Morton's neuroma by neurolysis. *Foot Ankle Int.* 1997;18(5):284-287.
443. Mann RA, Reynolds JC. Interdigital neuroma—a critical clinical analysis. *Foot Ankle.* 1983;3(4):238-243.
444. Rompe JD. Plantar fasciopathy. *Sports Med Arthrosc.* 2009;17(2):100-104. doi:10.1097/JS.0b013e3181a3d60e.
445. Mosier SM, Lucas DR, Pomeroy G, Manoli A 2nd. Pathology of the posterior tibial tendon in posterior tibial tendon insufficiency. *Foot Ankle Int.* 1998;19(8):520-524.
446. Paavola M, Kannus P, Jarvinen TA, Khan K, Jozsa L, Jarvinen M. Achilles tendinopathy. *J Bone Joint Surg Am.* 2002;84-A(11):2062-2076.
447. Andersson G, Danielson P, Alfredson H, Forsgren S. Nerve-related characteristics of ventral paratenondinous tissue in chronic Achilles tendinosis. *Knee Surg Sports Traumatol Arthrosc.* 2007;15(10):1272-1279.
448. de Jonge S, van den Berg C, de Vos RJ, et al. Incidence of midportion Achilles tendinopathy in the general population. *Br J Sports Med.* 2011;45(13):1026-1028. doi:10.1136/bjsports-2011-090342.
449. Klein EE, Weil L Jr, Weil LS Sr, Fleischer AE. Body mass index and Achilles tendinitis: a 10-year retrospective analysis. *Foot Ankle Spec.* 2013;6(4):276-282. doi:10.1177/1938640013489343.
450. Van Ginckel A, Thijss Y, Hesar NG, et al. Intrinsic gait-related risk factors for Achilles tendinopathy in novice runners: a prospective study. *Gait Posture.* 2009;29(3):387-391. doi:10.1016/j.gaitpost.2008.10.058.
451. Williams S, Riemann BL. Vertical leg stiffness following ankle taping and bracing. *Int J Sports Med.* 2009;30(5):383-386. doi:10.1055/s-0028-1105930.
452. Courville FC, Coe MP, Hecht PJ. Current concepts review: noninsertional Achilles tendinopathy. *Foot Ankle Int.* 2009;30(11):1132-1142. doi:10.3113/FAI.2009.1132.
453. Sayana MK, Maffulli N. Eccentric calf muscle training in non-athletic patients with Achilles tendinopathy. *J Sci Med Sport.* 2007;10(1):52-58.
454. Davenport TE, Kulig K, Matharu Y, Blanco CE. The EdUReP model for nonsurgical management of tendinopathy. *Phys Ther.* 2005;85(10):1093-1103.
455. Alfredson H, Lorentzon R. Chronic Achilles tendinosis: recommendations for treatment and prevention. *Sports Med.* 2000;29(2):135-146.
456. Williams DS, Zambardino JA, Banning VA. Transverse-plane mechanics at the knee and tibia in runners with and without a history of achilles tendonopathy. *J Orthop Sports Phys Ther.* 2008;38(12):761-767. doi:10.2519/jospt.2008.2911.

457. Young B, Walker MJ, Strunce J, Boyles R. A combined treatment approach emphasizing impairment-based manual physical therapy for plantar heel pain: a case series. *J Orthop Sports Phys Ther.* 2004;34(11):725-733.
458. Rees JD, Wolman RL, Wilson A. Eccentric exercises: why do they work, what are the problems and how can we improve them? *Br J Sports Med.* 2009;43(4):242-246. doi:10.1136/bjsm.2008.052910.
459. Kingma JJ, de Krijger R, Wittink HM, Takken T. Eccentric overload training in patients with chronic Achilles tendinopathy: a systematic review. *Br J Sports Med.* 2007;41(6):e3.
460. Jonsson P, Alfredson H, Sunding K, Fahlstrom M, Cook J. New regimen for eccentric calf-muscle training in patients with chronic insertional Achilles tendinopathy: results of a pilot study. *Br J Sports Med.* 2008;42(9):746-749. doi:10.1136/bjsm.2007.039545.
461. Beyer R, Kongsgaard M, Hougs Kjaer B, Ohlenschlaeger T, Kjaer M, Magnusson SP. Heavy slow resistance versus eccentric training as treatment for achilles tendinopathy: a randomized controlled trial. *Am J Sports Med.* 2015;43(7):1704-1711. doi:10.1177/0363546515584760.
462. Alfredson H, Pietila T, Jonsson P, Lorentzon R. Heavy-load eccentric calf muscle training for the treatment of chronic Achilles tendinosis. *Am J Sports Med.* 1998;26(3):360-366.
463. Stevens M, Tan CW. Effectiveness of the Alfredson protocol compared with a lower repetition-volume protocol for midportion Achilles tendinopathy: a randomized controlled trial. *J Orthop Sports Phys Ther.* 2014;44(2):59-67. doi:10.2519/jospt.2014.4720.
464. Rowe V, Hemmings S, Barton C, Malliaras P, Maffulli N, Morrissey D. Conservative management of midportion Achilles tendinopathy: a mixed methods study, integrating systematic review and clinical reasoning. *Sports Med.* 2012;42(11):941-967. doi:10.2165/11635410-00000000-0000.
465. de Jonge S, de Vos RJ, Van Schie HT, Verhaar JA, Weir A, Tol JL. One-year follow-up of a randomised controlled trial on added splinting to eccentric exercises in chronic midportion Achilles tendinopathy. *Br J Sports Med.* 2010;44(9):673-677. doi:10.1136/bjsm.2008.052142.
466. van der Plas A, de Jonge S, de Vos RJ, et al. A 5-year follow-up study of Alfredson's heel-drop exercise programme in chronic midportion Achilles tendinopathy. *Br J Sports Med.* 2012;46(3):214-218. doi:10.1136/bjsports-2011-090035.
467. McGuigan TC, Aierstok MD. Disorders of the Achilles tendon and its insertion. *Current Opinion in Orthopaedics.* 2005;16(2):65-71.
468. Wiegerinck JL, Kerkhoffs GM, van Sterkenburg MN, Siereveld IN, van Dijk CN. Treatment for insertional Achilles tendinopathy: a systematic review. *Knee Surg Sports Traumatol Arthrosc.* 2013;21(6):1345-1355. doi:10.1007/s00167-012-2219-8.
469. Nicholson CW, Berlet GC, Lee TH. Prediction of the success of nonoperative treatment of insertional Achilles tendinosis based on MRI. *Foot Ankle Int.* 2007;28(4):472-477.
470. Ganestam A, Kallemose T, Troelsen A, Barfod KW. Increasing incidence of acute Achilles tendon rupture and a noticeable decline in surgical treatment from 1994 to 2013. A nationwide registry study of 33,160 patients. *Knee Surg Sports Traumatol Arthrosc.* 2015.
471. Huttunen TT, Kannus P, Rolf C, Fellander-Tsai L, Mattila VM. Acute Achilles tendon ruptures: incidence of injury and surgery in Sweden between 2001 and 2012. *Am J Sports Med.* 2014;42(10):2419-2423. doi:10.1177/0363546514540599.
472. Chioldo CP, Glazebrook M, Bluman EM, et al. Diagnosis and treatment of acute Achilles tendon rupture. *J Am Acad Orthop Surg.* 2010;18(8):503-510.
473. Young SW, Patel A, Zhu M, et al. Weight-bearing in the nonoperative treatment of acute achilles tendon ruptures: a randomized controlled trial. *J Bone Joint Surg Am.* 2014;96(13):1073-1079.
474. van der Eng DM, Schepers T, Goslings JC, Schep NW. Rerupture rate after early weightbearing in operative versus conservative treatment of Achilles tendon ruptures: a meta-analysis. *J Foot Ankle Surg.* 2013;52(5):622-628. doi:10.1053/j.jfas.2013.03.027.
475. Holm C, Kjaer M, Eliasson P. Achilles tendon rupture-treatment and complications: a systematic review. *Scand J Med Sci Sports.* 2015;25(1):e1-10. doi:10.1111/sms.12209.
476. Brumman M, Baumbach SF, Mutschler W, Polzer H. Accelerated rehabilitation following Achilles tendon repair after acute rupture-development of an evidence-based treatment protocol. *Injury.* 2014;45(11):1782-1790. doi:10.1016/j.injury.2014.06.022.
477. Hutchison AM, Topliss C, Beard D, Evans RM, Williams P. The treatment of a rupture of the Achilles tendon using a dedicated management programme. *Bone Joint J.* 2015;97-B(4):510-515. doi:10.1302/0301-620X.97B4.35314.
478. O'Connor K, Baumhauer J, Houck JR. Patient factors in the selection of operative versus nonoperative treatment for posterior tibial tendon dysfunction. *Foot Ankle Int.* 2010;31(3):197-202. doi:10.3113/FAI.2010.0197.
479. Kohli-Gatzoulis JA, Sakellariou A. Re: Adult acquired flatfoot deformity following tibialposterior to dorsumtransfer: a case report. *Foot Ankle Int.* 2008;29(11):1168; author reply 1169. doi:10.3113/FAI.2008.1168.
480. Wacker J, Calder JD, Engstrom CM, Saxby TS. MR morphometry of posterior tibialis muscle in adult acquired flat foot. *Foot Ankle Int.* 2003;24(4):354-357.
481. Bluman EM, Title CI, Myerson MS. Posterior tibial tendon rupture: a refined classification system. *Foot Ankle Clin.* 2007;12(2):233-249, v.
482. Lin JL, Balbas J, Richardson EG, Lin JL, Balbas J, Richardson EG. Results of non-surgical treatment of stage II posterior tibial tendon dysfunction: a 7- to 10-year followup. *Foot Ankle Int.* 2008;29(8):781-786. doi:10.3113/FAI.2008.0781.
483. Alvarez RG, Marini A, Schmitt C, Saltzman CL. Stage I and II posterior tibial tendon dysfunction treated by a structured nonoperative management protocol: an orthosis and exercise program. *Foot Ankle Int.* 2006;27(1):2-8.
484. Neville CG, Houck JR. Choosing among 3 ankle-foot orthoses for a patient with stage II posterior tibial tendon dysfunction. *J Orthop Sports Phys Ther.* 2009;39(11):816-824. doi:10.2519/jospt.2009.3107.
485. Augustin JF, Lin SS, Berberian WS, Johnson JE. Nonoperative treatment of adult acquired flat foot with the Arizona brace. *Foot Ankle Clin.* 2003;8(3):491-502.
486. Bel N, Ozur A, Kaynak Y, Uygun F. The effect of orthotic treatment of posterior tibial tendon insufficiency on pain and disability. *Pain Clin.* 2003;15(3):345-350. doi:10.1163/156856903767650907
487. Kulig K, Reischl SF, Pomrancz AB, et al. Nonsurgical management of posterior tibial tendon dysfunction with orthoses and resistive exercise: a randomized controlled trial. *Phys Ther.* 2009;89(1):26-37. doi:10.2522/pj.20070242.
488. Kulig K, Lederhaus ES, Reischl S, Arya S, Bashford G. Effect of eccentric exercise program for early tibialis posterior tendinopathy. *Foot Ankle Int.* 2009;30(9):877-885. doi:10.3113/FAI.2009.0877.
489. Ziai P, Benca E, Wenzel F, et al. Peroneal tendinosis as a predisposing factor for the acute lateral ankle sprain in runners. *Knee Surg Sports Traumatol Arthrosc.* 2016;24(4):1175-1179.
490. Court-Brown CM, Caesar B. Epidemiology of adult fractures: A review. *Injury.* 2006;37(8):691-697.
491. Stevens JE, Pathare NC, Tillman SM, et al. Relative contributions of muscle activation and muscle size to plantarflexor torque during rehabilitation after immobilization. *J Orthop Res.* 2006;24(8):1729-1736.
492. Lin CW, Moseley AM, Reishauge KM. Rehabilitation for ankle fractures in adults. *Cochrane Database Syst Rev.* 2008(3):CD005595. doi:10.1002/14651858.CD005595.
493. Bessen T, Clark R, Shakib S, Hughes. A multifaceted strategy for implementation of the Ottawa ankle rules in two emergency departments. *BMJ.* 2009;339:b3056. doi:10.1136/bmj.b3056.
494. Dowling S, Spooner CH, Liang Y, et al. Accuracy of Ottawa Ankle Rules to exclude fractures of the ankle and midfoot in children: a meta-analysis. *Acad Emerg Med.* 2009;16(4):277-287. doi:10.1111/j.1535-2712.2008.00333.x
495. Smeeing DP, Houwert RM, Briet JP, et al. Weight-bearing and mobilization in the postoperative care of ankle fractures: a systematic review and meta-analysis of randomized controlled trials and cohort studies. *PLoS One.* 2015;10(2):e0118320. doi:10.1371/journal.pone.0118320.
496. Shibuya N, Davis ML, Jupiter DC. Epidemiology of foot and ankle fractures in the United States: an analysis of the National Trauma Data Bank (2007 to 2011). *J Foot Ankle Surg.* 2014;53(5):606-608. doi:10.1053/j.jfas.2014.03.011.
497. Olsen JR, Hunter J, Baumhauer JF. Osteoporotic ankle fractures. *Orthop Clin North Am.* 2013;44(2):225-241. doi:10.1016/j.ocl.2013.01.010.
498. Beckenkamp PR, Lin CW, Chagpar S, Herbert RD, van der Ploeg HP, Moseley AM. Prognosis of physical function following ankle fracture: a systematic review with meta-analysis. *J Orthop Sports Phys Ther.* 2014;44(11):841-851. doi:10.2519/jospt.2014.5199.
499. Hong CC, Roy SP, Nashi N, Tan KJ. Functional outcome and limitation of sporting activities after bimalleolar and trimalleolar ankle fractures. *Foot Ankle Int.* 2013;34(6):805-810. doi:10.1177/1071100712472490.
500. Egol KA, Tejwani NC, Walsh MG, Capla EL, Koval KJ. Predictors of short-term functional outcome following ankle fracture surgery. *J Bone Joint Surg Am.* 2006;88-A(5):974-979.
501. Stukens SA, van den Bekker MP, Kerkhoffs GM, Hintermann B, Van Dijk CN. Long-term outcome after 1822 operatively treated ankle fractures: a systematic review of the literature. *Injury.* 2011;42(2):119-127. doi:10.1016/j.injury.2010.04.006.
502. Segal G, Elbaz A, Parsi A, et al. Clinical outcomes following ankle fracture: a cross-sectional observational study. *J Foot Ankle Res.* 2014;7(1):50. doi:10.1186/s13047-014-0050-9.
503. Wang YC, Hart DL, Stratford PW, Mioduski JE. Clinical interpretation of computerized adaptive test outcome measures in patients with foot/ankle impairments. *J Orthop Sports Phys Ther.* 2009;39(10):753-764. doi:10.2519/jospt.2009.3122.
504. McPhail SM, Williams CM, Schuetz M, Baxter B, Tonks P, Haines TP. Development and validation of the ankle fracture outcome of rehabilitation measure (A-FORM). *J Orthop Sports Phys Ther.* 2014;44(7):488-499, B1-2. doi:10.2519/jospt.2014.4980.
505. Moseley AM, Herbert RD, Nightingale EJ, et al. Passive stretching does not enhance outcomes in patients with plantarflexion contracture after cast immobilization for ankle fracture: a randomized controlled trial. *Arch Phys Med Rehabil.* 2005;86(6):1118-1126.
506. Nightingale EJ, Moseley AM, Herbert RD. Passive dorsiflexion flexibility after cast immobilization for ankle fracture. *Clin Orthop Relat Res.* 2007;456:65-69.
507. Painter EE, Deyle GD, Allen C, Petersen EJ, Croy T, Rivera KP. Manual Physical Therapy Following Immobilization for Stable Ankle Fracture: A Case Series. *J Orthop Sports Phys Ther.* 2015;45(9):665-674. doi:10.2519/jospt.2015.5981.
508. Shaffer MA, Okereke E, Esterhai JL Jr, et al. Effects of immobilization on plantarflexion torque, fatigue resistance, and functional ability following an ankle fracture. *Phys Ther.* 2000;80(8):769-780.
509. Psathas M, Wu Z, Gammie FM, et al. A longitudinal MRI study of muscle atrophy during lower leg immobilization following ankle fracture. *J Magn Reson Imaging.* 2012;35(3):686-695. doi:10.1002/jmri.22864.
510. Stevens JE, Walter GA, Okereke E, et al. Muscle adaptations with immobilization and rehabilitation after ankle fracture. *Med Sci Sports Exerc.* 2004;36(10):1695-1701.
511. Nilsson GM, Jonsson K, Ekholm CS, Eneroth M. Effects of a training program after surgically treated ankle fracture: a prospective randomised controlled trial. *BMC Musculoskeletal Disord.* 2009;10:118. doi:10.1186/1471-2474-10-118.
512. Dudek K, Druzbicki M, Przysada G, Spiewak D. Assessment of standing balance in patients after ankle fractures. *Acta Bioeng Biomech.* 2014;16(4):59-65.

513. Nilsson G, Ageberg E, Ekdahl C, Eneroth M. Balance in single-limb stance after surgically treated ankle fractures: a 14-month follow-up. *BMC Musculoskelet Disord.* 2006;7:35.
514. Keene DJ, Williamson E, Bruce J, Willett K, Lamb SE. Early ankle movement versus immobilization in the postoperative management of ankle fracture in adults: a systematic review and meta-analysis. *J Orthop Sports Phys Ther.* 2014;44(9):690-701, C691-697. doi:10.2519/jospt.2014.5294.
515. Lin CW, Donkers NA, Refshauge KM, Beckenkamp PR, Khera K, Moseley AM. Rehabilitation for ankle fractures in adults. *Cochrane Database Syst Rev.* 2012;11:CD005595. doi:10.1002/14651858.CD005595.
516. Firoozabadi R, Harnden E, Krieg JC. Immediate weight-bearing after ankle fracture fixation. *Adv Orthop.* 2015;2015:491976. doi:10.1155/2015/491976.
517. Porter DA, May BD, Berney T. Functional outcome after operative treatment for ankle fractures in young athletes: a retrospective case series. *Foot Ankle Int.* 2008;29(9):887-894. doi:10.3113/FAI.2008.0887.
518. Briet JP, Houwert RM, Smeling DP, et al. Weight bearing or non-weight bearing after surgically fixed ankle fractures, the WOW! Study: study protocol for a randomized controlled trial. *Trials.* 2015;16(1):175. doi:10.1186/s13063-015-0714-1.
519. Lin CW, Haas M, Moseley AM, Herbert RD, Refshauge KM. Cost and utilisation of healthcare resources during rehabilitation after ankle fracture are not linked to health insurance, income, gender, or pain: an observational study. *Aust J Physiother.* 2008;54(3):201-208.
520. Lin CW, Moseley AM, Herbert RD, Refshauge KM. Pain and dorsiflexion range of motion predict short- and medium-term activity limitation in people receiving physiotherapy intervention after ankle fracture: an observational study. *Aust J Physiother.* 2009;55(1):31-37.
521. Lin CW, Moseley AM, Haas M, Refshauge KM, Herbert RD. Manual therapy in addition to physiotherapy does not improve clinical or economic outcomes after ankle fracture. *J Rehabil Med.* 2008;40(6):433-439. doi:10.2340/16501977-0187.
522. Beckenkamp PR, Lin CC, Herbert RD, Haas M, Khera K, Moseley AM. EXACT: exercise or advice after ankle fracture. Design of a randomised controlled trial. *BMC Musculoskelet Disord.* 2011;12:148. doi:10.1186/1471-2474-12-148.
523. Davidovitch RI, Walsh M, Spitzer A, Egol KA. Functional outcome after operatively treated ankle fractures in the elderly. *Foot Ankle Int.* 2009;30(8):728-733. doi:10.3113/FAI.2009.0728.
524. Buckley R. Calcaneal fractures: to fix or not to fix: Opinion: nonoperative approach. *J Orthop Trauma.* 2005;19(5):357-359.
525. Gougloulias N, Khanna A, McBride DJ, Maffulli N. Management of calcaneal fractures: systematic review of randomized trials. *Br Med Bull.* 2009;92:153-167. doi:10.1093/bmb/ldp030.
526. Rammelt S, Zwipp H. Calcaneus fractures: facts, controversies and recent developments. *Injury.* 2004;35(5):443-461.
527. Romeo G, Martinelli N, Bonifacini C, Bianchi A, Sartorelli E, Malerba F. Recreational sports activities after calcaneal fractures and subsequent subtalar joint arthrodesis. *J Foot Ankle Surg.* 2015;54(6):1057-1061. doi:10.1053/j.jfas.2015.05.003.
528. O'Brien J, Buckley R, McCormack R, et al. Personal gait satisfaction after displaced intraarticular calcaneal fractures: a 2-8 year followup. *Foot Ankle Int.* 2004;25(9):657-665.
529. Harrast MA, Colombo D. Stress fractures in runners. *Clin Sports Med.* 2010;29(3):399-416. doi:10.1016/j.csm.2010.03.001.
530. Zadpoor AA, Nikooyan AA. The relationship between lower-extremity stress fractures and the ground reaction force: a systematic review. *Clin Biomech (Bristol, Avon).* 2011;26(1):23-28. doi:10.1016/j.clinbiomech.2010.08.005.
531. Zifchock RA, Davis I, Hamill J. Kinetic asymmetry in female runners with and without retrospective tibial stress fractures. *J Biomed.* 2006;39(15):2792-2797.
532. Anderson RB, Hunt KJ, McCormick JJ. Management of common sports-related injuries about the foot and ankle. *J Am Acad Orthop Surg.* 2010;18(9):546-556.
533. Sueki D, Brechter J. *Orthopedic Rehabilitation Clinical Advisor.* Vol 1 ed. The Netherlands: Mosby; 2009.
534. Yates B, Allen MJ, Barnes MR. Outcome of surgical treatment of medial tibial stress syndrome. *J Bone Joint Surg Am.* 2003;85-A(10):1974-1980.
535. Beck BR. Tibial stress injuries. An aetiological review for the purposes of guiding management. *Sports Med.* 1998;26(4):265-279.
536. Tweed JL, Campbell JA, Avil SJ. Biomechanical risk factors in the development of medial tibial stress syndrome in distance runners. *J Am Podiatr Med Assoc.* 2008;98(6):436-444.
537. Yates B, White S. The incidence and risk factors in the development of medial tibial stress syndrome among naval recruits. *Am J Sports Med.* 2004;32(3):772-780.
538. Detmer DE. Chronic shin splints. Classification and management of medial tibial stress syndrome. *Sports Med.* 1986;3(6):436-446.
539. Loudon JK, Dolphino MR. Use of foot orthoses and calf stretching for individuals with medial tibial stress syndrome. *Foot Ankle Spec.* 2010;3(1):15-20. doi:10.1177/1938640009355659.
540. Plisky MS, Rauh MJ, Heiderscheit B, Underwood FB, Tank RT. Medial tibial stress syndrome in high school cross-country runners: incidence and risk factors. *J Orthop Sports Phys Ther.* 2007;37(2):40-47.
541. Moen MH, Tol JL, Weir A, Steunebrink M, De Winter TC. Medial tibial stress syndrome: a critical review. *Sports Med.* 2009;39(7):523-546. doi:10.2165/00007256-200939070-00002.
542. Bennett JE, Reinking MF, Pluemer B, Pentel A, Seaton M, Killian C. Factors contributing to the development of medial tibial stress syndrome in high school runners. *J Orthop Sports Phys Ther.* 2001;31(9):504-510.
543. Tweed JL, Avil SJ, Campbell JA, Barnes MR. Etiologic factors in the development of medial tibial stress syndrome: a review of the literature. *J Am Podiatr Med Assoc.* 2008;98(2):107-111.
544. Beck BR, Osterling LR. Medial tibial stress syndrome. The location of muscles in the leg in relation to symptoms. *J Bone Joint Surg Am.* 1994;76(7):1057-1061.
545. Bouche RT, Johnson CH. Medial tibial stress syndrome (tibial fasciitis): a proposed pathomechanical model involving fascial traction. *J Am Podiatr Med Assoc.* 2007;97(1):31-36.
546. Saxena A, O'Brien T, Bunce D. Anatomic dissection of the tibialis posterior muscle and its correlation to medial tibial stress syndrome. *J Foot Surg.* 1990;29(2):105-108.
547. Magnusson HI, Westlin NE, Nyqvist F, Gardsell P, Seeman E, Karlsson MK. Abnormally decreased regional bone density in athletes with medial tibial stress syndrome. *Am J Sports Med.* 2001;29(6):712-715.
548. Hamstra-Wright KL, Bliven KC, Bay C. Risk factors for medial tibial stress syndrome in physically active individuals such as runners and military personnel: a systematic review and meta-analysis. *Br J Sports Med.* 2015;49(6):362-369. doi:10.1136/bjsports-2014-093462.
549. Craig DL. Medial tibial stress syndrome: evidence-based prevention. *J Athl Train.* 2008;43(3):316-318. doi:10.4085/1062-6050-43.3.316.
550. Galbraith RM, Lavallee ME. Medial tibial stress syndrome: conservative treatment options. *Curr Rev Musculoskelet Med.* 2009;2(3):127-133. doi:10.1007/s12178-009-9055-6.
551. Pell RF 4th, Khanuja HS, Cooley GR. Leg pain in the running athlete. *J Am Acad Orthop Surg.* 2004;12(6):396-404.
552. Winters M, Eskes M, Weir A, Moen MH, Backx FJ, Bakker EW. Treatment of medial tibial stress syndrome: a systematic review. *Sports Med.* 2013;43(12):1315-1333. doi:10.1007/s40279-013-0087-0.
553. Solomon LB, Ruhli FJ, Taylor J, Ferris L, Pope R, Henneberg M. A dissection and computer tomograph study of tarsal coalitions in 100 cadaver feet. *J Orthop Res.* 2003;21(2):352-358.
554. Wickiewicz TL, Roy RR, Powell PL, Edgerton VR. Muscle architecture of the human lower limb. *Clin Orthop Relat Res.* 1983;(179):275-283.
555. Fukunaga T, Roy RR, Shellock FG, et al. Physiological cross-sectional area of human leg muscles based on magnetic resonance imaging. *J Orthop Res.* 1992;10(6):928-934.
556. Friederich JA, Brand RA. Muscle fiber architecture in the human lower limb. *J Biomech.* 1990;23(1):91-95.
557. Allard P, Stokes I, Blanck JP. *Three-Dimensional Analysis of Human Movement.* Champaign, IL: Human Kinetics; 1995.
558. Silver RL, de la Garza J, Rang M. The myth of muscle balance. A study of relative strengths and excursions of normal muscles about the foot and ankle. *J Bone Joint Surg Br.* 1985;67(3):432-437.
559. Chuckpaiwong B, Nunley JA 2nd, Queen RM. Correlation between static foot type measurements and clinical assessments. *Foot Ankle Int.* 2009;30(3):205-212. doi:10.3113/FAI.2009.0205.
560. Genova JM, Gross MT. Effect of foot orthotics on clacaneal eversion during standing and treadmill walking for subjects with abnormal pronation. *J Orthop Sports Phys Ther.* 2000;30(11):664-675.
561. Brushoj C, Larsen K, Albrecht-Beste E, Nielsen MB, Loye F, Holmich P. Prevention of overuse injuries by a concurrent exercise program in subjects exposed to an increase in training load: a randomized controlled trial of 1020 army recruits. *Am J Sports Med.* 2008;36(4):663-670. doi:10.1177/0363546508315469.
562. Buchanan KR, Davis I. The relationship between forefoot, midfoot, and rearfoot static alignment in pain-free individuals. *J Orthop Sports Phys Ther.* 2005;35(9):559-566.
563. Tsai LC, Yu B, Mercer VS, Gross MT. Comparison of different structural foot types for measures of standing postural control. *J Orthop Sports Phys Ther.* 2006;36(12):942-953.
564. Cavanagh PR, Rodgers MM. The arch index: a useful measure from footprints. *J Biomech.* 1987;20(5):547-551.
565. Olmsted LC, Carcia CR, Hertel J, Shultz SJ. Efficacy of the Star Excursion Balance Tests in Detecting Reach Deficits in Subjects with Chronic Ankle Instability. *J Athl Train.* 2002;37(4):501-506.
566. Lunsford BR, Perry J. The standing heel-rise test for ankle plantar flexion: criterion for normal. *Phys Ther.* 1995;75(8):694-698.
567. Jan MH, Chai HM, Lin YF, et al. Effects of age and sex on the results of an ankle plantar-flexor manual muscle test. *Phys Ther.* 2005;85(10):1078-1084.
568. Moseley AM, Crosbie J, Adams R. Normative data for passive ankle plantarflexion--dorsiflexion flexibility. *Clin Biomech (Bristol, Avon).* 2001;16(6):514-521.
569. Jonson SR, Gross MT. Intraexaminer reliability, interexaminer reliability, and mean values for nine lower extremity skeletal measures in healthy naval midshipmen. *J Orthop Sports Phys Ther.* 1997;25(4):253-263.