



Current Concepts of **Orthopaedic Physical Therapy**

Independent Study Course 26.2.9

4th Edition

**The Pelvis and Sacroiliac
Joint: Physical Therapy
Patient Management
Using Current
Evidence**

Richard Jackson, PT, OCS
Kris Porter, PT, DPT, OCS
The Jackson Clinics, LP
Middleburg, Virginia



REFERENCES

1. Vleeming A, Schuenke MD, Masi AT, Carreiro JE, Daniels L, Willard FH. The sacroiliac joint: An overview of its anatomy, function and potential clinical implications. *J Anat.* 2012;221(6):537–567. doi:10.1111/j.1469-7580.2012.01564.x. Epub 2012 Sep 19.
2. Huang BK, Campos JC, Michael Peschka PG, et al. Injury of the gluteal aponeurotic fascia and proximal iliotibial band: anatomy, pathologic conditions, and MR imaging. *Radiographics.* 2013;33(5):1437–1552. doi:10.1148/rg.335125171.
3. Sung PS. A kinematic analysis for shoulder and pelvis coordination during axial trunk rotation in subjects with and without recurrent low back pain. *Gait Posture.* 2014;40(4):493–498. doi:10.1016/j.gaitpost.2014.06.001.
4. Barker PJ, Hapuarachchi KS, Ross JA, Sambaiw E, Ranger TA, Briggs CA. Anatomy and biomechanics of gluteus maximus and the thoracolumbar fascia at the sacroiliac joint. *Clin Anat.* 2014;27(2):234–240. doi:10.1002/ca.22233.
5. Antonio S, Wolfgang G, Robert H, Fullerton B, Carla S. The anatomical and functional relation between gluteus maximus and fascia lata. *J Bodyw Mov Ther.* 2013;17(4):512–517. doi:10.1016/j.jbmt.2013.04.004.
6. Yang YT, Yoshida Y, Hortobágyi T, Suzuki S. Interaction between thorax, lumbar, and pelvis movements in the transverse plane during gait at three velocities. *J Appl Biomech.* 2013;29(3):261–269.
7. Aldabe D, Milosavljevic S, Bussey MD. Is pregnancy related pelvic girdle pain associated with altered kinematic, kinetic and motor control of the pelvis? A systematic review. *Eur Spine J.* 2012;21(9):1777–1787. doi:10.1007/s00586-012-2401-1.
8. Bussey MD, Milosavljevic S. Asymmetric pelvic bracing and altered kinematics in patients with posterior pelvic pain who present with postural muscle delay. *Clin Biomech.* 2015;30(1):71–77. doi:10.1016/j.clinbiomech.2014.11.002.
9. Waddell G. *The Back Pain Revolution.* 2nd ed. Edinburgh; New York, NY: Churchill Livingstone; 2007.
10. Khamis S, Yizhar Z. Effect of feet hyperpronation on pelvic alignment in a standing position. *Gait Posture.* 2007;25(1):127–134. doi:10.1016/j.gaitpost.2006.02.005.
11. Vleeming A, Stoeckart R. The role of the pelvic girdle in coupling the spine and the legs: a clinical-anatomical perspective on pelvic stability. In: Vleeming A, Mooney V, Stoeckart R, eds. *Movement, Stability and Lumbopelvic Pain: Integration of Research and Therapy.* 2nd ed. Edinburgh; New York, NY: Churchill Livingstone Elsevier; 2007:113–137.
12. Marshall PWM, McKee AD, Murphy BA. Impaired trunk and ankle stability in subjects with functional ankle instability. *Med Sci Sports Exerc.* 2009;41(8):1549–1557. doi:10.1249/MSS.0b013e31819d82e2.
13. Willson JD, Dougherty CP, Ireland ML, Davis IM. Core stability and its relationship to lower extremity function and injury. *J Am Acad Orthop Surg.* 2005;13(5):316–325.
14. Cibulka MT. Low back pain and its relation to the hip and foot. *J Orthop Sports Phys Ther.* 1999;29(10):595–601.
15. Kibler W Ben, Press J, Sciascia A. The role of core stability in athletic function. *Sports Med.* 2006;36(3):189–198.
16. Schwab F, Lafage V, Patel A, Farcy JP. Sagittal plane considerations and the pelvis in the adult patient. *Spine (Phila Pa 1976).* 2009;34(17):1828–1833. doi:10.1097/BRS.0b013e3181a13c08.
17. Leetun DT, Ireland ML, Willson JD, Ballantyne BT, Davis IM. Core stability measures as risk factors for lower extremity injury in athletes. *Med Sci Sports Exerc.* 2004;36(6):926–934.
18. Geraci MC, Brown W. Evidence-based treatment of hip and pelvic injuries in runners. *Phys Med Rehabil Clin N Am.* 2005;16(3):711–747. doi:10.1016/j.pmr.2005.02.004.
19. Goode A, Hegedus EJ, Sizer P, Brismee J-M, Linberg A, Cook CE. Three-dimensional movements of the sacroiliac joint: a systematic review of the literature and assessment of clinical utility. *J Man Manip Ther.* 2008;16(1):25–38.
20. Snijders CJ, Vleeming A, Stoeckart R. Transfer of lumbosacral load to iliac bones and legs. Part 1: Biomechanics of self-bracing of the sacroiliac joints and its significance for treatment and exercise. *Clin Biomech.* 1993;8(6):285–294. doi:10.1016/0268-0033(93)90002-Y.
21. Snijders CJ, Vleeming A, Stoeckart R. Transfer of lumbosacral load to iliac bones and legs. Part 2: Loading of the sacroiliac joints when lifting in a stooped posture. *Clin Biomech.* 1993;8(6):295–301. doi:10.1016/0268-0033(93)90003-Z.
22. Hoek van Dijke GA, J. Snijders C, Stoeckart R, Stam HJ. A biomechanical model on muscle forces in the transfer of spinal load to the pelvis and legs. *J Biomech.* 1999;32(9):927–933.
23. Vleeming A, Pool-Goudzwaard AL, Stoeckart R, van Wingerden JP, Snijders CJ. The posterior layer of the thoracolumbar fascia. Its function in load transfer from spine to legs. *Spine (Phila Pa 1976).* 1995;20(7):753–758.
24. Erhard R, Bowling R. The recognition and management of the pelvic component of low back and sciatic pain. *Bulletin of the Orthopaedic Section.* 1977;2(3):4–14.
25. Vleeming A, Albert HB, Ostgaard HC, et al. European guidelines for the diagnosis and treatment of pelvic girdle pain. *Eur Spine J.* 2008;17(6):794–819. doi:10.1007/s00586-008-0602-4.
26. Hancock MJ, Maher CG, Latimer J, et al. Systematic review of tests to identify the disc, SIJ or facet joint as the source of low back pain. *Eur Spine J.* 2007;16(10):1539–1550. doi:10.1007/s00586-007-0391-1.
27. Slipman CW, Jackson HB, Lipetz JS, Chan KT, Lenrow D, Vresilovic EH. Sacroiliac joint pain referral zones. *Arch Phys Med Rehabil.* 2000;81(3):334–338.
28. van der Wurff P, Buijs EJ, Groen GJ. Intensity mapping of pain referral areas in sacroiliac joint pain patients. *J Manipulative Physiol Ther.* 2006;29(3):190–195. doi:10.1016/j.jmpt.2006.01.007.

29. Laslett M, April CN, McDonald B. Provocation sacroiliac joint tests have validity in the diagnosis of sacroiliac joint pain. *Arch Phys Med Rehabil.* 2006;87(6):874–875.
30. Berthelot JM, Labat JJ, LeGoff B, Gouin F, Maugars Y. Provocative sacroiliac joint maneuvers and sacroiliac joint block are unreliable for diagnosing sacroiliac joint pain. *Joint Bone Spine.* 2006;73(1):17–23. doi:10.1016/j.jbspin.2004.08.003.
31. Hungerford B, Gillear W, Hodges P. Evidence of altered lumbopelvic muscle recruitment in the presence of sacroiliac joint pain. *Spine (Phila Pa 1976).* 2003;28(14):1593–1600.
32. O'Sullivan PB, Twomey L, Allison GT. Altered abdominal muscle recruitment in patients with chronic back pain following a specific exercise intervention. *J Orthop Sports Phys Ther.* 1998;27(2):114–124.
33. Hodges PW, Richardson CA. Altered trunk muscle recruitment in people with low back pain with upper limb movement at different speeds. *Arch Phys Med Rehabil.* 1999;80(9):1005–1012.
34. Ferreira PH, Ferreira ML, Hodges PW. Changes in recruitment of the abdominal muscles in people with low back pain: ultrasound measurement of muscle activity. *Spine (Phila Pa 1976).* 2004;29(22):2560–2566.
35. Silfies SP, Mehta R, Smith SS, Karduna AR. Differences in feedforward trunk muscle activity in subgroups of patients with mechanical low back pain. *Arch Phys Med Rehabil.* 2009;90(7):1159–1169. doi:10.1016/j.apmr.2008.10.033.
36. Hodges PW, Richardson CA. Delayed postural contraction of transversus abdominis in low back pain associated with movement of the lower limb. *J Spinal Disord.* 1998;11(1):46–56.
37. Hodges PW, Richardson CA. Inefficient muscular stabilization of the lumbar spine associated with low back pain. A motor control evaluation of transversus abdominis. *Spine (Phila Pa 1976).* 1996;21(22):2640–2650.
38. Cowan SM, Schache AG, Brukner P, et al. Delayed onset of transversus abdominus in long-standing groin pain. *Med Sci Sports Exerc.* 2004;36(12):2040–2045.
39. Cholewicki J, Silfies SP, Shah R a, et al. Delayed trunk muscle reflex responses increase the risk of low back injuries. *Spine (Phila Pa 1976).* 2005;30(23):2614–2620.
40. van Wingerden J-P, Vleeming A, Ronchetti I. Differences in standing and forward bending in women with chronic low back or pelvic girdle pain: indications for physical compensation strategies. *Spine (Phila Pa 1976).* 2008;33(11):E334–341. doi:10.1097/BRS.0b013e-318170fcf6.
41. Porter JL, Wilkinson A. Lumbar-hip flexion motion. A comparative study between asymptomatic and chronic low back pain in 18- to 36-year-old men. *Spine (Phila Pa 1976).* 1997;22(13):1508–1513; discussion 1513–1514.
42. Beales DJ, O'Sullivan PB, Briffa NK. Motor control patterns during an active straight leg raise in chronic pelvic girdle pain subjects. *Spine (Phila Pa 1976).* 2009;34(9):861–870. doi:10.1097/BRS.0b013e318198d212.
43. Anders C, Scholle HC, Wagner H, Puta C, Grassme R, Petrovitch A. Trunk muscle co-ordination during gait: relationship between muscle function and acute low back pain. *Pathophysiol Off J Int Soc Pathophysiol / ISP.* 2005;12(4):243–247. doi:10.1016/j.pathophys.2005.09.001.
44. Delphinus EM, Sayers MGL. The interrelationship of the thorax and pelvis under varying task constraints. *Ergonomics.* 2013;56(4):659–666. doi:10.1080/00140139.2012.760755.
45. Dreyfuss P, Michaelsen M, Pauza K, McLarty J, Bogduk N. The value of medical history and physical examination in diagnosing sacroiliac joint pain. *Spine (Phila Pa 1976).* 1996;21(22):2594–2602.
46. Bussey MD, Bell ML, Milosavljevic S. The influence of hip abduction and external rotation on sacroiliac motion. *Man Ther.* 2009;14(5):520–525. doi:10.1016/j.math.2008.08.009.
47. Adhia DB, Bussey MD, Ribeiro DC, Tumilty S, Milosavljevic S. Validity and reliability of palpation-digitization for non-invasive kinematic measurement - a systematic review. *Man Ther.* 2013;18(1):26–34. doi:10.1016/j.math.2012.06.004.
48. Adhia DB, Bussey MD, Mani R, Jayakaran P, Aldabe D, Milosavljevic S. Inter-tester reliability of non-invasive technique for measurement of innominate motion. *Man Ther.* 2012;17(1):71–76. doi:10.1016/j.math.2011.09.005.
49. Adhia DB, Mani R, Milosavljevic S, Tumilty S, Bussey MD. Does repeated palpation-digitization of pelvic landmarks for measurement of innominate motion introduce a systematic error? – A psychometric investigation. *Man Ther.* 2015;21:282–286. doi:10.1016/j.math.2015.09.009.
50. Adhia DB, Milosavljevic S, Tumilty S, Bussey MD. Innominate movement patterns, rotation trends and range of motion in individuals with low back pain of sacroiliac joint origin. *Man Ther.* 2015;21:100–108. doi:10.1016/j.math.2015.06.004.
51. Robinson HS, Brox JI, Robinson R, Bjelland E, Solem S, Telje T. The reliability of selected motion-and pain provocation tests for the sacroiliac joint. *Man Ther.* 2007;12(1):72–79. doi:10.1016/j.math.2005.09.004.
52. Dreyfuss P, Dryer S, Griffin J, Hoffman J, Walsh N. Positive sacroiliac screening tests in asymptomatic adults. *Spine (Phila Pa 1976).* 1994;19(10):1138–1143.
53. Meijne W, van Neerbos K, Aufdemkampe G, van der Wurff P. Intraexaminer and interexaminer reliability of the Gillet test. *J Manipulative Physiol Ther.* 1999;22(1):4–9.
54. Vincent-Smith B, Gibbons P. Inter-examiner and intra-examiner reliability of the standing flexion test. *Man Ther.* 1999;4(2):87–93. doi:10.1054/math.1999.0173.
55. Levangie PK. Four clinical tests of sacroiliac joint dysfunction: the association of test results with innominate torsion among patients with and without low back pain. *Phys Ther.* 1999;79(11):1043–1057.
56. Riddle DL, Freburger JK, others. Evaluation of the presence of sacroiliac joint region dysfunction using a combination of tests: a multicenter intertester reliability study. *Phys Ther.* 2002;82(8):772.
57. van der Wurff P, Hagmeijer RH, Meyne W. Clinical tests of the sacroiliac joint. A systematic methodological review. Part 1: Reliability. *Man Ther.* 2000;5(1):30–36. doi:10.1054/math.1999.0228.
58. Potter NA, Rothstein JM. Intertester reliability for selected clinical tests of the sacroiliac joint. *Phys Ther.* 1985;65(11):1671–1675.

59. Freburger JK, Riddle DL. Measurement of sacroiliac joint dysfunction: a multicenter intertester reliability study. *Phys Ther*. 1999;79(12):1134–1141.
60. Herzog W, Read LJ, Conway PJ, Shaw LD, McEwen MC. Reliability of motion palpation procedures to detect sacroiliac joint fixations. *J Manipulative Physiol Ther*. 1989;12(2):86–92.
61. Laslett M, Huijbregts PA, Laslett M. Evidence-based diagnosis and treatment of the painful sacroiliac joint. *J Man Manip Ther*. 2008;16(3):142–152.
62. O'Haire C, Gibbons P. Inter-examiner and intra-examiner agreement for assessing sacroiliac anatomical landmarks using palpation and observation: pilot study. *Man Ther*. 2000;5(1):13–20. doi:10.1054/math.1999.0203.
63. DeRosa C, Porterfield JA. Anatomical linkages and muscle slings of the lumbopelvic region. In: Vleeming A, Mooney V, Stoeckart R, eds. *Movement, Stability and Lumbopelvic Pain: Integration of Research and Therapy*. 2nd ed. Edinburgh; New York, NY: Churchill Livingstone Elsevier; 2007:47–62.
64. Willard F. The muscular, ligamentous, and neural structure of the lumbosacrum and its relationship to low back pain. In: Vleeming A, Mooney V, Stoeckart R, eds. *Movement, Stability and Lumbopelvic Pain: Integration of Research and Therapy*. 2nd ed. Edinburgh; New York, NY: Churchill Livingstone Elsevier; 2007:5–45.
65. Hodges PW, Richardson CA. Contraction of the abdominal muscles associated with movement of the lower limb. *Phys Ther*. 1997;77(2):132–142; discussion 142–144.
66. Vleeming A, Mooney V, Stoeckart R, et al. *Movement, Stability & Lumbopelvic Pain: Integration of Research and Therapy*. 2nd ed. Edinburgh; New York, NY: Churchill Livingstone Elsevier; 2007.
67. Hodges PW, Cholewicki J. Functional control of the spine. In: Vleeming A, Mooney V, Stoeckart R, eds. *Movement, Stability and Lumbopelvic Pain: Integration of Research and Therapy*. 2nd ed. Edinburgh; New York, NY: Churchill Livingstone Elsevier; 2007:489–512.
68. Cholewicki J, McGill SM. Mechanical stability of the in vivo lumbar spine: implications for injury and chronic low back pain. *Clin Biomech (Bristol, Avon)*. 1996;11(1):1–15.
69. McGill SM, Grenier S, Kavcic N, Cholewicki J. Coordination of muscle activity to assure stability of the lumbar spine. *J Electromyogr Kinesiol*. 2003;13(4):353–359.
70. Gracovetsky S. Stability or controlled instability? In: Vleeming A, Mooney V, Stoeckart R, eds. *Movement, Stability & Lumbopelvic Pain: Integration of Research and Therapy*. 2nd ed. Edinburgh; New York, NY: Churchill Livingstone Elsevier; 2007.
71. Danneels LA, Vanderstraeten GG, Cambier DC, Witvrouw EE, Stevens VK, De Cuyper HJ. A functional subdivision of hip, abdominal, and back muscles during asymmetric lifting. *Spine (Phila Pa 1976)*. 2001;26(6):E114–121.
72. Comerford MJ, Mottram SL. Functional stability re-training: principles and strategies for managing mechanical dysfunction. *Man Ther*. 2001;6(1):3–14. doi:10.1054/math.2000.0389.
73. Cholewicki J, VanVliet JJ. Relative contribution of trunk muscles to the stability of the lumbar spine during isometric exertions. *Clin Biomech (Bristol, Avon)*. 2002;17(2):99–105.
74. Bergmark A. Stability of the lumbar spine. A study in mechanical engineering. *Acta Orthop Scand Suppl*. 1989;230:1–54.
75. Pool-Goudzwaard A, Hoek van Dijke G, Mulder P, Spoor C, Snijders C, Stoeckart R. The iliolumbar ligament: its influence on stability of the sacroiliac joint. *Clin Biomech*. 2003;18(2):99–105.
76. Luk KD, Ho HC, Leong JC. The iliolumbar ligament. A study of its anatomy, development and clinical significance. *J Bone Joint Surg Br*. 1986;68(2):197–200.
77. Leong JC, Luk KD, Chow DH, Woo CW. The biomechanical functions of the iliolumbar ligament in maintaining stability of the lumbosacral junction. *Spine (Phila Pa 1976)*. 1987;12(7):669–674.
78. Yamamoto I, Panjabi MM, Oxland TR, Crisco JJ. The role of the iliolumbar ligament in the lumbosacral junction. *Spine (Phila Pa 1976)*. 1990;15(11):1138–1141.
79. Chow DH, Luk KD, Leong JC, Woo CW. Torsional stability of the lumbosacral junction. Significance of the iliolumbar ligament. *Spine (Phila Pa 1976)*. 1989;14(6):611–615.
80. Badii M, Shin S, Torreggiani WC, et al. Pelvic bone asymmetry in 323 study participants receiving abdominal CT scans. *Spine (Phila Pa 1976)*. 2003;28(12):1335–1339. doi:10.1097/01.BRS.0000065480.44620.C5.
81. van de Meent H, Jansen H, van der Linde H. Study of the human pelvis using CAT-scan: gender differences and anatomy of the ramus ossis ischii. *Prosthet Orthot Int*. 2008;32(4):385–389. doi:10.1080/03093640802258629.
82. Vleeming A, Stoeckart R, Volkers AC, Snijders CJ. Relation between form and function in the sacroiliac joint. Part I: Clinical anatomical aspects. *Spine (Phila Pa 1976)*. 1990;15(2):130–132.
83. Vleeming A, Volkers AC, Snijders CJ, Stoeckart R. Relation between form and function in the sacroiliac joint. Part II: Biomechanical aspects. *Spine (Phila Pa 1976)*. 1990;15(2):133–136.
84. Ingber DE. Tensegrity and mechanotransduction. *J Bodyw Mov Ther*. 2008;12(3):198–200. doi:10.1016/j.jbmt.2008.04.038.
85. Levin S. A suspensory system for the sacrum in pelvic mechanics: biotensegrity. In: *Movement, Stability & Lumbopelvic Pain : Integration of Research and Therapy*. 2nd ed. Vleeming A, Mooney V, Stoeckart R, eds. Edinburgh; New York, NY: Churchill Livingstone Elsevier; 2007.
86. Abitbol MM. Evolution of the sacrum in hominoids. *Am J Phys Anthropol*. 1987;74(1):65–81. doi:10.1002/ajpa.1330740107.
87. van Wingerden JP, Vleeming A, Buyruk HM, Raissadat K. Stabilization of the sacroiliac joint in vivo: verification of muscular contribution to force closure of the pelvis. *Eur Spine J*. 2004;13(3):199–205. doi:10.1007/s00586-003-0575-2.
88. Dalstra M, Huiskes R. Load transfer across the pelvic bone. *J Biomech*. 1995;28(6):715–724.
89. Pel JJM, Spoor CW, Pool-Goudzwaard AL, Hoek van Dijke GA, Snijders CJ. Biomechanical analysis of reducing sacroiliac joint shear load by optimization of pelvic muscle and ligament forces. *Ann Biomed Eng*. 2008;36(3):415–424. doi:10.1007/s10439-007-9385-8.

90. Longo UG, Loppini M, Berton a., Laverde I, Maffulli N, Denaro V. Degenerative changes of the sacroiliac joint after spinal fusion: an evidence-based systematic review. *Br Med Bull.* 2014;112(1):47–56. doi:10.1093/bmb/ldu030.
91. Ciarpaglini R, Otten P, Sutter P, Duy VQ, Gautier E, Mae-stretti G. Sacroiliac joint syndrome 10 years after lumbar arthroplasty: the importance of spinopelvic alignment. *Eur Spine J.* 2014;23(S6):720–724. doi:10.1007/s00586-014-3547-9.
92. Mens JMA, Vleeming A, Snijders CJ, Koes BW, Stam HJ. Validity of the active straight leg raise test for measuring disease severity in patients with posterior pelvic pain after pregnancy. *Spine (Phila Pa 1976).* 2002;27(2):196–200.
93. Shadmehr A, Jafarian Z, Talebian S. Changes in recruitment of pelvic stabilizer muscles in people with and without sacroiliac joint pain during the active straight-leg-raise test. *J Back Musculoskelet Rehabil.* 2012;25(1):27–32. doi:10.3233/BMR-2012-0307.
94. Maigne JY, Aivaliklis A, Pfefer F. Results of sacroiliac joint double block and value of sacroiliac pain provocation tests in 54 patients with low back pain. *Spine (Phila Pa 1976).* 1996;21(16):1889–1892.
95. Gemmell HA, Jacobson BH. Incidence of sacroiliac joint dysfunction and low back pain in fit college students. *J Manipulative Physiol Ther.* 1990;13(2):63–67.
96. Madani SP, Dadian M, Firouznia K, Alalawi S. Sacroiliac joint dysfunction in patients with herniated lumbar disc: A cross-sectional study. *J Back Musculoskelet Rehabil.* 2013;26(3):273–279. doi:10.3233/BMR-130376.
97. Borowsky CD, Fagen G. Sources of sacroiliac region pain: insights gained from a study comparing standard intra-articular injection with a technique combining intra-and peri-articular injection. *Arch Phys Med Rehabil.* 2008;89(11):2048–2056.
98. Palsson TS, Graven-Nielsen T. Experimental pelvic pain facilitates pain provocation tests and causes regional hyperalgesia. *Pain.* 2012;153(11):2233–2240. doi:10.1016/j.pain.2012.07.013.
99. Simopoulos TT, Manchidanti L, Singh V, et al. A systematic evaluation of prevalence and diagnostic accuracy of sacroiliac joint interventions. *Pain Physician.* 2012;15(3):E305–344.
100. Visser LH, Nijssen PGN, Tijssen CC, Van Middendorp JJ, Schieving J. Sciatica-like symptoms and the sacroiliac joint: Clinical features and differential diagnosis. *Eur Spine J.* 2013;22(7):1657–1664. doi:10.1007/s00586-013-2660-5.
101. Masi AT, Benjamin M, Vleeming A. Anatomical, biomechanical, and clinical perspectives on sacroiliac joints: an integrative synthesis of biodynamic mechanisms related to ankylosing spondylitis. In: Vleeming A, Mooney V, Stoeckart R, eds. *Movement, Stability & Lumbopelvic Pain: Integration of Research and Therapy.* 2nd ed. Edinburgh; New York, NY: Churchill Livingstone Elsevier; 2007.
102. Walker JM. Age-related differences in the human sacroiliac joint: a histological study; - implications for therapy. *J Orthop Sports Phys Ther.* 1986;7(6):325–334.
103. Bowen V, Cassidy JD. Macroscopic and microscop-ic anatomy of the sacroiliac joint from embryonic life until the eighth decade. *Spine (Phila Pa 1976).* 1981;6(6):620–628.
104. Weisl H. The articular surfaces of the sacro-iliac joint and their relation to the movements of the sacrum. *Acta Anat (Basel).* 1954;22(1):1–14.
105. Sashin D. A critical analysis of the anatomy and the pathologic changes of the sacro-iliac joints. *J Bone Joint Surg.* 1930;12:891–910.
106. Weisl H. The movements of the sacro-iliac joint. *Cells Tissues Organs.* 1955;23(1):80–91. doi:10.1159/000140984.
107. McLauchlan G, Gardner D. Sacral and iliac articular cartilage thickness and cellularity: relationship to sub-chondral bone end-plate thickness and cancellous bone density. *Rheumatology.* 2002;41(4):375–380.
108. Williams PL, Bannister LH, Berry MM, et al. *Gray's Anatomy : The Anatomical Basis of Medicine and Surgery.* 38th ed. New York, NY: Churchill Livingstone; 1995.
109. Schwarzer AC, April CN, Bogduk N. The sacroiliac joint in chronic low back pain. *Spine (Phila Pa 1976).* 1995;20(1):31–37.
110. Fortin JD, Washington WJ, Falco FJ. Three pathways between the sacroiliac joint and neural structures. *AJNR Am J Neuroradiol.* 1999;20(8):1429–1434.
111. Bani D. Relaxin: A pleiotropic hormone. *Gen Pharmacol Vasc Syst.* 1997;28(1):13–22. doi:10.1016/S0306-3623(96)00171-1.
112. Aldabe D, Ribeiro DC, Milosavljevic S, Bussey MD. Pregnancy-related pelvic girdle pain and its relationship with relaxin levels during pregnancy: A systematic review. *Eur Spine J.* 2012;21(9):1769–1776. doi:10.1007/s00586-012-2162-x.
113. Vukicević S, Marusić A, Stavljenić A, Vujičić G, Skavić J, Vukicević D. Holographic analysis of the human pelvis. *Spine (Phila Pa 1976).* 1991;16(2):209–214. Ć
114. Chen CS, Ingber DE. Tensegrity and mechanoregulation: from skeleton to cytoskeleton. *Osteoarthritis Cartilage.* 1999;7(1):81–94. doi:10.1053/joca.1998.0164.
115. Ingber DE. Tensegrity-based mechanosensing from macro to micro. *Prog Biophys Mol Biol.* 2008;97(2-3):163–179. doi:10.1016/j.pbiomolbio.2008.02.005.
116. Galli C, Guizzardi S, Passeri G, Macaluso GM, Scandroglio R. Life on the wire: on tensegrity and force balance in cells. *Acta Bio-Medica Atenei Parm.* 2005;76(1):5–12.
117. Fuller B. Tensegrity. *Portf Artnews Annu.* 1961;4:112–127.
118. Barker PJ, Briggs CA, Bogeski G. Tensile transmission across the lumbar fasciae in unembalmed cadavers: effects of tension to various muscular attachments. *Spine (Phila Pa 1976).* 2004;29(2):129–138. doi:10.1097/01.BRS.0000107005.62513.32.
119. Lavignolle B, Vital JM, Senegas J, et al. An approach to the functional anatomy of the sacroiliac joints in vivo. *Anat Clin.* 1983;5(3):169–176.
120. Miller JA, Schultz AB, Andersson GB. Load-displace-ment behavior of sacroiliac joints. *J Orthop Res Off Publ Orthop Res Soc.* 1987;5(1):92–101. doi:10.1002/jor.1100050112.
121. Egund N, Olsson TH, Schmid H, Selvik G. Movements in the sacroiliac joints demonstrated with roentgen stereophotogrammetry. *Acta Radiol Diagn (Stockh).* 1978;19(5):833–846.
122. Sturesson B, Selvik G, Udén A. Movements of the sacro-iliac joints. A roentgen stereophotogrammetric analysis. *Spine (Phila Pa 1976).* 1989;14(2):162–165.

123. Walheim G, Olerud S, Ribbe T. Mobility of the pubic symphysis. Measurements by an electromechanical method. *Acta Orthop Scand*. 1984;55(2):203–208. doi:10.3109/17453678408992338.
124. Walheim GG, Selvik G. Mobility of the pubic symphysis. In vivo measurements with an electromechanic method and a roentgen stereophotogrammetric method. *Clin Orthop Relat Res*. 1984;(191):129–135.
125. Walheim GG, Olerud S, Ribbe T. Motion of the pubic symphysis in pelvic instability. *Scand J Rehabil Med*. 1984;16(4):163–169.
126. Sturesson B, Uden A, Vleeming A. A radiostereometric analysis of movements of the sacroiliac joints during the standing hip flexion test. *Spine (Phila Pa 1976)*. 2000;25(3):364–368.
127. Berezin D. Pelvic insufficiency during pregnancy and after parturition; a clinical study. *Acta Obstet Gynecol Scand Suppl*. 1954;33(3):3–119.
128. Vleeming A, Pool-Goudzwaard AL, Hammudoglu D, Stoeckart R, Snijders CJ, Mens JM. The function of the long dorsal sacroiliac ligament: its implication for understanding low back pain. *Spine (Phila Pa 1976)*. 1996;21(5):556–562.
129. Vleeming A, de Vries HJ, Mens JMA, van Wingerden JP. Possible role of the long dorsal sacroiliac ligament in women with peripartum pelvic pain. *Acta Obstet Gynecol Scand*. 2002;81(5):430–436.
130. van Wingerden JP, Vleeming A, Snijders CJ, Stoeckart R. A functional-anatomical approach to the spine-pelvis mechanism: interaction between the biceps femoris muscle and the sacrotuberous ligament. *Eur Spine J*. 1993;2(3):140–144.
131. Dujardin FH, Roussignol X, Hossenboccus M, Thominne JM. Experimental study of the sacroiliac joint micromotion in pelvic disruption. *J Orthop Trauma*. 2002;16(2):99–103.
132. Vleeming A, Van Wingerden JP, Snijders CJ, Stoeckart R, Stijnen T. Load application to the sacrotuberous ligament; influences on sacroiliac joint mechanics. *Clin Biomech*. 1989;4(4):204–209. doi:10.1016/0268-0033(89)90003-X.
133. Borrelli J, Koval KJ, Helfet DL. Operative stabilization of fracture dislocations of the sacroiliac joint. *Clin Orthop Relat Res*. 1996;(329):141–146.
134. Vleeming A, Stoeckart R, Snijders CJ. The sacrotuberous ligament: a conceptual approach to its dynamic role in stabilizing the sacroiliac joint. *Clin Biomech*. 1989;4(4):201–203. doi:10.1016/0268-0033(89)90002-8.
135. Snijders CJ, Hermans PF., Niesing R, Spoor CW, Stoeckart R. The influence of slouching and lumbar support on iliolumbar ligaments, intervertebral discs and sacroiliac joints. *Clin Biomech*. 2004;19(4):323–329.
136. Vrahlas M, Hern TC, Diangelo D, Kellam J, Tile M. Ligamentous contributions to pelvic stability. *Orthopedics*. 1995;18(3):271–274.
137. McGrath C, Nicholson H, Hurst P. The long posterior sacroiliac ligament: a histological study of morphological relations in the posterior sacroiliac region. *Joint Bone Spine*. 2009;76(1):57–62. doi:10.1016/j.jbspin.2008.02.015.
138. Grob KR, Neuhuber WL, Kissling RO. Innervation of the sacroiliac joint of the human. *Z Rheumatol*. 1995;54(2):117–122.
139. Dreyfuss P, Henning T, Malladi N, Goldstein B, Bogduk N. The ability of multi-site, multi-depth sacral lateral branch blocks to anesthetize the sacroiliac joint complex. *Pain Med*. 2009;10(4):679–688. doi:10.1111/j.1526-4637.2009.00631.x.
140. Ikeda R. Innervation of the sacroiliac joint. Macroscopic and histological studies. *Nippon Ika Daigaku Zasshi*. 1991;58(5):587–596.
141. Vilensky JA, O'Connor BL, Fortin JD, et al. Histologic analysis of neural elements in the human sacroiliac joint. *Spine (Phila Pa 1976)*. 2002;27(11):1202–1207.
142. Szadek KM, Hoogland PVJM, Zuurmond WWA, De Lange JJ, Perez RSGM. Possible nociceptive structures in the sacroiliac joint cartilage: An immunohistochemical study. *Clin Anat*. 2010;23(2):192–198. doi:10.1002/ca.20908.
143. Kurosawa D, Murakami E, Aizawa T. Referred pain location depends on the affected section of the sacroiliac joint. *Eur Spine J*. 2014;24(3):521–527. doi:10.1007/s00586-014-3604-4.
144. Vera-Garcia FJ, Elvira JL., Brown SH., McGill SM. Effects of abdominal stabilization maneuvers on the control of spine motion and stability against sudden trunk perturbations. *J Electromyogr Kinesiol*. 2007;17(5):556–567.
145. Buyruk HM, Stam HJ, Snijders CJ, Laméris JS, Holland WP, Stijnen TH. Measurement of sacroiliac joint stiffness in peripartum pelvic pain patients with Doppler imaging of vibrations (DIV). *Eur J Obstet Gynecol Reprod Biol*. 1999;83(2):159–163.
146. Damen L, Buyruk HM, Güler-Uysal F, Lotgering FK, Snijders CJ, Stam HJ. The prognostic value of asymmetric laxity of the sacroiliac joints in pregnancy-related pelvic pain. *Spine (Phila Pa 1976)*. 2002;27(24):2820–2824. doi:10.1097/01.BRS.0000035726.65066.57.
147. Wnek G. *Encyclopedia of Biomaterials and Biomedical Engineering*. 2nd ed. New York, NY: Informa Healthcare USA; 2008.
148. Bogduk N, Macintosh JE. The applied anatomy of the thoracolumbar fascia. *Spine (Phila Pa 1976)*. 1984;9(2):164–170.
149. Barker PJ, Briggs CA. Attachments of the posterior layer of lumbar fascia. *Spine (Phila Pa 1976)*. 1999;24(17):1757–1764.
150. Barker PJ, Briggs CA. Anatomy and biomechanics of the lumbar fasciae. In: Vleeming A, Mooney V, Stoeckart R, eds. *Movement, Stability & Lumbopelvic Pain: Integration of Research and Therapy*. 2nd ed. Edinburgh; New York, NY: Churchill Livingstone Elsevier; 2007.
151. Mooney V, Pozos R, Vleeming A, Gulick J, Swenski D. Exercise treatment for sacroiliac pain. *Orthopedics*. 2001;24(1):29–32.
152. Bogduk N. A reappraisal of the anatomy of the human lumbar erector spinae. *J Anat*. 1980;131(Pt 3):525–540.
153. McGill S. *Low Back Disorders: Evidence-Based Prevention and Rehabilitation*. 2nd ed. Champaign, IL: Human Kinetics; 2007.
154. Hides JA, Richardson CA, Jull GA. Multifidus muscle recovery is not automatic after resolution of acute, first-episode low back pain. *Spine (Phila Pa 1976)*. 1996;21(23):2763–2769.
155. Parkkola R, Rytökoski U, Kormano M. Magnetic resonance imaging of the discs and trunk muscles in patients with chronic low back pain and healthy control subjects. *Spine (Phila Pa 1976)*. 1993;18(7):830–836.

156. Hides JA, Stokes MJ, Saide M, Jull GA, Cooper DH. Evidence of lumbar multifidus muscle wasting ipsilateral to symptoms in patients with acute/subacute low back pain. *Spine (Phila Pa 1976)*. 1994;19(2):165–172.
157. Paris S, Viti J. Differential Diagnosis of Low Back Pain. In: Vleeming A, Mooney V, Stoeckart R, eds. *Movement, stability & Lumbopelvic Pain: Integration of Research and Therapy*. 2nd ed. Edinburgh; New York, NY: Churchill Livingstone Elsevier; 2007:381–390.
158. Lehto M, Hurme M, Alaranta H, et al. Connective tissue changes of the multifidus muscle in patients with lumbar disc herniation. An immunohistologic study of collagen types I and III and fibronectin. *Spine (Phila Pa 1976)*. 1989;14(3):302–309.
159. Mattila M, Hurme M, Alaranta H, et al. The multifidus muscle in patients with lumbar disc herniation. A histochemical and morphometric analysis of intraoperative biopsies. *Spine (Phila Pa 1976)*. 1986;11(7):732–738.
160. Yoshihara K, Shirai Y, Nakayama Y, Uesaka S. Histochemical changes in the multifidus muscle in patients with lumbar intervertebral disc herniation. *Spine (Phila Pa 1976)*. 2001;26(6):622–626.
161. McGill S. *Ultimate Back Fitness and Performance*. Waterloo, Ontario: Wabuno Publishers; 2004.
162. McGill S, Juher D, Kropf P. Quantitative intramuscular myoelectric activity of quadratus lumborum during a wide variety of tasks. *Clin Biomech (Bristol, Avon)*. 1996;11(3):170–172.
163. Gibbons S. Clinical anatomy and function of psoas major and deep sacral gluteus maximus. In: Vleeming A, Mooney V, Stoeckart R, eds. *Movement, Stability and Lumbopelvic Pain: Integration of Research and Therapy*. 2nd ed. Edinburgh; New York, NY: Churchill Livingstone Elsevier; 2007:95–102.
164. Leinonen V, Kankaanpää M, Airaksinen O, Hänninen O. Back and hip extensor activities during trunk flexion/extension: Effects of low back pain and rehabilitation. *Arch Phys Med Rehabil*. 2000;81(1):32–37. doi:10.1016/S0003-9993(00)90218-1.
165. Nelson-Wong E, Alex B, Csepe D, Lancaster D, Callaghan JP. Altered muscle recruitment during extension from trunk flexion in low back pain developers. *Clin Biomech*. 2012;27(10):994–998. doi:10.1016/j.clinbiomech.2012.07.007.
166. Vogt L, Pfeifer K, Banzer W. Neuromuscular control of walking with chronic low-back pain. *Man Ther*. 2003;8(1):21–28. doi:10.1054/math.2002.0476.
167. Hammer N, Steinke H, Lingslebe U, et al. Ligamentous influence in pelvic load distribution. *Spine J*. 2013;13(10):1321–1330. doi:10.1016/j.spinee.2013.03.050.
168. Hodges PW. Changes in motor planning of feedforward postural responses of the trunk muscles in low back pain. *Exp Brain Res*. 2001;141(2):261–266. doi:10.1007/s002210100873.
169. Dangaria TR, Naesh O. Changes in cross-sectional area of psoas major muscle in unilateral sciatica caused by disc herniation. *Spine (Phila Pa 1976)*. 1998;23(8):928–931.
170. Barker KL, Shamley DR, Jackson D. Changes in the cross-sectional area of multifidus and psoas in patients with unilateral back pain: the relationship to pain and disability. *Spine (Phila Pa 1976)*. 2004;29(22):E515–519.
171. Norton-Old KJ, Schache AG, Barker PJ, Clark RA, Harrison SM, Briggs CA. Anatomical and mechanical relationship between the proximal attachment of adductor longus and the distal rectus sheath. *Clin Anat*. 2013;26(4):522–530. doi:10.1002/ca.22116.
172. Aston B. Postnatal pelvic floor dysfunction: conservative treatment and management options. *J Fam Health Care*. 2010;20(3):90–92.
173. Goode PS, Burgio KL, Richter HE, Markland AD. Incontinence in older women. *J Am Med Assoc*. 2010;303(21):2172–2181. doi:10.1001/jama.2010.749.
174. Heilbrun ME, Nygaard IE, Lockhart ME, et al. Correlation between levator ani muscle injuries on magnetic resonance imaging and fecal incontinence, pelvic organ prolapse, and urinary incontinence in primiparous women. *Am J Obstet Gynecol*. 2010;202(5):488.e1–6. doi:10.1016/j.ajog.2010.01.002.
175. Sjödahl J. Pregnancy-related pelvic girdle pain and its relation to muscle function. *Br Med J*. 2010;(1188). doi:10.1136/bmj.38397.507014.E0.
176. Stuge B, Sætre K, Brækken IH. The association between pelvic floor muscle function and pelvic girdle pain - A matched case control 3D ultrasound study. *Man Ther*. 2012;17(2):150–156. doi:10.1016/j.math.2011.12.004.
177. Fitzgerald CM, Santos LR, Mallinson T. The association between pelvic girdle pain and urinary incontinence among pregnant women in the second trimester. *Int J Gynecol Obstet*. 2012;117(3):248–250. doi:10.1016/j.ijgo.2012.01.014.
178. O'Sullivan PB, Beales DJ, Beetham JA, et al. Altered motor control strategies in subjects with sacroiliac joint pain during the active straight-leg-raise test. *Spine (Phila Pa 1976)*. 2002;27(1):E1.
179. O'Sullivan PB, Beales DJ. Changes in pelvic floor and diaphragm kinematics and respiratory patterns in subjects with sacroiliac joint pain following a motor learning intervention: a case series. *Man Ther*. 2007;12(3):209–218. doi:10.1016/j.math.2006.06.006.
180. Hay-Smith J, Mørkved S, Fairbrother KA, Herbison GP. Pelvic floor muscle training for prevention and treatment of urinary and faecal incontinence in antenatal and postnatal women. *Cochrane Database Syst Rev*. 2008;(4):CD007471. doi:10.1002/14651858.CD007471.
181. Bø K. Pelvic floor muscle training is effective in treatment of female stress urinary incontinence, but how does it work? *Int Urogynecol J Pelvic Floor Dysfunct*. 2004;15(2):76–84. doi:10.1007/s00192-004-1125-0.
182. Bø K. Pelvic floor muscle training in treatment of female stress urinary incontinence, pelvic organ prolapse and sexual dysfunction. *World J Urol*. 2012;30(4):437–443. doi:10.1007/s00345-011-0779-8.
183. Sapsford R. Rehabilitation of pelvic floor muscles utilizing trunk stabilization. *Man Ther*. 2004;9(1):3–12. doi:10.1016/S1356-689X(03)00131-0.
184. Hung HC, Hsiao SM, Chih SY, Lin HH, Tsauo JY. An alternative intervention for urinary incontinence: Retraining diaphragmatic, deep abdominal and pelvic floor muscle coordinated function. *Man Ther*. 2010;15(3):273–279. doi:10.1016/j.math.2010.01.008.
185. Schamberger W. *The Malalignment Syndrome : Implications for Medicine and Sport*. Edinburgh; London, UK: Churchill Livingstone; 2002.

186. Bussey MD, Yanai T, Milburn P. A non-invasive technique for assessing innominate bone motion. *Clin Biomech (Bristol, Avon)*. 2004;19(1):85–90.
187. Stevens A. Side-bending and axial rotation of the sacrum inside the pelvic girdle. In: *First Interdisciplinary World Congress on Low Back Pain and its Relation to the Sacroiliac Joint*. San Diego, CA; 1992.
188. Mitchell F. The balanced pelvis and its relationship to reflexes. *AAO Yearbook*. 1948:146–151.
189. Mens JM, Vleeming A, Snijders CJ, Stam HJ, Ginai AZ. The active straight leg raising test and mobility of the pelvic joints. *Eur Spine J*. 1999;8(6):468–473.
190. Damen L, Spoor CW, Snijders CJ, Stam HJ. Does a pelvic belt influence sacroiliac joint laxity? *Clin Biomech (Bristol, Avon)*. 2002;17(7):495–498.
191. Richardson CA, Snijders CJ, Hides JA, Damen L, Pas MS, Storm J. The relation between the transversus abdominis muscles, sacroiliac joint mechanics, and low back pain. *Spine (Phila Pa 1976)*. 2002;27(4):399–405.
192. Hungerford B, Gillett W, Lee D. Altered patterns of pelvic bone motion determined in subjects with posterior pelvic pain using skin markers. *Clin Biomech (Bristol, Avon)*. 2004;19(5):456–464. doi:10.1016/j.clinbiomech.2004.02.004.
193. Beales DJ, O'Sullivan PB, Briffa NK. The effect of increased physical load during an active straight leg raise in pain free subjects. *J Electromyogr Kinesiol*. 2010;20(4):710–718. doi:10.1016/j.jelekin.2009.12.004.
194. Wu WH, Meijer OG, Bruijn SM, et al. Gait in pregnancy-related pelvic girdle pain: amplitudes, timing, and coordination of horizontal trunk rotations. *Eur Spine J*. 2008;17(9):1160–1169. doi:10.1007/s00586-008-0703-0.
195. Roussel N, Nijs J, Truijen S, Vervecken L, Mottram S, Stassijns G. Altered breathing patterns during lumbopelvic motor control tests in chronic low back pain: a case-control study. *Eur Spine J*. 2009;18(7):1066–1073. doi:10.1007/s00586-009-1020-y.
196. Berg G, Hammar M, Möller-Nielsen J, Lindén U, Thorblad J. Low back pain during pregnancy. *Obstet Gynecol*. 1988;71(1):71–75.
197. Verma E, Mittal R, Weeks A. Pelvic girdle pain and low back pain in pregnancy: a review. *Pain Pract*. 2010;10(1):60–71. doi:10.1111/j.1533-2500.2009.00327.x.
198. Röst CC, Jacqueline J, Kaiser A, Verhagen AP, Koes BW. Pelvic pain during pregnancy: a descriptive study of signs and symptoms of 870 patients in primary care. *Spine (Phila Pa 1976)*. 2004;29(22):2567–2572.
199. Gutke A, Ostgaard HC, Oberg B. Predicting persistent pregnancy-related low back pain. *Spine (Phila Pa 1976)*. 2008;33(12):E386–393. doi:10.1097/BRS.0b013e31817331a4.
200. Wu WH, Meijer OG, Uegaki K, et al. Pregnancy-related pelvic girdle pain (PPP), I: Terminology, clinical presentation, and prevalence. *Eur Spine J*. 2004;13(7):575–589. doi:10.1007/s00586-003-0615-y.
201. Porterfield JA, DeRosa C. Conditions of weight bearing: asymmetrical overload syndrome (AOS). In: Movement, stability & lumbopelvic pain : integration of research and therapy. In: Vleeming A, Mooney V, Stoeckart R, eds. *Movement, Stability and Lumbopelvic Pain: Integration of Research and Therapy*. 2nd ed. Edinburgh; New York, NY: Churchill Livingstone Elsevier; 2007:391–403.
202. Eichenseer PH, Sybert DR, Cotton JR. A finite element analysis of sacroiliac joint ligaments in response to different loading conditions. *Spine (Phila Pa 1976)*. 2011;36(22):E1446–E1452. doi:10.1097/BRS.0b013e31820bc705.
203. Jackson R. *Differential Diagnosis of Pubic Symphysis Hypermobility*. In: *4th Interdisciplinary World Congress on Low Back & Pelvic Pain: "Moving From Structure to Function," November 8-10, 2001, Montreal, Canada*. (Vleeming A, Mooney V, Gracovetsky S, et al., eds.). San Diego, CA: University of California San Diego Office of Continuing Medical Education; 2001.
204. Chamberlin W. The x-ray examination of the sacroiliac joint. *Delaware Med J*. 1932;4(9).
205. Vora AJ, Doerr KD, Wolfer LR. Functional anatomy and pathophysiology of axial low back pain: disc, posterior elements, sacroiliac joint, and associated pain generators. *Phys Med Rehabil Clin N Am*. 2010;21(4):679–709. doi:10.1016/j.pmr.2010.07.005.
206. Robinson JP, Apriarian AV. Low Back Pain. In: Mayer E, Bushnell M, eds. *Functional Pain Syndromes, Presentation and Pathophysiology*. 1st ed. International Assn for the Study of Pain; 2009:23–53.
207. Wand BM, Parkitny L, O'Connell NE, et al. Cortical changes in chronic low back pain: current state of the art and implications for clinical practice. *Man Ther*. 2011;16(1):15–20. doi:10.1016/j.math.2010.06.008.
208. Apkarian AV, Baliki MN, Geha PY. Towards a theory of chronic pain. *Prog Neurobiol*. 2009;87(2):81–97. doi:10.1016/j.pneurobio.2008.09.018.
209. Palsson TS, Beales D, Slater H, O'Sullivan P, Graven-Nielsen T. Pregnancy is characterized by widespread deep-tissue hypersensitivity independent of lumbopelvic pain intensity, a facilitated response to manual orthopedic tests, and poorer self-reported health. *J Pain*. 2015;16(3):270–282. doi:10.1016/j.jpain.2014.12.002.
210. Mens JMA, Vleeming A, Snijders CJ, Ronchetti I, Stam HJ. Reliability and validity of hip adduction strength to measure disease severity in posterior pelvic pain since pregnancy. *Spine (Phila Pa 1976)*. 2002;27(15):1674–1679.
211. Hungerford BA, Gillett W, Moran M, Emmerson C. Evaluation of the ability of physical therapists to palpate intrapelvic motion with the Stork test on the support side. *Phys Ther*. 2007;87(7):879–887. doi:10.2522/pjt.20060014.
212. Laslett M, Young SB, Aprill CN, McDonald B. Diagnosing painful sacroiliac joints: A validity study of a McKenzie evaluation and sacroiliac provocation tests. *Aust J Physiother*. 2003;49(2):89–97.
213. Young S, Aprill C, Laslett M. Correlation of clinical examination characteristics with three sources of chronic low back pain. *Spine J*. 2003;3(6):460–465.
214. Murakami E, Aizawa T, Noguchi K, Kanno H, Okuno H, Uozumi H. Diagram specific to sacroiliac joint pain site indicated by one-finger test. *J Orthop Sci*. 2008;13(6):492–497. doi:10.1007/s00776-008-1280-0.
215. Laslett M, Aprill CN, McDonald B, Young SB. Diagnosis of sacroiliac joint pain: validity of individual provocation tests and composites of tests. *Man Ther*. 2005;10(3):207–218.

216. van der Wurff P, Buijs EJ, Groen GJ. A multitest regimen of pain provocation tests as an aid to reduce unnecessary minimally invasive sacroiliac joint procedures. *Arch Phys Med Rehabil.* 2006;87(1):10–14. doi:10.1016/j.apmr.2005.09.023.
217. Kokmeyer DJ, Van der Wurff P, Aufdemkampe G, Fickenscher TCM. The reliability of multitest regimens with sacroiliac pain provocation tests. *J Manipulative Physiol Ther.* 2002;25(1):42–48.
218. Adhia DB, Tumilty S, Mani R, Milosavljevic S, Bussey MD. Can hip abduction and external rotation discriminate sacroiliac joint pain? *Man Ther.* 2015;1–7. doi:10.1016/j.math.2015.08.002.
219. Fortin JD, Falco FJ. The Fortin finger test: an indicator of sacroiliac pain. *Am J Orthop (Belle Mead NJ).* 1997;26(7):477–480.
220. Torstensson T, Butler S, Lindgren A, Peterson M, Eriksson M, Kristiansson P. Referred pain patterns provoked on intra-pelvic structures among women with and without chronic pelvic pain: a descriptive study. *PLoS One.* 2015;10(3):e0119542. doi:10.1371/journal.pone.0119542.
221. Lesher JM, Dreyfuss P, Hager N, Kaplan M, Furman M. Hip joint pain referral patterns: a descriptive study. *Pain Med.* 2008;9(1):22–25. doi:10.1111/j.1526-4637.2006.00153.x.
222. Fortin JD, Dwyer AP, West S, Pier J. Sacroiliac joint: pain referral maps upon applying a new injection/arthrography technique. Part I: Asymptomatic volunteers. *Spine (Phila Pa 1976).* 1994;19(13):1475–1482.
223. Simpson R, Gemmell H. Accuracy of spinal orthopaedic tests: a systematic review. *Chiropr Osteopat.* 2006;14(31):26.
224. Fortin JD, Aprill CN, Ponthieux B, Pier J. Sacroiliac joint: pain referral maps upon applying a new injection/arthrography technique. Part II: Clinical evaluation. *Spine (Phila Pa 1976).* 1994;19(13):1483–1489.
225. Dreyfuss P, Snyder BD, Park K, Willard F, Carreiro J, Bogduk N. The ability of single site, single depth sacral lateral branch blocks to anesthetize the sacroiliac joint complex. *Pain Med.* 2008;9(7):844–850.
226. Cohen SP. Sacroiliac Joint Pain: A Comprehensive Review of Anatomy, Diagnosis, and Treatment. *Anesth Analg.* 2005;101(5):1440–1453. doi:10.1213/01.ANE.0000180831.60169.EA.
227. Cibulka MT, Koldehoff R. Clinical usefulness of a cluster of sacroiliac joint tests in patients with and without low back pain. *J Orthop Sports Phys Ther.* 1999;29(2):83–89; discussion 90–92.
228. Sizer PS, McGaillard MK, Azebedo E, et al. The Pelvis and Sacroiliac Joint: Physical Therapy Patient Management Utilizing Current Evidence. In: *Current Concepts of Orthopaedic Physical Therapy*, 3rd Ed. Vol 21.2.2. Current Concepts of Orthopedic Physical Therapy. La Crosse, WI: APTA Orthopaedic Section; 2011.
229. Delitto A, Cibulka MT, Erhard RE, Bowling RW, Tenhula JA. Evidence for use of an extension-mobilization category in acute low back syndrome: a prescriptive validation pilot study. *Phys Ther.* 1993;73(4):216–222; discussion 223–228.
230. Cibulka MT, Delitto A, Koldehoff RM. Changes in innominate tilt after manipulation of the sacroiliac joint in patients with low back pain. An experimental study. *Phys Ther.* 1988;68(9):1359–1363.
231. Tong HC, Heyman OG, Lado DA, Isser MM. Interexaminer reliability of three methods of combining test results to determine side of sacral restriction, sacral base position, and innominate bone position. *J Am Osteopath Assoc.* 2006;106(8):464–468.
232. McGrath MC. Palpation of the sacroiliac joint: An anatomical and sensory challenge. *Int J Osteopath Med.* 2006;9(3):103–107.
233. Beal MC. The sacroiliac problem: review of anatomy, mechanics, and diagnosis. *J Am Osteopath Assoc.* 1982;81(10):667–679.
234. Palsson TS, Hirata RP, Graven-Nielsen T. Experimental pelvic pain impairs the performance during the active straight leg raise test and causes excessive muscle stabilization. *Clin J Pain.* 2015;31(7):642–651 doi:10.1097/AJP.0000000000000139.
235. Broadhurst NA, Bond MJ. Pain provocation tests for the assessment of sacroiliac joint dysfunction. *J Spinal Disord.* 1998;11(4):341–345. doi:10.1097/000002517-199908000-00014.
236. Cook C, Massa L, Harm-Ernandes I, et al. Interrater reliability and diagnostic accuracy of pelvic girdle pain classification. *J Manipulative Physiol Ther.* 2007;30(4):252–258. doi:10.1016/j.jmpt.2007.03.008.
237. Ostgaard HC, Zetherström G, Roos-Hansson E. Back pain in relation to pregnancy: a 6-year follow-up. *Spine (Phila Pa 1976).* 1997;22(24):2945–2950.
238. MacLennan AH, MacLennan SC. Symptom-giving pelvic girdle relaxation of pregnancy, postnatal pelvic joint syndrome and developmental dysplasia of the hip. The Norwegian Association for Women with Pelvic Girdle Relaxation (Landforeningen for Kvinner Med Bekkenløsningsplager). *Acta Obstet Gynecol Scand.* 1997;76(8):760–764.
239. Nilsson-Wikmar L, Holm K, Oijerstedt R, Harms-Ringdahl K. Effect of three different physical therapy treatments on pain and activity in pregnant women with pelvic girdle pain: a randomized clinical trial with 3, 6, and 12 months follow-up postpartum. *Spine (Phila Pa 1976).* 2005;30(8):850–856.
240. Haugland KS, Rasmussen S, Daltveit AK. Group intervention for women with pelvic girdle pain in pregnancy. A randomized controlled trial. *Acta Obstet Gynecol Scand.* 2006;85(11):1320–1326. doi:10.1080/00016340600780458.
241. Elden H, Hagberg H, Olsen MF, Ladfors L, Ostgaard HC. Regression of pelvic girdle pain after delivery: follow-up of a randomised single blind controlled trial with different treatment modalities. *Acta Obstet Gynecol Scand.* 2008;87(2):201–208. doi:10.1080/00016340701823959.
242. Vøllestad NK, Stuge B. Prognostic factors for recovery from postpartum pelvic girdle pain. *Eur Spine J.* 2009;18(5):718–726. doi:10.1007/s00586-009-0911-2.
243. Jackson R. Documenting innominate superior shear dysfunction and efficacy of limited treatment on pain and disability. European Conference Organizers. Rotterdam; 1998.
244. Shultz SJ, Nguyen A-D. Bilateral asymmetries in clinical measures of lower-extremity anatomic characteristics. *Clin J Sport Med.* 2007;17(5):357–361. doi:10.1097/JSM.0b013e31811df950.

245. Golightly YM, Tate JJ, Burns CB, Gross MT. Changes in pain and disability secondary to shoe lift intervention in subjects with limb length inequality and chronic low back pain: a preliminary report. *J Orthop Sports Phys Ther.* 2007;37(7):380-388.
246. Juhl JH, Ippolito Cremin TM, Russell G. Prevalence of frontal plane pelvic postural asymmetry--part 1. *J Am Osteopath Assoc.* 2004;104(10):411–421.
247. Irvin RE. Why and how to optimize posture. In: Vleeming A, Mooney V, Stoeckart R, eds. *Movement, Stability and Lumbopelvic Pain: Integration of Research and Therapy.* 2nd ed. New York, NY: Churchill Livingstone; 2007:239–252.
248. Friberg O. Clinical symptoms and biomechanics of lumbar spine and hip joint in leg length inequality. *Spine (Phila Pa 1976).* 1983;8(6):643–651.
249. Irvin RE. Reduction of lumbar scoliosis by use of a heel lift to level the sacral base. *J Am Osteopath Assoc.* 1991;91(1):34, 37–44.
250. Schultz IZ, Crook JM, Berkowitz J, et al. Biopsychosocial multivariate predictive model of occupational low back disability. *Spine (Phila Pa 1976).* 2002;27(23):2720–2725. doi:10.1097/01.BRS.0000035323.16390.B5.
251. Sizer Jr PS, Brismee JM, Cook C. Medical screening for red flags in the diagnosis and management of musculoskeletal spine pain. *Pain Pract.* 2007;7(1):53-71.
252. Leerar PJ, Boissonnault W, Domholdt E, Roddey T. Documentation of red flags by physical therapists for patients with low back pain. *J Man Manip Ther.* 2007;15(1):42–49.
253. Van den Bosch M, Hollingworth W, Kinmonth AL, Dixon AK. Evidence against the use of lumbar spine radiography for low back pain. *Clin Radiol.* 2004;59(1):69–76.
254. Deyo RA, Diehl AK. Cancer as a cause of back pain: frequency, clinical presentation, and diagnostic strategies. *J Gen Intern Med.* 1988;3(3):230–238.
255. Henschke N, Maher CG, Refshauge KM. Screening for malignancy in low back pain patients: a systematic review. *Eur Spine J.* 2007;16(10):1673–1679. doi:10.1007/s00586-007-0412-0.
256. Ebell MH, Culp MB, Radke TJ. A Systematic Review of Symptoms for the Diagnosis of Ovarian Cancer. *Am J Prev Med.* 2016;50(3):384-394. doi:10.1016/j.amepre.2015.09.023.
257. Fraser S, Roberts L, Murphy E. Cauda equina syndrome: a literature review of its definition and clinical presentation. *Arch Phys Med Rehabil.* 2009;90(11):1964–1968. doi:10.1016/j.apmr.2009.03.021.
258. Henschke N, Maher C, Refshauge K. A systematic review identifies five “red flags” to screen for vertebral fracture in patients with low back pain. *J Clin Epidemiol.* 2008;61(2):110–118.e1. doi:10.1016/j.jclinepi.2007.04.013.
259. Williams CM, Henschke N, Maher CG, et al. Red flags to screen for vertebral fracture in patients presenting with low-back pain. *Cochrane Database Syst Rev.* 2013;(1):CD008643 doi:10.1002/14651858.CD008643.pub2.
260. Ham SJ, Van Walsum ADP, Vierhout PAM. Predictive value of the hip flexion test for fractures of the pelvis. *Injury.* 1996;27(8):543–544.
261. Duane TM, Dechert T, Wolfe LG, et al. Clinical examination is superior to plain films to diagnose pelvic fractures compared to CT. *Am Surg.* 2008;74(6):476–480.
262. Hilibrand MJ, Hammoud S, Bishop M, Woods D, Fredrick RW, Dodson CC. Common injuries and ailments of the female athlete; pathophysiology, treatment and prevention. *Phys Sportsmed.* 2015;43(4):403-411. doi:10.1080/00913847.2015.1092856.
263. Egol KA, Koval KJ, Kummer F, Frankel VH. Stress fractures of the femoral neck. *Clin Orthop Relat Res.* 1998;(348):72–78.
264. Wall J, Feller JF. Imaging of stress fractures in runners. *Clin Sport Med.* 2006;25(4):781–802.
265. Ivkovic A, Bojanic I, Pecina M. Stress fractures of the femoral shaft in athletes: a new treatment algorithm. *Br J Sports Med.* 2006;40(6):518–520, discussion 520. doi:10.1136/bjsm.2005.023655.
266. Pantano KJ. Strategies used by physical therapists in the US for treatment and prevention of the female athlete triad. *Phys Ther Sport.* 2009;10(1):3–11.
267. Ford LS, Ellis AM, Allen HW, Campbell DE. Osteomyelitis and pyogenic sacroiliitis: A difficult diagnosis. *J Paediatr Child Health.* 2004;40(5-6):317–319.
268. Laslett M. Bilateral buttock pain caused by aortic stenosis: a case report of claudication of the buttock. *Man Ther.* 2000;5(4):227–233. doi:10.1054/math.2000.0368.
269. Gray JC. Diagnosis of intermittent vascular claudication in a patient with a diagnosis of sciatica. *Phys Ther.* 1999;79(6):582-590.
270. Chen SJ, Pipinos I, Johanning J, et al. Bilateral claudication results in alterations in the gait biomechanics at the hip and ankle joints. *J Biomech.* 2008;41(11):2506–2514.
271. Laslett M, Öberg B, Aprill CN, McDonald B. Centralization as a predictor of provocation discography results in chronic low back pain, and the influence of disability and distress on diagnostic power. *Spine J.* 2005;5(4):370–380.
272. Berthelot JM, Delecrin J, Maugars Y, Passuti N. Contribution of centralization phenomenon to the diagnosis, prognosis, and treatment of diskogenic low back pain. *Joint Bone Spine.* 2007;74(4):319–323.
273. Chicago BR, Center DC, Baltimore MD, et al. Systematic review of lumbar discography as a diagnostic test for chronic low back pain. *Pain Physician.* 2009;12(3):541–559.
274. Carragee EJ, Don AS, Hurwitz EL, et al. 2009 ISSLS Prize Winner: Does discography cause accelerated progression of degeneration changes in the lumbar disc: a ten-year matched cohort study. *Spine (Phila Pa 1976).* 2009;34(21):2338–2345. doi:10.1097/BRS.0b013e-3181ab5432.
275. Broetz D, Burkard S, Weller M. A prospective study of mechanical physiotherapy for lumbar disk prolapse: Five year follow-up and final report. *NeuroRehabilitation.* 2010;26(2):155–158.
276. Long A, Donelson R, Fung T. Does it matter which exercise? A randomized control trial of exercise for low back pain. *Spine (Phila Pa 1976).* 2004;29(23):2593–2602.
277. Laslett M, Öberg B, Aprill CN, McDonald B. Zygapophyseal joint blocks in chronic low back pain: a test of Revel’s model as a screening test. *BMC Musculoskeletal Disord.* 2004;5(1):43.
278. Staal JB, de Bie RA, de Vet HCW, Hildebrandt J, Nelemans P. Injection therapy for subacute and chronic low back pain: an updated Cochrane review. *Spine*

- (*Phila Pa* 1976). 2009;34(1):49–59. doi:10.1097/BRS.0b013e3181909558.
279. Laslett M, McDonald B, Aprill CN, Tropp H, Oberg B. Clinical predictors of screening lumbar zygapophyseal joint blocks: development of clinical prediction rules. *Spine J.* 2006;6(4):370–379. doi:10.1016/j.spinee.2006.01.004.
280. Revel M, Poiradeau S, Auleley GR, et al. Capacity of the clinical picture to characterize low back pain relieved by facet joint anesthesia. Proposed criteria to identify patients with painful facet joints. *Spine (Phila Pa 1976)*. 1998;23(18):1972–1976; discussion 1977.
281. Manchikanti L, Pampati V, Fellows B, Baha AG. The inability of the clinical picture to characterize pain from facet joints. *Pain Physician*. 2000;3(2):158–166.
282. Hall AM, Maher CG, Latimer J, Ferreira ML, Costa LO. The patient-specific functional scale is more responsive than the Roland Morris disability questionnaire when activity limitation is low. *Eur Spine J.* 2011;20(1):79–86. doi:10.1007/s00586-010-1521-8.
283. Albert H, Godsken M, Westergaard J. Evaluation of clinical tests used in classification procedures in pregnancy-related pelvic joint pain. *Eur Spine J.* 2000;9(2):161–166.
284. Lequesne M, Mathieu P, Vuillemin-Bodaghi V, Bard H, Djian P. Gluteal tendinopathy in refractory greater trochanter pain syndrome: Diagnostic value of two clinical tests. *Arthritis Rheum*. 2008;59(2):241–246. doi:10.1002/art.23354.
285. Browder DA, Erhard RE. Decision making for a painful hip: a case requiring referral. *J Orthop Sports Phys Ther*. 2005;35(11):738–744.
286. Bird PA, Oakley SP, Shnier R, Kirkham BW. Prospective evaluation of magnetic resonance imaging and physical examination findings in patients with greater trochanteric pain syndrome. *Arthritis Rheum*. 2001;44(9):2138–2145.
287. Greenwood MJ, Erhard RE, Jones DL. Differential diagnosis of the hip vs. lumbar spine: five case reports. *J Orthop Sports Phys Ther*. 1998;27(4):308–315.
288. VanWye WR. Patient Screening by a Physical Therapist for Nonmusculoskeletal Hip Pain. *Phys Ther*. 2009;89(3):248–256. doi: 10.2522/ptj.20070366. Epub 2009 Jan 23.
289. Narvani AA, Tsiridis E, Kendall S, Chaudhuri R, Thomas P. A preliminary report on prevalence of acetabular labrum tears in sports patients with groin pain. *Knee Surg Sport Traumatol Arthrosc*. 2003;11(6):403–408. doi:10.1007/s00167-003-0390-7.
290. Philippon MJ, Maxwell RB, Johnston TL, Schenker M, Briggs KK. Clinical presentation of femoroacetabular impingement. *Knee Surgery Sport Traumatol Arthrosc*. 2007;15(8):1041–1047. doi:10.1007/s00167-007-0348-2.
291. Macfarlane RJ, Haddad FS. The diagnosis and management of femoro-acetabular impingement. *Ann R Coll Surg Engl*. 2010;92(5):363–367. doi:10.1308/003588410X12699663903791.
292. Birrell F. Predicting radiographic hip osteoarthritis from range of movement. — *Rheumatology (Oxford)*. 2001;40(5):506–512. doi:10.1093/rheumatology/40.5.506.
293. Cook C. *Orthopedic Physical Examination Tests : An Evidence-Based Approach*. 2nd ed. Upper Saddle River, NJ: PearsonPrentice Hall; 2012.
294. Waddell G. 1987 Volvo award in clinical sciences. A new clinical model for the treatment of low-back pain. *Spine (Phila Pa 1976)*. 1987;12(7):632–644.
295. Indahl A, Velund L, Reikeraas O. Good prognosis for low back pain when left untampered. A randomized clinical trial. *Spine (Phila Pa 1976)*. 1995;20(4):473–477.
296. Bergquist-Ullman M, Larsson U. Acute low back pain in industry. A controlled prospective study with special reference to therapy and confounding factors. *Acta Orthop Scand*. 1977;170(1):1–117.
297. Troup JD, Martin JW, Lloyd DC. Back pain in industry. A prospective survey. *Spine (Phila Pa 1976)*. 1981;6(1):61–69.
298. Hides JA, Jull GA, Richardson CA. Long-term effects of specific stabilizing exercises for first-episode low back pain. *Spine (Phila Pa 1976)*. 2001;26(11):E243–248.
299. Croft P, Macfarlane G, Papageorgiou A, Thomas E, Silman A. Outcome of low back pain in general practice: a prospective study. *BMJ*. 1998;316:1356–1359.
300. Ayala F, de Baranda Andújar PS. Effect of 3 different active stretch durations on hip flexion range of motion. *J Strength Cond Res*. 2010;24(2):430–436. doi:10.1519/JSC.0b013e3181c0674f.
301. Stevens VK, Coorevits PL, Bouche KG, Mahieu NN, Vanderstraeten GG, Danneels LA. The influence of specific training on trunk muscle recruitment patterns in healthy subjects during stabilization exercises. *Man Ther*. 2007;12(3):271–279.
302. Stuge B, Laerum E, Kirkesola G, Vollestad N. The efficacy of a treatment program focusing on specific stabilizing exercises for pelvic girdle pain after pregnancy: a randomized controlled trial. *Spine (Phila Pa 1976)*. 2004;29(4):351.
303. Danneels LA, Vanderstraeten GG, Cambier DC, et al. Effects of three different training modalities on the cross sectional area of the lumbar multifidus muscle in patients with chronic low back pain. *Br J Sports Med*. 2001;35(3):186–191.
304. Stevens VK, Vleeming A, Bouche KG, Mahieu NN, Vanderstraeten GG, Danneels LA. Electromyographic activity of trunk and hip muscles during stabilization exercises in four-point kneeling in healthy volunteers. *Eur Spine J*. 2007;16(5):711–718. doi:10.1007/s00586-006-0181-1.
305. O’Sullivan PB, Phyty GD, Twomey LT, Allison GT. Evaluation of specific stabilizing exercise in the treatment of chronic low back pain with radiologic diagnosis of spondylolysis or spondylolisthesis. *Spine (Phila Pa 1976)*. 1997;22(24):2959–2967.
306. Ferreira PH, Ferreira ML, Maher CG, Herbert RD, Refshauge K. Specific stabilisation exercise for spinal and pelvic pain: A systematic review. *Aust J Physiother*. 2006;52(2):79–88.
307. Teyhen DS, Bluemle LN, Dolbeer JA, et al. Changes in lateral abdominal muscle thickness during the abdominal drawing-in maneuver in those with lumbopelvic pain. *J Orthop Sports Phys Ther*. 2009;39(11):791–798. doi:10.2519/jospt.2009.3128.
308. Sapsford RR, Hodges PW. Contraction of the pelvic floor muscles during abdominal maneuvers. *Arch Phys*

- Med Rehabil.* 2001;82(8):1081–1088. doi:10.1053/apmr.2001.24297.
309. van Tulder M, Malmivaara A, Esmail R, Koes B. Exercise therapy for low back pain: a systematic review within the framework of the cochrane collaboration back review group. *Spine (Phila Pa 1976)*. 2000;25(21):2784.
310. Pillastrini P, Ferrari S, Rattin S, Cupello A, Villafaña JH, Vanti C. Exercise and tropism of the multifidus muscle in low back pain: a short review. *J Phys Ther Sci.* 2015;27(3):943–945. doi:10.1589/jpts.27.943.
311. Vera-Garcia FJ, Moreside JM, McGill SM. Abdominal muscle activation changes if the purpose is to control pelvis motion or thorax motion. *J Electromyogr Kinesiol.* 2011;21(6):893–903. doi:10.1016/j.jelekin.2011.08.003.
312. Son J-H, Park GD, Park HS, Article O. The effect of sacroiliac joint mobilization on pelvic deformation and the static balance ability of female university students with si joint dysfunction. *J Phys Ther Sci.* 2014;26(6):845–848. doi:10.1589/jpts.26.845.
313. Visser LH, Woudenberg NP, De Bont J, et al. Treatment of the sacroiliac joint in patients with leg pain: A randomized-controlled trial. *Eur Spine J.* 2013;22(10):2310–2317. doi:10.1007/s00586-013-2833-2.
314. Kamali F, Shokri E. The effect of two manipulative therapy techniques and their outcome in patients with sacroiliac joint syndrome. *J Bodyw Mov Ther.* 2012;16(1):29–35. doi:10.1016/j.jbmt.2011.02.002.
315. van Benten E, Pool J, Mens J, Pool-Goudzwaard A. Recommendations for physical therapists on the treatment of lumbopelvic pain during pregnancy: a systematic review. *J Orthop Sport Phys Ther.* 2014;44(7):464–473, A1–15. doi:10.2519/jospt.2014.5098.
316. Pennick V, Liddle SD. Interventions for preventing and treating pelvic and back pain in pregnancy. *Cochrane Database Syst Rev.* 2013;8(2):CD001139. doi:10.1002/14651858.CD001139.pub3.
317. Janssens L, McConnell AK, Pijnenburg M, et al. Inspiratory muscle training affects proprioceptive use and low back pain. *Med Sci Sports Exerc.* 2014;(19):12–19. doi:10.1249/MSS.0000000000000385.
318. Vibe Fersum K, O'Sullivan P, Skouen JS, et al. Efficacy of classification-based cognitive functional therapy in patients with non-specific chronic low back pain: A randomized controlled trial. *Eur J Pain.* 2013;17(6):916–928. doi:10.1002/j.1532-2149.2012.00252.x.
319. Shearar KA, Colloca CJ, White HL. A randomized clinical trial of manual versus mechanical force manipulation in the treatment of sacroiliac joint syndrome. *J Manipulative Physiol Ther.* 2005;28(7):493–501.
320. Marshall P, Murphy B. The effect of sacroiliac joint manipulation on feed-forward activation times of the deep abdominal musculature. *J Manipulative Physiol Ther.* 2006;29(3):196–202.
321. McGill S. Evidence Based stabilization and rehabilitation of the low back patient. A Summary by Richard Jackson from “Low Back Disorders” and “Ultimate Back Fitness.” *Jackson Clin Contin Educ Progr.* 2010.
322. Pel JJ, Spoor CW, Goossens RHM, Pool-Goudzwaard AL. Biomechanical model study of pelvic belt influence on muscle and ligament forces. *J Biomech.* 2008;41(9):1878–1884.
323. Vleeming A, Buyruk HM, Stoeckart R, Karamursel S, Snijders CJ. An integrated therapy for peripartum pelvic instability: a study of the biomechanical effects of pelvic belts. *Am J Obstet Gynecol.* 1992;166(4):1243–1247.
324. Hu H, Meijer OG, van Dieën JH, et al. Muscle activity during the active straight leg raise (ASLR), and the effects of a pelvic belt on the ASLR and on treadmill walking. *J Biomech.* 2010;43(3):532–539. doi:10.1016/j.jbiomech.2009.09.035.
325. Pool-Goudzwaard AL, Vleeming A, Stoeckart R, Snijders CJ, Mens JM. Insufficient lumbopelvic stability: a clinical, anatomical and biomechanical approach to “a-specific” low back pain. *Man Ther.* 1998;3(1):12–20. doi:10.1054/math.1998.0311.
326. Soisson O, Lube J, Germano A, et al. Pelvic belt effects on pelvic morphometry, muscle activity and body balance in patients with sacroiliac joint dysfunction. *PLoS One.* 2015;10(3):e0116739. doi:10.1371/journal.pone.0116739.
327. Arumugam A, Milosavljevic S, Woodley S, Sole G. Effects of external pelvic compression on form closure, force closure, and neuromotor control of the lumbopelvic spine - A systematic review. *Man Ther.* 2012;17(4):275–284. doi:10.1016/j.math.2012.01.010.
328. Sichting F, Rossol J, Soisson O, Klima S, Milani T, Hammer N. Pelvic belt effects on sacroiliac joint ligaments: a computational approach to understand therapeutic effects of pelvic belts. *Pain Physician.* 2014;17(1):43–51.
329. Brizzolara K, Wang-price S, Roddey T. The effect of pelvic compression on deep abdominal muscle thickness during the active straight leg raise test. *Int J Ther Rehabil.* 2015;22(2):60–67. doi: 10.12968/ijtr.2015.22.2.60
330. Simopoulos TT, Manchikanti L, Gupta S, et al. Systematic review of the diagnostic accuracy and therapeutic effectiveness of sacroiliac joint interventions. *Pain Physician.* 2015;18(5):E713–E756.
331. Dreyfuss P, Stout A, April C, Pollei S, Johnson B, Bogduk N. The significance of multifidus atrophy after successful radiofrequency neurotomy for low back pain. *PM R J Inj Funct Rehabil.* 2009;1(8):719–722. doi:10.1016/j.pmrj.2009.05.014.
332. Cohen SP, Strassels SA, Kurihara C, et al. Outcome predictors for sacroiliac joint (lateral branch) radiofrequency denervation. *Reg Anesth Pain Med.* 2009;34(3):206–214. doi:10.1097/AAP.0b013e3181958f4b.
333. Rabago D, Best TM, Zgierska AE, Zeisig E, Ryan M, Crane D. A systematic review of four injection therapies for lateral epicondylitis: prolotherapy, polidocanol, whole blood and platelet-rich plasma. *Br J Sports Med.* 2009;43(7):471–481. doi:10.1136/bjsm.2008.052761.
334. Topol GA, Reeves KD, Hassanein KM. Efficacy of dextrose prolotherapy in elite male kicking-sport athletes with chronic groin pain. *Arch Phys Med Rehabil.* 2005;86(4):697–702. doi:10.1016/j.apmr.2004.10.007.
335. Centeno CJ, Elliott J, Elkins WL, Freeman M. Fluoroscopically guided cervical prolotherapy for instability with blinded pre and post radiographic reading. *Pain Physician.* 2005;8(1):67–72.
336. Chou R, Atlas SJ, Stanos SP, Rosenquist RW. Nonsurgical interventional therapies for low back pain: a review of the evidence for an American Pain Society clinical practice guideline. *Spine (Phila Pa 1976).* 2009;34(10):1078–1093. doi:10.1097/BRS.0b013e-3181a103b1.

337. Rabago D, Slattengren A, Zgierska A. Prolotherapy in primary care practice. *Prim Care*. 2010;37(1):65–80. doi:10.1016/j.pop.2009.09.013.
338. Yelland MJ, Sweeting KR, Lyftogt JA, Ng SK, Scuffham PA, Evans KA. Prolotherapy injections and eccentric loading exercises for painful Achilles tendinosis: a randomised trial. *Br J Sports Med*. 2011;45(5):421–428. doi:10.1136/bjsm.2009.057968.
339. Dagenais S, Yelland MJ, Del Mar C, Schoene ML. Prolotherapy injections for chronic low-back pain. *Cochrane Database Syst Rev*. 2007;(2):CD004059. doi:10.1002/14651858.CD004059.pub3.
340. Cusi M, Saunders J, Hungerford B, Wisbey-Roth T, Lucas P, Wilson S. The use of prolotherapy in the sacro-iliac joint. *Br J Sports Med*. 2008;44(2):100–104. doi:10.1136/bjsm.2007.042044.
341. Kim WM, Lee HG, Jeong CW, Kim CM, Yoon MH. A randomized controlled trial of intra-articular prolotherapy versus steroid injection for sacroiliac joint pain. *J Altern Complement Med*. 2010;16(12):1285–1290. doi:10.1089/acm.2010.0031.
342. Lyras DN, Kazakos K, Verettas D, et al. The influence of platelet-rich plasma on angiogenesis during the early phase of tendon healing. *Foot Ankle Int*. 2009;30(11):1101–1106. doi:10.3113/FAI.2009.1101.
343. Kon E, Buda R, Filardo G, et al. Platelet-rich plasma: intra-articular knee injections produced favorable results on degenerative cartilage lesions. *Knee Surgery Sport Traumatol Arthrosc*. 2010;18(4):472–479. doi:10.1007/s00167-009-0940-8.
344. Knezevic NN, Candido KD, Desai R, Kaye AD. Is platelet-rich plasma a future therapy in pain management? *Med Clin North Am*. 2016;100(1):199–217. doi:10.1016/j.mcna.2015.08.014.
345. Zhang Q, Chen W, Liu H, Su Y, Pan J, Zhang Y. The anterior dislocation of the sacroiliac joint: a report of four cases and review of the literature and treatment algorithm. *Arch Orthop Trauma Surg*. 2009;129(7):941–947. doi:10.1007/s00402-009-0853-5.
346. Bouguennec N, Gouin F, Piétu G. Isolated anterior unilateral sacroiliac dislocation without pubic arch disjunction. *Orthop Traumatol Surg Res*. 2012;98(3):359–362. doi:10.1016/j.otsr.2011.10.013.
347. Baskin KM, Cahill AM, Kaye RD, Born CT, Grudziak JS, Towbin RB. Closed reduction with CT-guided screw fixation for unstable sacroiliac joint fracture-dislocation. *Pediatr Radiol*. 2004;34(12):963–969. doi:10.1007/s00247-004-1291-8.
348. Wise CL, Dall BE. Minimally invasive sacroiliac arthrodesis: outcomes of a new technique. *J Spinal Disord Tech*. 2008;21(8):579–584. doi:10.1097/BSD.0b013e31815ec4b.
349. Khurana A, Guha AR, Mohanty K, Ahuja S. Percutaneous fusion of the sacroiliac joint with hollow modular anchorage screws: clinical and radiological outcome. *J Bone Joint Surg Br*. 2009;91(5):627–631. doi:10.1302/0301-620X.91B5.21519.
350. Heiney J, Capobianco R, Cher D. A systematic review of minimally invasive sacroiliac joint fusion utilizing a lateral transarticular technique. *Int J Spine Surg*. 2015;9. doi:10.14444/2040.
351. Ivanov AA, Kiapour A, Ebraheim NA, Goel V. Lumbar fusion leads to increases in angular motion and stress across sacroiliac joint: a finite element study. *Spine (Phila Pa 1976)*. 2009;34(5):E162–169. doi:10.1097/BRS.0b013e3181978ea3.
352. Ha K-Y, Lee J-S, Kim K-W. Degeneration of sacroiliac joint after instrumented lumbar or lumbosacral fusion: a prospective cohort study over five-year follow-up. *Spine (Phila Pa 1976)*. 2008;33(11):1192–1198. doi:10.1097/BRS.0b013e318170fd35.
353. Pelletier R, Higgins J, Bourbonnais D. Is neuroplasticity in the central nervous system the missing link to our understanding of chronic musculoskeletal disorders? *BMC Musculoskelet Disord*. 2015;16(1):25. doi:10.1186/s12891-015-0480-y.
354. Moseley GL. Widespread brain activity during an abdominal task markedly reduced after pain physiology education: fMRI evaluation of a single patient with chronic low back pain. *Aust J Physiother*. 2005;51:49–52.
355. Moseley GL, Nicholas MK, Hodges PW. A randomized controlled trial of intensive neurophysiology education in chronic low back pain. *Clin J Pain*. 2004;20(5):324–330.
356. van der Windt D, Hay E, Jellema P, Main C. Psychosocial interventions for low back pain in primary care: lessons learned from recent trials. *Spine (Phila Pa 1976)*. 2008;33(1):81.
357. Gutke A, Josefsson A, Öberg B. Pelvic girdle pain and lumbar pain in relation to postpartum depressive symptoms. *Spine (Phila Pa 1976)*. 2007;32(13):1430.
358. Daffada PJ, Walsh N, McCabe CS, Palmer S. The impact of cortical remapping interventions on pain and disability in chronic low back pain: A systematic review. *Physiotherapy*. 2015;101(1):25–33. doi:10.1016/j.physio.2014.07.002.
359. Moseley GL. Evidence for a direct relationship between cognitive and physical change during an education intervention in people with chronic low back pain. *Eur J Pain*. 2004;8(1):39–45. doi:10.1016/S1090-3801(03)00063-6.
360. Bowering KJ, O’Connell NE, Tabor A, et al. The effects of graded motor imagery and its components on chronic pain: a systematic review and meta-analysis. *J Pain*. 2013;14(1):3–13. doi:10.1016/j.jpain.2012.09.007.
361. Wand BM, Connell NE, Di Pietro F, Bulsara M. Back pain with a sensorimotor retraining approach: exploratory multiple-baseline study of 3 participants. *Phys Ther*. 2011;91(4):536–546. doi: 10.2522/ptj.20100150. Epub 2011 Feb 24.
362. Altman R, Alarcón G, Appelrouth D, et al. The American College of Rheumatology criteria for the classification and reporting of osteoarthritis of the hip. *Arthritis Rheum*. 1991;34(5):505–514.
363. Caesar BC, Roberts SJ. Stress Fractures of the Femoral Diaphysis. *Oper Tech Sports Med*. 2009;17(2):94–99. doi:10.1053/j.otsm.2009.05.008.
364. Christodoulides AN. Ipsilateral sciatica on femoral nerve stretch test is pathognomonic of an L4/5 disc protrusion. *J Bone Joint Surg Br*. 1989;71(1):88–89.