

Blood Flow Restricted Exercise: Physical Therapy Patient Management Using Current Evidence

Independent Study
Course 30.3.3

Johnny G. Owens, MPT
Owens Recovery Science
San Antonio, TX, USA

Luke Hughes, PhD
Faculty of Sport, Health and Applied Science
St. Mary's University, London, UK

Stephen D. Patterson, PhD
Faculty of Sport, Health and Applied Science
St. Mary's University, London, UK

CONTINUING PHYSICAL THERAPY EDUCATION

ACADEMY OF
**ORTHOPAEDIC
PHYSICAL THERAPY**



APTA
American Physical Therapy Association

REFERENCES

1. The Nobel Prize in Physiology or Medicine 2019. NobelPrize.org. <https://www.nobelprize.org/prizes/medicine/2019/advanced-information/>. Accessed December 1, 2019.
2. Alam M, Smirk FH. Observations in man upon a blood pressure raising reflex arising from the voluntary muscles. *J Physiol*. 1937;89(4):372-383. doi: 10.1113/jphysiol.1937.sp003485.
3. Collens WS, Wilensky ND. Intermittent venous occlusion in treatment of peripheral vascular disease: an experience with one hundred and twenty-four cases. *JAMA*. 1937;109(26):2125-2130. doi:10.1001/jama.1937.02780520015004.
4. Murry CE, Jennings RB, Reimer KA. Preconditioning with ischemia: a delay of lethal cell injury in ischemic myocardium. *Circulation*. 1986;74(5):1124-1136. doi: 10.1161/01.cir.74.5.1124.
5. Eiken O, Bjurstedt H. Dynamic exercise in man as influenced by experimental restriction of blood flow in the working muscles. *Acta Physiol Scand*. 1987;131(3):339-345. doi: 10.1111/j.1748-1716.1987.tb08248.x.
6. Takarada Y, Nakamura Y, Aruga S, Onda T, Miyazaki S, Ishii N. Rapid increase in plasma growth hormone after low-intensity resistance exercise with vascular occlusion. *J Appl Physiol*. 2000;88(1):61-65. doi: 10.1152/jap-2000.88.1.61.
7. Slys J, Stultz J, Burr JF. The efficacy of blood flow restricted exercise: A systematic review and meta-analysis. *J Sci Med Sport*. 2016;19(8):669-675. doi: 10.1016/j.jsams.2015.09.005.
8. Centner C, Wiegel P, Gollhofer A, König D. Effects of blood flow restriction training on muscular strength and hypertrophy in older individuals: a systematic review and meta-analysis. *Sports Med*. 2018;49(1):95-108. doi:10.1007/s40279-018-0994-1
9. Patterson SD, Hughes L, Warmington S, et al. Blood flow restriction exercise position stand: considerations of methodology, application, and safety. *Front Physiol*. 2019;10:533. doi: 10.3389/fphys.2019.00533
10. Manini TM, Clark BC. Blood flow restricted exercise and skeletal muscle health. *Exerc Sport Sci Rev*. 2009;37(2):78-85. doi: 10.1097/JES.0b013e31819c2e5c.
11. Goldberg AL, Etlinger JD, Goldspink DF, Jablecki C. Mechanism of work-induced hypertrophy of skeletal muscle. *Med Sci Sports*. 1975;7(3):185-198.
12. Schoenfeld BJ. The mechanisms of muscle hypertrophy and their application to resistance training. *J Strength Cond Res*. 2010;24(10):2857-2872. doi: 10.1519/JSC.0b013e3181e840f3.
13. Zou K, Meador BM, Johnson B, et al. The alpha(7) beta(1)-integrin increases muscle hypertrophy following multiple bouts of eccentric exercise. *J Appl Physiol (1985)*. 2011;111(4):1134-1141. doi: 10.1152/jap-physiol.00081.2011.
14. Adams GR. Autocrine and/or paracrine insulin-like growth factor-I activity in skeletal muscle. *Clin Orthop Relat Res*. 2002;(403 Suppl):S188-96. doi: 10.1097/00003086-200210001-00022.
15. Tatsumi R, Liu X, Pulido A, et al. Satellite cell activation in stretched skeletal muscle and the role of nitric oxide and hepatocyte growth factor. *Am J Physiol Cell Physiol*. 2006;290(6):1487-1494. doi: 10.1152/ajpcell.00513.2005.
16. Uchiyama S, Tsukamoto H, Yoshimura S, Tamaki T. Relationship between oxidative stress in muscle tissue and weight-lifting-induced muscle damage. *Pflugers Arch*. 2006;452(1):109-116. doi: 10.1007/s00424-005-0012-y.
17. Suga T, Okita K, Morita N, et al. Intramuscular metabolism during low-intensity resistance exercise with blood flow restriction. *J Appl Physiol (1985)*. 2009;106(4):1119-1124. doi: 10.1152/jap-physiol.90368.2008.
18. Cook SB, Kanaley JA, Ploutz-Snyder LL. Neuromuscular function following muscular unloading and blood flow restricted exercise. *Eur J Appl Physiol*. 2014;114(7):1357-1365. doi: 10.1007/s00421-014-2864-3.
19. Baar K, Esser K. Phosphorylation of p70(S6k) correlates with increased skeletal muscle mass following resistance exercise. *Am J Physiol*. 1999;276(1):120. doi: 10.1152/ajpcell.1999.276.1.C120.
20. Bodine SC, Stitt TN, Gonzalez M, et al. Akt/mTOR pathway is a crucial regulator of skeletal muscle hypertrophy and can prevent muscle atrophy in vivo. *Nat Cell Biol*. 2001;3(11):1014-1019. doi: 10.1038/ncb1101-1014.
21. Goto K, Ishii N, Kizuka T, Takamatsu K. The impact of metabolic stress on hormonal responses and muscular adaptations. *Med Sci Sports Exerc*. 2005;37(6):955-963. doi: 00005768-200506000-00009 [pii].
22. Suga T, Okita K, Takada S, et al. Effect of multiple set on intramuscular metabolic stress during low-intensity resistance exercise with blood flow restriction. *Eur J Appl Physiol*. 2012;112(11):3915-3920. doi: 10.1007/s00421-012-2377-x.
23. Kon M, Ikeda T, Homma T, Suzuki Y. Effects of low-intensity resistance exercise under acute systemic hypoxia on hormonal responses. *J Strength Cond Res*. 2012;26(3):611-617. doi: 10.1519/JSC.0b013e3182281c69.

24. Reeves GV, Kraemer RR, Hollander DB, et al. Comparison of hormone responses following light resistance exercise with partial vascular occlusion and moderately difficult resistance exercise without occlusion. *J Appl Physiol (1985)*. 2006;101(6):1616-1622. doi: 00440.2006 [pii].
25. Loenneke JP, Fahs CA, Thiebaud RS, et al. The acute muscle swelling effects of blood flow restriction. *Acta Physiol Hung*. 2012;99(4):400-410. doi: 10.1556/APhysiol.99.2012.4.4.
26. Fujita S, Abe T, Drummond MJ, et al. Blood flow restriction during low-intensity resistance exercise increases S6K1 phosphorylation and muscle protein synthesis. *J Appl Physiol (1985)*. 2007;103(3):903-910. doi: 10.1152/jappphysiol.00195.2007.
27. Fry CS, Glynn EL, Drummond MJ, et al. Blood flow restriction exercise stimulates mTORC1 signaling and muscle protein synthesis in older men. *J Appl Physiol (1985)*. 2010;108(5):1199-1209. doi: 10.1152/jappphysiol.01266.2009.
28. Laurentino GC, Ugrinowitsch C, Roschel H, et al. Strength training with blood flow restriction diminishes myostatin gene expression. *Med Sci Sports Exerc*. 2012;44(3):406-412. doi: 10.1249/MSS.0b013e318233b4bc.
29. Takarada Y, Sato Y, Ishii N. Effects of resistance exercise combined with vascular occlusion on muscle function in athletes. *Eur J Appl Physiol*. 2002;86(4):308-314. doi: 10.1007/s00421-001-0561-5
30. Yasuda T, Brechue WF, Fujita T, Shirakawa J, Sato Y, Abe T. Muscle activation during low-intensity muscle contractions with restricted blood flow. *J Sports Sci*. 2009;27(5):479-489. doi: 10.1080/02640410802626567.
31. Takano H, Morita T, Iida H, et al. Hemodynamic and hormonal responses to a short-term low-intensity resistance exercise with the reduction of muscle blood flow. *Eur J Appl Physiol*. 2005;95(1):65-73. doi: 10.1007/s00421-005-1389-1.
32. West DW, Phillips SM. Associations of exercise-induced hormone profiles and gains in strength and hypertrophy in a large cohort after weight training. *Eur J Appl Physiol*. 2012;112(7):2693-2702. doi: 10.1007/s00421-011-2246-z.
33. Mitchell CJ, Churchward-Venne TA, Bellamy L, Parise G, Baker SK, Phillips SM. Muscular and systemic correlates of resistance training-induced muscle hypertrophy. *PLoS One*. 2013;8(10):e78636. doi: 10.1371/journal.pone.0078636.
34. Lang F, Busch GL, Ritter M, et al. Functional significance of cell volume regulatory mechanisms. *Physiol Rev*. 1998;78(1):247-306. doi: 10.1152/physrev.1998.78.1.247.
35. Loenneke JP, Fahs CA, Rossow LM, Abe T, Bemben MG. The anabolic benefits of venous blood flow restriction training may be induced by muscle cell swelling. *Med Hypotheses*. 2012;78(1):151-154. doi: 10.1016/j.mehy.2011.10.014.
36. Lang F. Mechanisms and significance of cell volume regulation. *J Am Coll Nutr*. 2007;26(5 Suppl):613S-623S. doi: 10.1080/07315724.2007.10719667.
37. Low SY, Rennie MJ, Taylor PM. Signaling elements involved in amino acid transport responses to altered muscle cell volume. *FASEB J*. 1997;11(13):1111-1117. doi: 10.1096/fasebj.11.13.9367345.
38. Dangott B, Schultz E, Mozdziak PE. Dietary creatine monohydrate supplementation increases satellite cell mitotic activity during compensatory hypertrophy. *Int J Sports Med*. 2000;21(1):13-16. doi: 10.1055/s-2000-8848.
39. Gundermann DM, Fry CS, Dickinson JM, et al. Reactive hyperemia is not responsible for stimulating muscle protein synthesis following blood flow restriction exercise. *J Appl Physiol (1985)*. 2012;112(9):1520-1528. doi: 10.1152/jappphysiol.01267.2011.
40. Serrano AL, Baeza-Raja B, Perdiguero E, Jardi M, Munoz-Canoves P. Interleukin-6 is an essential regulator of satellite cell-mediated skeletal muscle hypertrophy. *Cell Metab*. 2008;7(1):33-44. doi: 10.1016/j.cmet.2007.11.011.
41. Nielsen AR, Pedersen BK. The biological roles of exercise-induced cytokines: IL-6, IL-8, and IL-15. *Appl Physiol Nutr Metab*. 2007;32(5):833-839. doi: h07-054 [pii].
42. Sorichter S, Mair J, Koller A, et al. Skeletal troponin I as a marker of exercise-induced muscle damage. *J Appl Physiol (1985)*. 1997;83(4):1076-1082. doi: 10.1152/jappphysiol.1997.83.4.1076.
43. Hather BM, Tesch PA, Buchanan P, Dudley GA. Influence of eccentric actions on skeletal muscle adaptations to resistance training. *Acta Physiol Scand*. 1991;143(2):177-185. doi: 10.1111/j.1748-1716.1991.tb09219.x.
44. Roig M, O'Brien K, Kirk G, et al. The effects of eccentric versus concentric resistance training on muscle strength and mass in healthy adults: A systematic review with meta-analysis. *Br J Sports Med*. 2009;43(8):556-568. doi: 10.1136/bjbm.2008.051417.
45. McHugh MP, Connolly DA, Eston RG, Gleim GW. Exercise-induced muscle damage and potential mechanisms for the repeated bout effect. *Sports Med*. 1999;27(3):157-170. doi: 10.2165/00007256-199927030-00002.
46. Thiebaud RS, Yasuda T, Loenneke JP, Abe T. Effects of low-intensity concentric and eccentric exercise combined with blood flow restriction on indices of exercise-induced muscle damage. *Interv Med Appl Sci*. 2013;5(2):53-59. doi: 10.1556/IMAS.5.2013.2.1.

47. Umbel JD, Hoffman RL, Dearth DJ, Chleboun GS, Manini TM, Clark BC. Delayed-onset muscle soreness induced by low-load blood flow-restricted exercise. *Eur J Appl Physiol*. 2009;107(6):687-695. doi: 10.1007/s00421-009-1175-6.
48. Henneman E, Somjen G, Carpenter DO. Functional significance of cell size in spinal motoneurons. *J Neurophysiol*. 1965;28:560-580. doi: 10.1152/jn.1965.28.3.560.
49. Wernbom M, Augustsson J, Raastad T. Ischemic strength training: A low-load alternative to heavy resistance exercise? *Scand J Med Sci Sports*. 2008;18(4):401-416. doi: 10.1111/j.1600-0838.2008.00788.x.
50. Fatela P, Reis JF, Mendonca GV, Avela J, Mil-Homens P. Acute effects of exercise under different levels of blood-flow restriction on muscle activation and fatigue. *Eur J Appl Physiol*. 2016;116(5):985-995. doi: 10.1007/s00421-016-3359-1.
51. Lauver JD, Cayot TE, Rotarius T, Scheuermann BW. The effect of eccentric exercise with blood flow restriction on neuromuscular activation, microvascular oxygenation, and the repeated bout effect. *Eur J Appl Physiol*. 2017;117(5):1005-1015. doi: 10.1007/s00421-017-3589-x.
52. Yasuda T, Brechue WF, Fujita T, Sato Y, Abe T. Muscle activation during low-intensity muscle contractions with varying levels of external limb compression. *J Sports Sci Med*. 2008;7(4):467-474. PMID: PMC3761922.
53. Wernbom M, Apro W, Paulsen G, Nilsen TS, Blomstrand E, Raastad T. Acute low-load resistance exercise with and without blood flow restriction increased protein signalling and number of satellite cells in human skeletal muscle. *Eur J Appl Physiol*. 2013;113(12):2953-2965. doi: 10.1007/s00421-013-2733-5.
54. Reynolds TH, Bodine SC, Lawrence JC. Control of Ser2448 phosphorylation in the mammalian target of rapamycin by insulin and skeletal muscle load. *J Biol Chem*. 2002;277(20):17657-17662. doi: 10.1074/jbc.M201142200.
55. Lee SJ, McPherron AC. Myostatin and the control of skeletal muscle mass. *Curr Opin Genet Dev*. 1999;9(5):604-607. doi: 10.1016/s0959-437x(99)00004-0.
56. Lee SJ, McPherron AC. Regulation of myostatin activity and muscle growth. *Proc Natl Acad Sci U S A*. 2001;98(16):9306-9311. doi: 10.1073/pnas.151270098.
57. McPherron AC, Lee SJ. Double muscling in cattle due to mutations in the myostatin gene. *Proc Natl Acad Sci U S A*. 1997;94(23):12457-12461. doi: 10.1073/pnas.94.23.12457.
58. McPherron AC, Lawler AM, Lee SJ. Regulation of anterior/posterior patterning of the axial skeleton by growth/differentiation factor 11. *Nat Genet*. 1999;22(3):260-264. doi: 10.1038/10320.
59. Lin J, Arnold HB, Della-Fera MA, Azain MJ, Hartzell DL, Baile CA. Myostatin knockout in mice increases myogenesis and decreases adipogenesis. *Biochem Biophys Res Commun*. 2002;291(3):701-706. doi: 10.1006/bbrc.2002.6500.
60. Rios R, Carneiro I, Arce VM, Devesa J. Myostatin regulates cell survival during C2C12 myogenesis. *Biochem Biophys Res Commun*. 2001;280(2):561-566. doi: 10.1006/bbrc.2000.4159.
61. Kawada S, Ishii N. Skeletal muscle hypertrophy after chronic restriction of venous blood flow in rats. *Med Sci Sports Exerc*. 2005;37(7):1144-1150. doi: 00005768-200507000-00010 [pii].
62. Drummond MJ, Fujita S, Abe T, Dreyer HC, Volpi E, Rasmussen BB. Human muscle gene expression following resistance exercise and blood flow restriction. *Med Sci Sports Exerc*. 2008;40(4):691-698. doi: 10.1249/MSS.0b013e318160ff84.
63. Ciciliot S, Schiaffino S. Regeneration of mammalian skeletal muscle. basic mechanisms and clinical implications. *Curr Pharm Des*. 2010;16(8):906-914. doi: 10.2174/138161210790883453.
64. Schiaffino S, Bormioli SP, Aloisi M. Cell proliferation in rat skeletal muscle during early stages of compensatory hypertrophy. *Virchows Arch B Cell Pathol*. 1972;11(3):268-273. doi: 10.1007/bf02889406.
65. Schiaffino S, Dyar KA, Ciciliot S, Blaauw B, Sandri M. Mechanisms regulating skeletal muscle growth and atrophy. *FEBS J*. 2013;280(17):4294-4314. doi: 10.1111/febs.12253.
66. Nielsen JL, Aagaard P, Bech RD, et al. Proliferation of myogenic stem cells in human skeletal muscle in response to low-load resistance training with blood flow restriction. *J Physiol (Lond)*. 2012;590(17):4351-4361. doi: 10.1113/jphysiol.2012.237008.
67. Bjørnsen T, Wernbom M, Løvstad AT, et al. Delayed myonuclear addition, myofiber hypertrophy and increases in strength with high-frequency low-load blood flow restricted training to volitional failure. *J Appl Physiol*. 2019;126(3):578-592. doi:10.1152/jappphysiol.00397.2018.
68. Scott BR, Loenneke JP, Slattery KM, Dascombe BJ. Exercise with blood flow restriction: An updated evidence-based approach for enhanced muscular development. *Sports Med*. 2015;45(3):313-325. doi: 10.1007/s40279-014-0288-1.
69. Kacin A, Rosenblatt B, Grapar Zargi T, Biswas A. Safety considerations with blood flow restricted resistance training. *Annales Kinesiologiae*. 2015;6(1):3. <https://www.dlib.si/stream/URN:NBN:SI:doc-MQ2NK37K/479cde7d-2300-4c99-b37a-1aa47cd157ce/PDF>. Accessed April 23, 2020.
70. Fahs CA, Loenneke JP, Rossow LM, Thiebaud RS, Bemben MG. Methodological considerations for blood flow

- restricted resistance exercise. *J Trainology*. 2012;1:14. http://trainology.org/PDF/4%20Methodological%20considerations%20for%20blood%20flow%20restricted%20resistance%20exercise%202012_1_14-22.pdf. Accessed April 23, 2020.
71. Abe T, Kearns CF, Sato Y. Muscle size and strength are increased following walk training with restricted venous blood flow from the leg muscle, kaatsu-walk training. *J Appl Physiol (1985)*. 2006;100(5):1460-1466. doi: 10.1152/jappphysiol.01267.2005.
 72. Takarada Y, Takazawa H, Ishii N. Applications of vascular occlusion diminish disuse atrophy of knee extensor muscles. *Med Sci Sports Exerc*. 2000;32(12):2035-2039. doi: 10.1097/00005768-200012000-00011.
 73. Kubota A, Sakuraba K, Sawaki K, Sumide T, Tamura Y. Prevention of disuse muscular weakness by restriction of blood flow. *Med Sci Sports Exerc*. 2008;40(3):529-534. doi: 10.1249/MSS.0b013e31815ddac6.
 74. Loenneke JP, Wilson JM, Marin PJ, Zourdos MC, Bemben MG. Low intensity blood flow restriction training: A meta-analysis. *Eur J Appl Physiol*. 2012;112(5):1849-1859. doi: 10.1007/s00421-011-2167-x.
 75. Lixandrao ME, Ugrinowitsch C, Berton R, et al. Magnitude of muscle strength and mass adaptations between high-load resistance training versus low-load resistance training associated with blood-flow restriction: A systematic review and meta-analysis. *Sports Med*. 2018;48(2):361-378. doi: 10.1007/s40279-017-0795-y.
 76. Hughes L, Paton B, Rosenblatt B, Gissane C, Paterson SD. Blood flow restriction training in clinical musculoskeletal rehabilitation: A systematic review and meta-analysis. *Br J Sports Med*. 2017;51(13):1003-1011. doi: 10.1136/bjsports-2016-097071.
 77. Burgomaster KA, Moore DR, Schofield LM, Phillips SM, Sale DG, Gibala MJ. Resistance training with vascular occlusion: Metabolic adaptations in human muscle. *Med Sci Sports Exerc*. 2003;35(7):1203-1208. doi: 10.1249/01.MSS.0000074458.71025.71.
 78. Moore DR, Burgomaster KA, Schofield LM, Gibala MJ, Sale DG, Phillips SM. Neuromuscular adaptations in human muscle following low intensity resistance training with vascular occlusion. *Eur J Appl Physiol*. 2004;92(4-5):399-406. doi: 10.1007/s00421-004-1072-y.
 79. Takarada Y, Takazawa H, Sato Y, Takebayashi S, Tanaka Y, Ishii N. Effects of resistance exercise combined with moderate vascular occlusion on muscular function in humans. *J Appl Physiol*. 2000;88(6):2097-2106. doi: 10.1152/jappl.2000.88.6.2097.
 80. Nielsen JL, Frandsen U, Prokhorova T, et al. Delayed effect of blood flow-restricted resistance training on rapid force capacity. *Med Sci Sports Exerc*. 2017;49(6):1157-1167. doi: 10.1249/MSS.0000000000001208.
 81. Abe T, Kawamoto K, Yasuda T, Kearns C, Midorikawa T, Sato Y. Eight days KAATSU-resistance training improved sprint but not jump performance in collegiate male track and field athletes. *Int J KAATSU Train Res*. 2005;1:19.
 82. Farup J, de Paoli F, Bjerg K, Riis S, Ringgard S, Vissing K. Blood flow restricted and traditional resistance training performed to fatigue produce equal muscle hypertrophy. *Scand J Med Sci Sports*. 2015;25(6):754-763. doi: 10.1111/sms.12396.
 83. Blazeovich AJ, Cannavan D, Coleman DR, Horne S. Influence of concentric and eccentric resistance training on architectural adaptation in human quadriceps muscles. *J Appl Physiol (1985)*. 2007;103(5):1565-1575. doi: 10.1152/jappphysiol.00578.2007.
 84. Damas F, Phillips SM, Lixandrao ME, et al. An inability to distinguish edematous swelling from true hypertrophy still prevents a completely accurate interpretation of the time course of muscle hypertrophy. *Eur J Appl Physiol*. 2016;116(2):445-446. doi: 10.1007/s00421-015-3287-5.
 85. Ladlow P, Coppack RJ, Dharm-Datta S, et al. Low-load resistance training with blood flow restriction improves clinical outcomes in musculoskeletal rehabilitation: a single-blind randomized controlled trial. *Front Physiol*. 2018;9:1269. doi: 10.3389/fphys.2018.01269.
 86. Manimmanakorn A, Hamlin MJ, Ross JJ, Taylor R, Manimmanakorn N. Effects of low-load resistance training combined with blood flow restriction or hypoxia on muscle function and performance in netball athletes. *J Sci Med Sport*. 2013;16(4):337-342. doi: 10.1016/j.jsams.2012.08.009.
 87. Thiebaud RS, Loenneke JP, Fahs CA, et al. The effects of elastic band resistance training combined with blood flow restriction on strength, total bone-free lean body mass and muscle thickness in postmenopausal women. *Clin Physiol Funct Imaging*. 2013;33(5):344-352. doi: 10.1111/cpf.12033.
 88. Libardi CA, Chacon-Mikahil MP, Cavaglieri CR, et al. Effect of concurrent training with blood flow restriction in the elderly. *Int J Sports Med*. 2015;36(5):395-399. doi: 10.1055/s-0034-1390496.
 89. Yasuda T, Fukumura K, Tomaru T, Nakajima T. Thigh muscle size and vascular function after blood flow-restricted elastic band training in older women. *Oncotarget*. 2016;7(23):33595-33607. doi: 10.18632/oncotarget.9564.
 90. Cook SB, LaRoche DP, Villa MR, Barile H, Manini TM. Blood flow restricted resistance training in older adults at risk of mobility limitations. *Exp Gerontol*. 2017;99:138-145. doi: 10.1016/j.exger.2017.10.004.
 91. Loenneke JP, Loprinzi PD, Abe T, et al. Arm circumference influences blood pressure even when applying the correct cuff size: Is a further correction needed? *Int J Cardiol*. 2016;202:743-744. doi: 10.1016/j.ijcard.2015.10.009.

92. Jessee MB, Buckner SL, Mouser JG, Mattocks KT, Loenneke JP. Letter to the editor: Applying the blood flow restriction pressure: the elephant in the room. *Am J Physiol Heart Circ Physiol*. 2016;310(1):132. doi: 10.1152/ajpheart.00820.2015.
93. Patterson SD, Hughes L, Head P, Warmington S, Brandner C. Blood flow restriction training: a novel approach to augment clinical rehabilitation: how to do it. *Br J Sports Med*. 2017;51(23):1648-1649. doi: 10.1136/bjsports-2017-097738.
94. Brandner CR, Warmington SA, Kidgell DJ. Corticomotor excitability is increased following an acute bout of blood flow restriction resistance exercise. *Front Hum Neurosci*. 2015;9:652. doi: 10.3389/fnhum.2015.00652.
95. Younger AS, McEwen JA, Inkpen K. Wide contoured thigh cuffs and automated limb occlusion measurement allow lower tourniquet pressures. *Clin Orthop Relat Res*. 2004;(428):286-93. doi: 10.1097/01.blo.0000142625.82654.b3.
96. Kubota A, Sakuraba K, Koh S, Ogura Y, Tamura Y. Blood flow restriction by low compressive force prevents disuse muscular weakness. *J Sci Med Sport*. 2011;14(2):95-99. doi: 10.1016/j.jsams.2010.08.007.
97. Cook SB, Clark BC, Ploutz-Snyder LL. Effects of exercise load and blood-flow restriction on skeletal muscle function. *Med Sci Sports Exerc*. 2007;39(10):1708-1713. doi: 10.1249/mss.0b013e31812383d6.
98. Mattocks KT, Jessee MB, Counts BR, et al. The effects of upper body exercise across different levels of blood flow restriction on arterial occlusion pressure and perceptual responses. *Physiol Behav*. 2017;171:181-186. doi: 10.1016/j.physbeh.2017.01.015.
99. Jessee MB, Dankel SJ, Buckner SL, Mouser JG, Mattocks KT, Loenneke JP. The cardiovascular and perceptual response to very low load blood flow restricted exercise. *Int J Sports Med*. 2017;38(8):597-603. doi: 10.1055/s-0043-109555.
100. Crenshaw AG, Hargens AR, Gershuni DH, Rydevik B. Wide tourniquet cuffs more effective at lower inflation pressures. *Acta Orthop Scand*. 1988;59(4):447-451. doi: 10.3109/17453678809149401.
101. Loenneke JP, Fahs CA, Rossow LM, et al. Effects of cuff width on arterial occlusion: Implications for blood flow restricted exercise. *Eur J Appl Physiol*. 2012;112(8):2903-2912. doi: 10.1007/s00421-011-2266-8.
102. Rossow LM, Fahs CA, Loenneke JP, et al. Cardiovascular and perceptual responses to blood-flow-restricted resistance exercise with differing restrictive cuffs. *Clin Physiol Funct Imaging*. 2012;32(5):331-337. doi: 10.1111/j.1475-097X.2012.01131.x.
103. Mouser JG, Ade CJ, Black CD, Bemben DA, Bemben MG. Brachial blood flow under relative levels of blood flow restriction is decreased in a nonlinear fashion. *Clin Physiol Funct Imaging*. 2018;38(3):425-430. doi: 10.1111/cpf.12432.
104. Mouser JG, Dankel SJ, Jessee MB, et al. A tale of three cuffs: the hemodynamics of blood flow restriction. *Eur J Appl Physiol*. 2017;117(7):1493-1499. doi: 10.1007/s00421-017-3644-7.
105. Kacin A, Strazar K. Frequent low-load ischemic resistance exercise to failure enhances muscle oxygen delivery and endurance capacity. *Scand J Med Sci Sports*. 2011;21(6):231. doi: 10.1111/j.1600-0838.2010.01260.x.
106. Ellefsen S, Hammarstrom D, Strand TA, et al. Blood flow-restricted strength training displays high functional and biological efficacy in women: a within-subject comparison with high-load strength training. *Am J Physiol Regul Integr Comp Physiol*. 2015;309(7):767. doi: 10.1152/ajpregu.00497.2014.
107. Laurentino GC, Loenneke JP, Teixeira EL, Nakajima E, Iared W, Tricoli V. The effect of cuff width on muscle adaptations after blood flow restriction training. *Med Sci Sports Exerc*. 2016;48(5):920-925. doi: 10.1249/MSS.0000000000000833.
108. Folland JP, Williams AG. The adaptations to strength training: morphological and neurological contributions to increased strength. *Sports Med*. 2007;37(2):145-168. doi: 10.2165/00007256-200737020-00004.
109. Kraemer WJ, Ratamess NA. Fundamentals of resistance training: progression and exercise prescription. *Med Sci Sports Exerc*. 2004;36(4):674-688. doi: 10.1249/01.mss.0000121945.36635.61.
110. Garber CE, Blissmer B, Deschenes MR, et al. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc*. 2011;43(7):1334-1359. doi: 10.1249/MSS.0b013e3181213febf.
111. Lixandrao ME, Ugrinowitsch C, Laurentino G, et al. Effects of exercise intensity and occlusion pressure after 12 weeks of resistance training with blood-flow restriction. *Eur J Appl Physiol*. 2015;115(12):2471-2480. doi: 10.1007/s00421-015-3253-2.
112. Counts BR, Dankel SJ, Barnett BE, et al. Influence of relative blood flow restriction pressure on muscle activation and muscle adaptation. *Muscle Nerve*. 2016;53(3):438-445. doi: 10.1002/mus.24756.
113. Dankel SJ, Jessee MB, Abe T, Loenneke JP. The effects of blood flow restriction on upper-body musculature located distal and proximal to applied pressure. *Sports Med*. 2016;46(1):23-33. doi: 10.1007/s40279-015-0407-7.
114. Yasuda T, Ogasawara R, Sakamaki M, Ozaki H, Sato Y, Abe T. Combined effects of low-intensity blood flow restriction training and high-intensity resistance

- training on muscle strength and size. *Eur J Appl Physiol.* 2011;111(10):2525-2533. doi: 10.1007/s00421-011-1873-8.
115. Yasuda T, Loenneke JP, Thiebaud RS, Abe T. Effects of blood flow restricted low-intensity concentric or eccentric training on muscle size and strength. *PLoS One.* 2012;7(12):e52843. doi: 10.1371/journal.pone.0052843.
 116. Yasuda T, Fujita S, Ogasawara R, Sato Y, Abe T. Effects of low-intensity bench press training with restricted arm muscle blood flow on chest muscle hypertrophy: a pilot study. *Clin Physiol Funct Imaging.* 2010;30(5):338-343. doi: 10.1111/j.1475-097X.2010.00949.x.
 117. Ozaki H, Yasuda T, Ogasawara R, Sakamaki-Sunaga M, Naito H, Abe T. Effects of high-intensity and blood flow-restricted low-intensity resistance training on carotid arterial compliance: role of blood pressure during training sessions. *Eur J Appl Physiol.* 2013;113(1):167-174. doi: 10.1007/s00421-012-2422-9.
 118. Yasuda T, Ogasawara R, Sakamaki M, Bemben MG, Abe T. Relationship between limb and trunk muscle hypertrophy following high-intensity resistance training and blood flow-restricted low-intensity resistance training. *Clin Physiol Funct Imaging.* 2011;31(5):347-351. doi: 10.1111/j.1475-097X.2011.01022.x.
 119. Madarame H, Neya M, Ochi E, Nakazato K, Sato Y, Ishii N. Cross-transfer effects of resistance training with blood flow restriction. *Med Sci Sports Exerc.* 2008;40(2):258-263. doi: 10.1249/mss.0b013e31815c6d7e.
 120. Manini TM, Yarrow JF, Buford TW, Clark BC, Conover CF, Borst SE. Growth hormone responses to acute resistance exercise with vascular restriction in young and old men. *Growth Horm IGF Res.* 2012;22(5):167-172. doi: 10.1016/j.ghir.2012.05.002.
 121. Fahs CA, Loenneke JP, Thiebaud RS, et al. Muscular adaptations to fatiguing exercise with and without blood flow restriction. *Clin Physiol Funct Imaging.* 2015;35(3):167-176. doi: 10.1111/cpf.12141.
 122. Loenneke JP, Fahs CA, Wilson JM, Bemben MG. Blood flow restriction: the metabolite/volume threshold theory. *Med Hypotheses.* 2011;77(5):748-752. doi: 10.1016/j.mehy.2011.07.029.
 123. Martin-Hernandez J, Marin PJ, Menendez H, Ferrero C, Loenneke JP, Herrero AJ. Muscular adaptations after two different volumes of blood flow-restricted training. *Scand J Med Sci Sports.* 2013;23(2):114. doi: 10.1111/sms.12036.
 124. Loenneke JP, Kearney ML, Thrower AD, Collins S, Pujol TJ. The acute response of practical occlusion in the knee extensors. *J Strength Cond Res.* 2010;24(10):2831-2834. doi: 10.1519/JSC.0b013e3181f0ac3a.
 125. Loenneke JP, Balapur A, Thrower AD, Barnes JT, Pujol TJ. The perceptual responses to occluded exercise. *Int J Sports Med.* 2011;32(3):181-184. doi: 10.1055/s-0030-1268472.
 126. Patterson SD, Ferguson RA. Enhancing strength and postocclusive calf blood flow in older people with training with blood-flow restriction. *J Aging Phys Act.* 2011;19(3):201-213. doi: 10.1123/japa.19.3.201.
 127. Patterson SD, Ferguson RA. Increase in calf post-occlusive blood flow and strength following short-term resistance exercise training with blood flow restriction in young women. *Eur J Appl Physiol.* 2010;108(5):1025-1033. doi: 10.1007/s00421-009-1309-x.
 128. Ladlow P, Coppack RJ, Dharm-Datta S, et al. Low-load resistance training with blood flow restriction improves clinical outcomes in musculoskeletal rehabilitation: a single-blind randomized controlled trial. *Front Physiol.* 2018;9:1269. doi: 10.3389/fphys.2018.01269.
 129. Yasuda T, Loenneke JP, Ogasawara R, Abe T. Influence of continuous or intermittent blood flow restriction on muscle activation during low-intensity multiple sets of resistance exercise. *Acta Physiol Hung.* 2013;100(4):419-426. doi: 10.1556/APhysiol.100.2013.4.6.
 130. Ohta H, Kurosawa H, Ikeda H, Iwase Y, Satou N, Nakamura S. Low-load resistance muscular training with moderate restriction of blood flow after anterior cruciate ligament reconstruction. *Acta Orthop Scand.* 2003;74(1):62-68. doi: 10.1080/00016470310013680.
 131. Kraemer WJ, Patton JF, Gordon SE, et al. Compatibility of high-intensity strength and endurance training on hormonal and skeletal muscle adaptations. *J Appl Physiol.* 1995;78(3):976-989. doi: 10.1152/jappl.1995.78.3.976.
 132. Bennett H, Slattery F. Effects of blood flow restriction training on aerobic capacity and performance: a systematic review. *J Strength Cond Res.* 2019;33(2):572-583. doi: 10.1519/JSC.0000000000002963.
 133. Abe T, Loenneke JP, Fahs CA. Exercise intensity and muscle hypertrophy in blood flow-restricted limbs and non-restricted muscles: a brief review. *Clin Physiol.* 2012;32(4):247-52. doi: 10.1111/j.1475-097X.2012.01126.x..
 134. Conceição MS, Junior EMM, Telles GD, et al. Augmented anabolic responses following 8-weeks cycling with blood flow restriction. *Med Sci Sports Exerc.* August 2018; 51(1):84-93. doi:10.1249/MSS.0000000000001755.
 135. Park S, Kim JK, Choi HM, Kim HG, Beekley MD, Nho H. Increase in maximal oxygen uptake following 2-week walk training with blood flow occlusion in athletes. *Eur J Appl Physiol.* 2010;109(4):591-600. doi: 10.1007/s00421-010-1377-y.
 136. Ilett M, Rantalainen T, Keske M, May A, Warming-ton S. The effects of restriction pressures on the acute responses to blood flow restriction exercise. *Front Physiol.* 2019;10:1018. doi: 10.3389/fphys.2019.01018

137. Abe T, Fujita S, Nakajima T, et al. Effects of low-intensity cycle training with restricted leg blood flow on thigh muscle volume and VO₂max in young men. *J Sports Sci Med.* 2010;9(3):452-458.
138. Tabata I, Atomi Y, Kanehisa H, Miyashita M. Effect of high-intensity endurance training on isokinetic muscle power. *Eur J Appl Physiol Occup Physiol.* 1990;60(4):254-258. doi: 10.1007/bf00379392.
139. Gaesser GA, Rich RG. Effects of high- and low-intensity exercise training on aerobic capacity and blood lipids. *Med Sci Sports Exerc.* 1984;16(3):269-274.
140. Wenger HA, Bell GJ. The interactions of intensity, frequency and duration of exercise training in altering cardiorespiratory fitness. *Sports Med.* 1986;3(5):346-356. doi: 10.2165/00007256-198603050-00004.
141. Ozaki H, Brechue WF, Sakamaki M, et al. Metabolic and cardiovascular responses to upright cycle exercise with leg blood flow reduction. *J Sports Sci Med.* 2010;9(2):224-230.
142. Paffenbarger RS Jr, Laughlin ME, Gima AS, Black RA. Work activity of longshoremen as related to death from coronary heart disease and stroke. *N Engl J Med.* 1970;282(20):1109-1114. doi: 10.1056/NEJM197005142822001.
143. Prado CM, Purcell SA, Alish C, et al. Implications of low muscle mass across the continuum of care: a narrative review. *Ann Med.* 2018;1-19. doi: 10.1080/07853890.2018.1511918.
144. Kim C-H, Wheatley CM, Behnia M, Johnson BD. The effect of aging on relationships between lean body mass and VO₂max in rowers. *PLoS One.* 2016;11(8):e0160275. doi: 10.1371/journal.pone.0160275.
145. Proctor DN, Joyner MJ. Skeletal muscle mass and the reduction of VO₂ max in trained older subjects. *J Appl Physiol.* 1997;82(5):1411-1415. doi: 10.1152/jap-1997.82.5.1411.
146. Ozaki H, Miyachi M, Nakajima T, Abe T. Effects of 10 weeks walk training with leg blood flow reduction on carotid arterial compliance and muscle size in the elderly adults. *Angiology.* 2011;62(1):81-86. doi: 10.1177/0003319710375942.
147. Sundberg CJ. Exercise and training during graded leg ischaemia in healthy man with special reference to effects on skeletal muscle. *Acta Physiol Scand Suppl.* 1994;615:1-50.
148. de Oliveira MFM, Caputo F, Corvino RB, Denadai BS. Short-term low-intensity blood flow restricted interval training improves both aerobic fitness and muscle strength. *Scand J Med Sci Sports.* 2016;26(9):1017-1025. doi: 10.1111/sms.12540.
149. Loenneke JP, Thrower AD, Balapur A, Barnes JT, Pujol TJ. Blood flow-restricted walking does not result in an accumulation of metabolites. *Clin Physiol Funct Imaging.* 2012;32(1):80-82. doi: 10.1111/j.1475-097X.2011.01059.x.
150. Christiansen D, Eibye KH, Rasmussen V, et al. Cycling with blood flow restriction improves performance and muscle K⁺ regulation and alters the effect of antioxidant infusion in humans. *J Physiol.* 2019;597(9):2421-2444. doi:10.1113/JP277657.
151. Pignanelli C, Petrick HL, Keyvani F, et al. Low-load resistance training to task-failure with and without blood flow restriction: muscular functional and structural adaptations. *Am J Physiol Regul Integr Comp Physiol.* 2020;318(2):R284-R295. doi:10.1152/ajp-regu.00243.2019.
152. William Ursprung TAU-SA, Smith JD. The effects of blood flow restriction training on vo₂max and 1.5 mile run performance. *Int J Exerc Sci: Conference Proceedings.* 2017;2(9):108.
153. Held S, Behringer M, Donath L. Low intensity rowing with blood flow restriction over 5 weeks increases V̇O₂max in elite rowers: a randomized controlled trial. *J Sci Med Sport.* 2020 Mar;23(3):304-308. doi:10.1016/j.jsams.2019.10.002.
154. Pope ZK, Willardson JM, Schoenfeld BJ. Exercise and blood flow restriction. *J Strength Cond Res.* 2013;27(10):2914-2926. doi:10.1519/JSC.0b013e3182874721.
155. Brandner CR, May AK, Clarkson MJ, Warmington SA. Reported side-effects and safety considerations for the use of blood flow restriction during exercise in practice and research. *Tech Orthop.* 2018;32(2):114-121. doi:10.1097/BTO.0000000000000259.
156. Nakajima T, Kurano M, Iida H, et al. Use and safety of KAATSU training: results of a national survey. *Int J KAATSU Train Res.* 2006;2(1):5-13. doi:10.3806/ijk-tr.2.5.
157. Patterson SD, Brandner CR. The role of blood flow restriction training for applied practitioners: a questionnaire-based survey. *J Sports Sci.* 2017;36(2):123-130. doi:10.1080/02640414.2017.1284341.
158. Brandner CR, Warmington SA. Delayed onset muscle soreness and perceived exertion after blood flow restriction exercise. *J Strength Cond Res.* 2017;31(11):3101-3108. doi:10.1519/JSC.0000000000001779.
159. Wernbom M, Järrebring R, Andreasson MA, Augustsson J. Acute effects of blood flow restriction on muscle activity and endurance during fatiguing dynamic knee extensions at low load. *J Strength Cond Res.* 2009;23(8):2389-2395. doi:10.1519/JSC.0b013e3181bc1c2a.
160. Wernbom M, Paulsen G, Nilsen TS, Hisdal J, Raastad T. Sarcolemmal permeability and muscle damage as hypertrophic stimuli in blood flow restricted resistance exercise (Reply to Loenneke and Abe). *Eur J Appl Physiol.*

- 2012;112(9):3447-3449. doi:10.1007/s00421-012-2309-9.
161. Karabulut M, Sherk VD, Bemben DA, Bemben MG. Inflammation marker, damage marker and anabolic hormone responses to resistance training with vascular restriction in older males. *Clin Physiol Funct Imaging*. 2013;33(5):393-399. doi:10.1111/cpf.12044.
 162. Iversen E, Rstad V. Low-load ischemic exercise-Induced rhabdomyolysis. *Clin J Sport Med*. 2010;20(3):218-219. doi:10.1097/JSM.0b013e3181df8d10.
 163. Tabata S, Suzuki Y, Azuma K, Matsumoto H. Rhabdomyolysis after performing blood flow restriction training: a case report. *J Strength Cond Res*. 2016;30(7):2064-2068. doi:10.1519/JSC.0000000000001295.
 164. Lee AJ, Maddix DS. Rhabdomyolysis secondary to a drug interaction between simvastatin and clarithromycin. *Ann Pharmacother*. 2001;35(1):26-31. doi:10.1345/aph.10177.
 165. Ozawa Y, Koto T, Shinoda H, Tsubota K. Vision loss by central retinal vein occlusion After Kaatsu training. *Medicine (Baltimore)*. 2015;94(36):e1515. doi:10.1097/MD.0000000000001515.
 166. Nakajima T, Morita T, Sato Y. Key considerations when conducting KAATSU training. *Int J KAATSU Train Res*. 2011;7(1):1-6. doi:10.3806/ijkr.7.1
 167. Kondo T. Rumpel-Leede phenomenon associated with blood flow restriction training. *QJM An Int J Med*. 2017;111(5):345. doi:10.1093/qjmed/hcx244.
 168. Martín-Hernández J, Santos-Lozano A, Foster C, Lucia A. Syncope episodes and blood flow restriction training. *Clin J Sport Med Off J Can Acad Sport Med*. 2017;0(0):1-3. doi:10.1097/JSM.0000000000000496.
 169. Noto T, Hashimoto G, Takagi T, et al. Paget-Schroetter syndrome resulting from thoracic outlet syndrome and KAATSU training. *Intern Med*. 2017;56(19):2595-2601. doi:10.2169/internalmedicine.7937-16.
 170. Loenneke JP, Wilson JM, Wilson GJ, Pujol TJ, Bemben MG. Potential safety issues with blood flow restriction training. *Scand J Med Sci Sport*. 2011;21(4):510-518. doi:10.1111/j.1600-0838.2010.01290.x.
 171. Estebe JP, Le Naoures A, Chemaly L, Ecoffey C. Tourniquet pain in a volunteer study: effect of changes in cuff width and pressure. *Anaesthesia*. 2000;55(1):21-26. doi:10.1046/j.1365-2044.2000.01128.x.
 172. McEwen JA, Inkpen K. Safety: preventing skin injuries. *Surg Technol*. 2002;34:7-15.
 173. Hollander DB, Reeves GV, Clavier JD, Francois MR, Thomas C, Kraemer RR. Partial occlusion during resistance exercise alters effort sense and pain. *J Strength Cond Res*. 2010;24(1):235-243. doi:10.1519/JSC.0b013e3181c7badf.
 174. Fitschen PJ, Kistler BM, Jeong JH, et al. Perceptual effects and efficacy of intermittent or continuous blood flow restriction resistance training. *Clin Physiol Funct Imaging*. 2014;34(5):356-363. doi:10.1111/cpf.12100.
 175. McHugh MP. Recent advances in the understanding of the repeated bout effect: The protective effect against muscle damage from a single bout of eccentric exercise. *Scand J Med Sci Sport*. 2003;13(2):88-97. doi:10.1034/j.1600-0838.2003.02477.x.
 176. Martín-Hernández J, Ruiz-Aguado J, Herrero AJ, et al. Adaptation of perceptual responses to low-load blood flow restriction training. *J Strength Cond Res*. 2016;31(3):765-772. doi:10.1519/JSC.0000000000001478.
 177. Hughes L, Patterson SD, Haddad F, et al. Examination of the comfort and pain experienced with blood flow restriction training during post-surgery rehabilitation of anterior cruciate ligament reconstruction patients: a UK National Health Service trial. *Phys Ther Sport*. 2019;39:90-98. doi:10.1016/j.ptsp.2019.06.014.
 178. Vieira PJC, Chiappa GR, Umpierre D, Stein R, Ribeiro JP. Hemodynamic responses to resistance exercise with restricted blood flow in young and older men. *J Strength Cond Res*. 2013;27(8):2288-2294. doi:10.1519/JSC.0b013e318278f21f.
 179. Downs ME, Hackney KJ, Martin D, et al. Acute vascular and cardiovascular responses to blood flow-restricted exercise. *Med Sci Sports Exerc*. 2014;46(8):1489-1497. doi:10.1249/MSS.0000000000000253.
 180. Araújo JP, Silva ED, Silva JCG, et al. The acute effect of resistance exercise with blood flow restriction with hemodynamic variables on hypertensive subjects. *J Hum Kinet*. 2014;43(1):79-85. doi:10.2478/hukin-2014-0092.
 181. Sakamaki M, Fujita S, Sato Y, Bemben MG, Abe T. Blood pressure response to slow walking combined with KAATSU in the elderly. *Int J KAATSU Train Res*. 2008;4(1):17-20. doi:10.3806/ijkr.4.17.
 182. Spranger MD, Krishnan AC, Levy PD, O'Leary DS, Smith SA. Blood flow restriction training and the exercise pressor reflex: a call for concern. *Am J Physiol - Hear Circ Physiol*. 2015;309(9):1440-1452. doi:10.1152/ajpheart.00208.2015.
 183. Neto GR, Sousa MSC, Costa e Silva GV, Gil ALS, Salles BF, Novaes JS. Acute resistance exercise with blood flow restriction effects on heart rate, double product, oxygen saturation and perceived exertion. *Clin Physiol Funct Imaging*. 2014;36(1):53-59. doi:10.1111/cpf.12193.
 184. Neto GR, Santos HH, Sousa JBC, et al. Effects of high-intensity blood flow restriction exercise on muscle fatigue. *J Hum Kinet*. 2014;41(1):163-172. doi:10.2478/hukin-2014-0044.
 185. Poton R, Polito MD. Hemodynamic response to resistance exercise with and without blood flow restriction in healthy subjects. *Clin Physiol Funct Imaging*. 2016;36(3):231-236. doi:10.1111/cpf.12218.

186. Pinto RR, Karabulut M, Poton R, Polito MD. Acute resistance exercise with blood flow restriction in elderly hypertensive women: haemodynamic, rating of perceived exertion and blood lactate. *Clin Physiol Funct Imaging*. 2018 Jan;38(1):17-24. doi:10.1111/cpf.12376.
187. Brandner CR, Kidgell DJ, Warmington SA. Unilateral bicep curl hemodynamics: low-pressure continuous vs high-pressure intermittent blood flow restriction. *Scand J Med Sci Sport*. 2015;25(6):770-777. doi:10.1111/sms.12297.
188. Lentini AC, McKelvie RS, McCartney N, Tomlinson CW, MacDougall JD. Left ventricular response in healthy young men during heavy-intensity weight-lifting exercise. *J Appl Physiol*. 1993;75(6):2703-2710. doi:10.1152/jappl.1993.75.6.2703.
189. Adams J, Cline MJ, Hubbard M, McCullough T, Hartman J. A new paradigm for post-cardiac event resistance exercise guidelines. *Am J Cardiol*. 2006;97(2):281-286. doi:10.1016/j.amjcard.2005.08.035.
190. Dankel S, Buckner S, Counts B, et al. The acute muscular response to two distinct blood flow restriction protocols. *Physiol Int*. 2017;104(1):64-76. doi:10.1556/2060.104.2017.1.1.
191. Jarrett P, Ritchie I, Albadran L, Glen S., Bridges A., M E. Do thigh tourniquets contribute to the formation of intra-operative venous emboli? *Acta Orthop Belg*. 2004;70(3):253-529.
192. Madarame H, Kurano M, Takano H, et al. Effects of low-intensity resistance exercise with blood flow restriction on coagulation system in healthy subjects. *Clin Physiol Funct Imaging*. 2010;30(3):210-213. doi:10.1111/j.1475-097X.2010.00927.x.
193. Clark BC, Manini TM, Hoffman RL, et al. Relative safety of 4 weeks of blood flow-restricted resistance exercise in young, healthy adults. *Scand J Med Sci Sports*. 2011;21(5):653-662. doi:10.1111/j.1600-0838.2010.01100.x.
194. Yasuda T, Fukumura K, Fukuda T, et al. Muscle size and arterial stiffness after blood flow-restricted low-intensity resistance training in older adults. *Scand J Med Sci Sport*. 2014;24(5):799-806. doi:10.1111/sms.12087.
195. Yasuda T, Fukumura K, Fukuda T, et al. Effects of low-intensity, elastic band resistance exercise combined with blood flow restriction on muscle activation. *Scand J Med Sci Sport*. 2014;24(1):55-61. doi:10.1111/j.1600-0838.2012.01489.x.
196. Madarame H, Kurano M, Fukumura K, Fukuda T, Nakajima T. Haemostatic and inflammatory responses to blood flow-restricted exercise in patients with ischaemic heart disease: A pilot study. *Clin Physiol Funct Imaging*. 2013;33(1):11-17. doi:10.1111/j.1475-097X.2012.01158.x.
197. Tennent DJ, Hylden CM, Johnson AE, Burns TC, Wilken JM, Owens JG. Blood flow restriction training after knee arthroscopy: a randomized controlled pilot study. *Clin J Sport Med*. 2017;27(3):245-252. doi:10.1097/JSM.0000000000000377.
198. Minniti MC, Statkevich AP, Kelly RL, et al. The safety of blood flow restriction training as a therapeutic intervention for patients with musculoskeletal disorders: a systematic review. *Am J Sports Med*. 2019:1-13. doi:10.1177/0363546519882652.
199. Giles L, Webster KE, McClelland J, Cook JL. Quadriceps strengthening with and without blood flow restriction in the treatment of patellofemoral pain: A double-blind randomised trial. *Br J Sports Med*. 2017;51(23):1688-1694. doi:10.1136/bjsports-2016-096329.
200. Ferraz RB, Gualano B, Rodrigues R, et al. Benefits of resistance training with blood flow restriction in knee osteoarthritis. *Med Sci Sports Exerc*. 2018;50(5):897-905. doi:10.1249/MSS.0000000000001530.
201. Rodrigues R, Ferraz RB, Kurimori CO, et al. Low-load resistance training with blood flow restriction increases muscle function, mass and functionality in women with rheumatoid arthritis. *Arthritis Care Res (Hoboken)*. 2019. doi:10.1002/acr.23911.
202. Hughes L, Rosenblatt B, Haddad F, et al. Comparing the effectiveness of blood flow restriction and traditional heavy load resistance training in the post-surgery rehabilitation of anterior cruciate ligament reconstruction patients: a UK National Health Service randomised controlled trial. *Sport Med*. 2019;49(11):1787-1805. doi:10.1007/s40279-019-01137-2.
203. Allison RC, Bedsole DL. The other medical causes of rhabdomyolysis. *Am J Med Sci*. 2003;326(2):79-88. doi:10.1097/00000441-200308000-00005.
204. Atherton PJ, Greenhaff PL, Phillips SM, Bodine SC, Adams CM, Lang CH. Control of skeletal muscle atrophy in response to disuse: clinical/preclinical contentions and fallacies of evidence. *Am J Physiol Endocrinol Metab*. 2016;311(3):594. doi: 10.1152/ajpendo.00257.2016.
205. Crossland H, Constantin-Teodosiu D, Greenhaff PL, Gardiner SM. Low-dose dexamethasone prevents endotoxaemia-induced muscle protein loss and impairment of carbohydrate oxidation in rat skeletal muscle. *J Physiol*. 2010;588(8):1333-1347. doi: 10.1113/jphysiol.2009.183699.
206. Snyder-Mackler L, Ladin Z, Schepsis AA, Young JC. Electrical stimulation of the thigh muscles after reconstruction of the anterior cruciate ligament. Effects of electrically elicited contraction of the quadriceps femoris and hamstring muscles on gait and on strength of the thigh muscles. *J Bone Joint Surg Am*. 1991;73(7):1025-1036.
207. Snyder-Mackler L, Delitto A, Bailey SL, Stralka SW. Strength of the quadriceps femoris muscle and functional recovery after reconstruction of the anterior

- cruciate ligament. A prospective, randomized clinical trial of electrical stimulation. *J Bone Joint Surg Am*. 1995;77(8):1166-1173. doi: 10.2106/00004623-199508000-00004.
208. Lewek M, Rudolph K, Axe M, Snyder-Mackler L. The effect of insufficient quadriceps strength on gait after anterior cruciate ligament reconstruction. *Clin Biomech (Bristol, Avon)*. 2002;17(1):56-63. doi: 10.1016/s0268-0033(01)00097-3.
 209. Eitzen I, Holm I, Risberg MA. Preoperative quadriceps strength is a significant predictor of knee function two years after anterior cruciate ligament reconstruction. *Br J Sports Med*. 2009;43(5):371-376. doi: 10.1136/bjism.2008.057059.
 210. Keller K, Engelhardt M. Muscle atrophy caused by limited mobilisation. *Sportverletz Sportschaden*. 2013;27(2):91-95. doi: 10.1055/s-0033-1335159.
 211. Wall BT, Dirks ML, Snijders T, Senden JM, Dolmans J, van Loon LJ. Substantial skeletal muscle loss occurs during only 5 days of disuse. *Acta Physiol (Oxf)*. 2014;210(3):600-611. doi: 10.1111/apha.12190.
 212. Fukunaga T, Johnson CD, Nicholas SJ, McHugh MP. Muscle hypotrophy, not inhibition, is responsible for quadriceps weakness during rehabilitation after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc*. 2019;27(2):573-579. doi: 10.1007/s00167-018-5166-1.
 213. Thomas AC, Wojtys EM, Brandon C, Palmieri-Smith RM. Muscle atrophy contributes to quadriceps weakness after anterior cruciate ligament reconstruction. *J Sci Med Sport*. 2016;19(1):7-11. doi: 10.1016/j.jsams.2014.12.009.
 214. Konishi Y, Ikeda K, Nishino A, Sunaga M, Aihara Y, Fukubayashi T. Relationship between quadriceps femoris muscle volume and muscle torque after anterior cruciate ligament repair. *Scand J Med Sci Sports*. 2007;17(6):656-661. doi: 10.1111/j.1600-0838.2006.00619.x.
 215. Konishi Y, Oda T, Tsukazaki S, Kinugasa R, Fukubayashi T. Relationship between quadriceps femoris muscle volume and muscle torque at least 18 months after anterior cruciate ligament reconstruction. *Scand J Med Sci Sports*. 2012;22(6):791-796. doi: 10.1111/j.1600-0838.2011.01332.x.
 216. Haggmark T, Jansson E, Eriksson E. Fiber type area and metabolic potential of the thigh muscle in man after knee surgery and immobilization. *Int J Sports Med*. 1981;2(1):12-17. doi: 10.1055/s-2008-1034577.
 217. Lindboe CF, Platou CS. Effect of immobilization of short duration on the muscle fibre size. *Clin Physiol*. 1984;4(2):183-188. doi: 10.1111/j.1475-097x.1984.tb00234.x.
 218. Bamman MM, Clarke MS, Feedback DL, et al. Impact of resistance exercise during bed rest on skeletal muscle sarcopenia and myosin isoform distribution. *J Appl Physiol (1985)*. 1998;84(1):157-163. doi: 10.1152/jap-1998.84.1.157.
 219. Hvid LG, Ortenblad N, Aagaard P, Kjaer M, Suetta C. Effects of ageing on single muscle fibre contractile function following short-term immobilisation. *J Physiol*. 2011;589(Pt 19):4745-4757. doi: 10.1113/jphysiol.2011.215434.
 220. de Boer MD, Selby A, Atherton P, et al. The temporal responses of protein synthesis, gene expression and cell signaling in human quadriceps muscle and patellar tendon to disuse. *J Physiol*. 2007;585(Pt 1):241-251. doi: jphysiol.2007.142828 [pii].
 221. Krawiec BJ, Frost RA, Vary TC, Jefferson LS, Lang CH. Hindlimb casting decreases muscle mass in part by proteasome-dependent proteolysis but independent of protein synthesis. *Am J Physiol Endocrinol Metab*. 2005;289(6):969. doi: 10.1152/ajpendo.00126.2005 .
 222. Atherton PJ, Greenhaff PL, Phillips SM, Bodine SC, Adams CM, Lang CH. Control of skeletal muscle atrophy in response to disuse: clinical/preclinical contentions and fallacies of evidence. *Am J Physiol Endocrinol Metab*. 2016;311(3):594. doi: 10.1152/ajpendo.00257.2016.
 223. Wall BT, Dirks ML, Snijders T, et al. Short-term muscle disuse lowers myofibrillar protein synthesis rates and induces anabolic resistance to protein ingestion. *Am J Physiol Endocrinol Metab*. 2016;310(2):137. doi: 10.1152/ajpendo.00227.2015.
 224. Fox DK, Ebert SM, Bongers KS, et al. p53 and ATF4 mediate distinct and additive pathways to skeletal muscle atrophy during limb immobilization. *Am J Physiol Endocrinol Metab*. 2014;307(3):245-261. doi: 10.1152/ajpendo.00010.2014.
 225. Fry CS, Johnson DL, Ireland ML, Noehren B. ACL injury reduces satellite cell abundance and promotes fibrogenic cell expansion within skeletal muscle. *J Orthop Res*. 2017;35(9):1876-1885. doi: 10.1002/jor.23502.
 226. Williams GN, Snyder-Mackler L, Barrance PJ, Axe MJ, Buchanan TS. Neuromuscular function after anterior cruciate ligament reconstruction with autologous semitendinosus-gracilis graft. *J Electromyogr Kinesiol*. 2005;15(2):170-180. doi: 10.1016/j.jelekin.2004.08.005.
 227. Drechsler WI, Cramp MC, Scott OM. Changes in muscle strength and EMG median frequency after anterior cruciate ligament reconstruction. *Eur J Appl Physiol*. 2006;98(6):613-623. doi: 10.1007/s00421-006-0311-9.
 228. Palmieri-Smith RM, Thomas AC, Wojtys EM. Maximizing quadriceps strength after ACL reconstruction. *Clin Sports Med*. 2008;27(3):405-ix. doi: 10.1016/j.csm.2008.02.001.
 229. Grapar Zargi T, Drobnic M, Jkoder J, Strazar K, Kacin A. The effects of preconditioning with ischemic exercise

- on quadriceps femoris muscle atrophy following anterior cruciate ligament reconstruction: a quasi-randomized controlled trial. *Eur J Phys Rehabil Med.* 2016;52(3):310-320.
230. Yasuda K, Ohkoshi Y, Tanabe Y, Kaneda K. Muscle weakness after anterior cruciate ligament reconstruction using patellar and quadriceps tendons. *Bull Hosp Jt Dis Orthop Inst.* 1991;51(2):175-185.
 231. de Jong SN, van Caspel DR, van Haeff MJ, Saris DB. Functional assessment and muscle strength before and after reconstruction of chronic anterior cruciate ligament lesions. *Arthroscopy.* 2007;23(1):21-3. doi: 10.1016/j.arthro.2006.08.024.
 232. Konishi Y, Ikeda K, Nishino A, Sunaga M, Aihara Y, Fukubayashi T. Relationship between quadriceps femoris muscle volume and muscle torque after anterior cruciate ligament repair. *Scand J Med Sci Sports.* 2007;17(6):656-661. doi: 10.1111/j.1600-0838.2006.00619.x.
 233. Hiemstra LA, Webber S, MacDonald PB, Kriellaars DJ. Knee strength deficits after hamstring tendon and patellar tendon anterior cruciate ligament reconstruction. *Med Sci Sports Exerc.* 2000;32(8):1472-1479. doi: 10.1097/00005768-200008000-00016.
 234. Konishi Y, Fukubayashi T. Relationship between muscle volume and muscle torque of the hamstrings after anterior cruciate ligament reconstruction. *J Sci Med Sport.* 2010;13(1):101-105. doi: 10.1016/j.jsams.2008.08.001.
 235. Kobayashi A, Higuchi H, Terauchi M, Kobayashi F, Kimura M, Takagishi K. Muscle performance after anterior cruciate ligament reconstruction. *Int Orthop.* 2004;28(1):48-51. doi: 10.1007/s00264-003-0502-5.
 236. Kilinç BE, Kara A, Camur S, Oc Y, Celik H. Isokinetic dynamometer evaluation of the effects of early thigh diameter difference on thigh muscle strength in patients undergoing anterior cruciate ligament reconstruction with hamstring tendon graft. *J Exerc Rehabil.* 2015;11(2):95-100. doi: 10.12965/jer.150100.
 237. Keays SL, Bullock-Saxton JE, Keays AC, Newcombe PA, Bullock MI. A 6-year follow-up of the effect of graft site on strength, stability, range of motion, function, and joint degeneration after anterior cruciate ligament reconstruction: patellar tendon versus semitendinosus and gracilis tendon graft. *Am J Sports Med.* 2007;35(5):729-739. doi: 10.1177/0363546506298277 .
 238. Barbalho M, Rocha AC, Seus TL, Raiol R, Del Vecchio FB, Coswig VS. Addition of blood flow restriction to passive mobilization reduces the rate of muscle wasting in elderly patients in the intensive care unit: a within-patient randomized trial. *Clin Rehabil.* 2019;33(2):233-240. doi: 10.1177/0269215518801440.
 239. Iversen E, Rostad V, Larmo A. Intermittent blood flow restriction does not reduce atrophy following anterior cruciate ligament reconstruction. *J Sport Health Sci.* 2016;5(1):115-118. doi: 10.1016/j.jshs.2014.12.005.
 240. Jeffries O, Waldron M, Pattison JR, Patterson SD. Enhanced local skeletal muscle oxidative capacity and microvascular blood flow following 7-day ischemic preconditioning in healthy humans. *Front Physiol.* 2018;9:463. doi: 10.3389/fphys.2018.00463.
 241. Jones H, Hopkins N, Bailey TG, Green DJ, Cable NT, Thijssen DH. Seven-day remote ischemic preconditioning improves local and systemic endothelial function and microcirculation in healthy humans. *Am J Hypertens.* 2014;27(7):918-925. doi: 10.1093/ajh/hpu004.
 242. Kimura M, Ueda K, Goto C, et al. Repetition of ischemic preconditioning augments endothelium-dependent vasodilation in humans: role of endothelium-derived nitric oxide and endothelial progenitor cells. *Arterioscler Thromb Vasc Biol.* 2007;27(6):1403-1410. doi: 10.1161/ATVBAHA.107.143578.
 243. Jones H, Nyakayiru J, Bailey TG, et al. Impact of eight weeks of repeated ischaemic preconditioning on brachial artery and cutaneous microcirculatory function in healthy males. *Eur J Prev Cardiol.* 2015;22(8):1083-1087. doi: 10.1177/2047487314547657.
 244. Berneis K, Ninnis R, Häussinger D, Keller U. Effects of hyper- and hyposmolality on whole body protein and glucose kinetics in humans. *Am J Physiol.* 1999;276(1):188. doi: 10.1152/ajpendo.1999.276.1.E188.
 245. Nakajima T, Yasuda T, Koide S, et al. Repetitive restriction of muscle blood flow enhances mTOR signaling pathways in a rat model. *Heart Vessels.* 2016;31(10):1685-1695. doi: 10.1007/s00380-016-0801-6.
 246. Hughes L, Jeffries O, Waldron M, et al. Influence and reliability of lower-limb arterial occlusion pressure at different body positions. *Peer J.* 2018;6:e4697. doi: 10.7717/peerj.4697.
 247. Dirks ML, Wall BT, Snijders T, Ottenbros CL, Verdijk LB, van Loon LJ. Neuromuscular electrical stimulation prevents muscle disuse atrophy during leg immobilization in humans. *Acta Physiol (Oxf).* 2014;210(3):628-641. doi: 10.1111/apha.12200.
 248. Natsume T, Ozaki H, Saito AI, Abe T, Naito H. Effects of electrostimulation with blood flow restriction on muscle size and strength. *Med Sci Sports Exerc.* 2015;47(12):2621-2627. doi: 10.1249/MSS.0000000000000722.
 249. Slysz JT, Burr JF. The effects of blood flow restricted electrostimulation on strength and hypertrophy. *J Sport Rehabil.* 2018;27(3):257-262. doi: 10.1123/jsr.2017-0002.
 250. Natsume T, Ozaki H, Kakigi R, Kobayashi H, Naito H. Effects of training intensity in electromyostimulation on human skeletal muscle. *Eur J Appl Physiol.* 2018;118(7):1339-1347. doi: 10.1007/s00421-018-3866-3.

251. Gorgey AS, Timmons MK, Dolbow DR, et al. Electrical stimulation and blood flow restriction increase wrist extensor cross-sectional area and flow mediated dilatation following spinal cord injury. *Eur J Appl Physiol*. 2016;116(6):1231-1244. doi: 10.1007/s00421-016-3385-z.
252. Koltyn KF, Brellenthin AG, Cook DB, Sehgal N, Hillard C. Mechanisms of exercise-induced hypoalgesia. *J Pain*. 2014;15(12):1294-1304. doi:10.1016/j.jpain.2014.09.006.
253. Koltyn KF. Exercise-induced hypoalgesia and intensity of exercise. *Sport Med*. 2002;32(8):477-487. doi: 10.2165/00007256-200232080-00001.
254. Cancio J, Sgromolo N, Rhee P. Blood flow restriction therapy after closed treatment of distal radius fractures. *J Wrist Surg*. 2019;8(4):288-294. doi:10.1055/s-0039-1685455.
255. Korakakis V, Whiteley R, Epameinontidis K. Blood flow restriction induces hypoalgesia in recreationally active adult male anterior knee pain patients allowing therapeutic exercise loading. *Phys Ther Sport*. 2018;32:235-243. doi:10.1016/j.ptsp.2018.05.021.
256. Korakakis V, Whiteley R, Giakas G. Low load resistance training with blood flow restriction decreases anterior knee pain more than resistance training alone. A pilot randomised controlled trial. *Phys Ther Sport*. 2018;34:121-128. doi:10.1016/j.ptsp.2018.09.007.
257. Bryk FF, dos Reis AC, Fingerhut D, et al. Exercises with partial vascular occlusion in patients with knee osteoarthritis: a randomized clinical trial. *Knee Surg Sport Traumatol Arthrosc*. 2016;24(5):1580-1586. doi:10.1007/s00167-016-4064-7.
258. Hughes L, Paton B, Haddad F, Rosenblatt B, Gissane C, Patterson SD. Comparison of the acute perceptual and blood pressure response to heavy load and light load blood flow restriction resistance exercise in anterior cruciate ligament reconstruction patients and non-injured populations. *Phys Ther Sport*. 2018;33:54-61. doi: 10.1016/j.ptsp.2018.07.002.
259. Hughes L, Patterson SD. Low intensity blood flow restriction exercise: rationale for a hypoalgesia effect. *Med Hypotheses*. 2019;132:109370. doi:10.1016/j.mehy.2019.109370.
260. Stagg NJ, Mata HP, Ibrahim MM, et al. Regular exercise reverses sensory hypersensitivity in a rat neuropathic pain model. *Anesthesiology*. 2011;114(4):940-948. doi:10.1097/ALN.0b013e318210f880.
261. Thoren P, Floras JS, Hoffman P, Seals DR. Endorphins and exercise: physiological mechanisms and clinical implications. *Med Sci Sport Exerc*. 1990;22(4):417-428. doi:10.1249/00005768-199008000-00001.
262. Kjaer M, Secher NH, Bach FW, Galbo H, Reeves DR, Mitchell JH. Hormonal, metabolic, and cardiovascular exercise in humans: influence of epidural responses to static anesthesia. *Am Physiol Soc*. 1991;261:E214-20. doi:10.1152/ajpendo.1991.261.2.E214.
263. Koltyn KF. Analgesia following exercise: a review. *Sport Med*. 2000;29(2):85-98. doi:10.2165/00007256-200029020-00002.
264. Dietrich A, McDaniel WF. Endocannabinoids and exercise. *Br J Sports Med*. 2004;38(5):536-541. doi:10.1136/bjism.2004.011718.
265. Hoffmann P, Skarphedinsson JO, Delle M, Thoren P. Electrical stimulation of the gastrocnemius muscle in the spontaneously hypertensive rat increases the pain threshold: Role of different serotonergic receptors. *Acta Physiol Scand*. 1990;138(2):125-131. doi:10.1111/j.1748-1716.1990.tb08824.x.
266. Mogil JS, Belknap JK. Sex and genotype determine the selective activation of neurochemically-distinct mechanisms of swim stress-induced analgesia. *Pharmacol Biochem Behav*. 1996;56(1):61-66. doi:10.1016/S0091-3057(96)00157-8.
267. Neto GR, Novaes JS, Dias I, Brown A, Vianna J, Cirilo-Sousa MS. Effects of resistance training with blood flow restriction on haemodynamics: a systematic review. *Clin Physiol Funct Imaging*. 2017;37(6):567-574. doi: 10.1111/cpf.12368.
268. Ellingson LD, Koltyn KF, Kim J, Cook DB. Does exercise induce hypoalgesia through conditioned pain modulation? *Psychophysiology*. 2014;51(3):267-276. doi:10.1111/psyp.12168.
269. Chitour D, Dickenson A, Le Bars D. Pharmacological evidence for the involvement of serotonergic mechanisms in diffuse noxious inhibitory controls (DNIC). *Brain Res*. 1982;236(2):329-337. doi: 10.1016/0006-8993(82)90718-1.
270. Lemley KJ, Hunter SK, Bement MKH. Conditioned pain modulation predicts exercise-induced hypoalgesia in healthy adults. *Med Sci Sport Exerc*. 2014;47(1):176-184. doi:10.1249/MSS.0000000000000381.
271. Le Bars D, Willer J. Diffuse noxious inhibitory controls (DNIC). In: Basbaum AI, Bushnell MC, eds. *Science of Pain*. Oxford, United Kingdom: Elsevier; 2009:763-773.
272. Vaegter HB, Handberg G, Graven-Nielsen T. Similarities between exercise-induced hypoalgesia and conditioned pain modulation in humans. *Pain*. 2014;155(1):158-167. doi:10.1016/j.pain.2013.09.023.
273. Willer J., De Broucker T, Le Bars D. Encoding of nociceptive thermal stimuli by diffuse noxious inhibitory controls in humans. *J Neurophysiol*. 1989;62:1028-1038. doi: 10.1152/jn.1989.62.5.1028.
274. Oono Y, Wang K, Svensson P, Arendt-Nielsen L. Conditioned pain modulation evoked by different intensities of mechanical stimuli applied to the craniofacial region in healthy men and women. *J Orofac Pain*. 2011;25(4):364-375.

275. Kosek E, Lundberg L. Segmental and plurisegmental modulation of pressure pain thresholds during static muscle contractions in healthy individuals. *Eur J Pain*. 2003;7:251-258. doi:10.1016/S1090-3801(02)00124-6.
276. Lemley KJ, Drewek B, Hunter SK, Bement MKH. Pain relief after isometric exercise is not task-dependent in older men and women. *Med Sci Sport Exerc*. 2013;46(1):185-191. doi:10.1249/MSS.0b013e3182a05de8.
277. Hoeger-Bement MK, Dicapò J, Rasiarmos R, Hunter SK. Dose response of isometric contractions on pain perception in healthy adults. *Med Sci Sport Exerc*. 2008;40(11):1880-1889. doi:10.1249/MSS.0b013e31817eeec.
278. Jessee MB, Mattocks KT, Buckner SL, et al. The acute muscular response to blood flow-restricted exercise with very low relative pressure. *Clin Physiol Funct Imaging*. 2018;38(2):304-311. doi:10.1111/cpf.12416.
279. Hoffman MD, Shepanski MA, Ruble SB, Valic Z, Buckwalter JB, Clifford PS. Intensity and duration threshold for aerobic exercise-induced analgesia to pressure pain. *Arch Phys Med Rehabil*. 2003;85(7):1183-1187. doi:10.1016/j.apmr.2003.09.010.
280. Koltyn KF, Arbogast RW. Perception of pain after resistance exercise. *Br J Sports Med*. 1998;32:20-24. doi:10.1136/bjism.32.1.20.
281. Henneman E, Somjen G, Carpenter DO. Excitability and inhibibility of motoneurons of different sizes. *J Neurophysiol*. 1965;28(3):599-620. doi:10.1152/jn.1965.28.3.599.
282. Pearson SJ, Hussain SR. A review on the mechanisms of blood-flow restriction resistance training-induced muscle hypertrophy. *Sport Med*. 2015;45(2):187-200. doi:10.1007/s40279-014-0264-9.
283. Moritani T, Sherman WM, Shibata M, Matsumoto T, Shinohara M. Oxygen availability and motor unit activity in humans. *Eur J Appl Physiol Occup Physiol*. 1992;64(6):552-556. doi:10.1007/BF00843767.
284. Yasuda T, Fujita T, Miyagi Y, et al. Electromyographic responses of arm and chest muscle during bench press exercise with and without KAATSU. *Int J KAATSU Train Res*. 2006;2(1):15-18. doi:10.3806/ijktr.2.15.
285. Choosing Wisely: An Initiative of the ABIM Foundation. <http://www.choosingwisely.org/societies/american-physical-therapy-association/>. Accessed December 3, 2019.
286. Hylden C, Burns T, Stinner D, Owens J. Blood flow restriction rehabilitation for extremity weakness: a case series. *J Spec Oper Med*. 2015;15(1):50-56.
287. Gualano B, Neves M Jr, Lima FR, et al. Resistance training with vascular occlusion in inclusion body myositis: a case study. *Med Sci Sports Exerc*. 2010;42(2):250-254. doi:10.1249/MSS.0b013e3181b18fb8.
288. Gualano B, Ugrinowitsch C, Neves M Jr, et al. Vascular occlusion training for inclusion body myositis: a novel therapeutic approach. *J Vis Exp*. 2010;(40). doi:10.3791/1894.
289. Jørgensen AN, Aagaard P, Nielsen JL, Frandsen U, Diederichsen LP. Effects of blood-flow-restricted resistance training on muscle function in a 74-year-old male with sporadic inclusion body myositis: a case report. *Clin Physiol Funct Imaging*. 2016;36(6):504-509. doi:10.1111/cpf.12259.
290. Santos AR, Neves MT Jr, Gualano B, et al. Blood flow restricted resistance training attenuates myostatin gene expression in a patient with inclusion body myositis. *Biol Sport*. 2014;31(2):121-124. doi:10.5604/20831862.1097479.
291. Mattocks KT, Grant Mouser J, Jessee MB, et al. Acute hemodynamic changes following high load and very low load lower body resistance exercise with and without the restriction of blood flow. *Physiol Meas*. 2018;39(12):125007. doi:10.1088/1361-6579/aaefc9.
292. da Cunha Nascimento D, Schoenfeld BJ, Prestes J. Potential implications of blood flow restriction exercise on vascular health: a brief review. *Sports Med*. 2020;50(1):73-81. doi:10.1007/s40279-019-01196-5.
293. Jessee MB, Buckner SL, Dankel SJ, Counts BR, Abe T, Loenneke JP. The influence of cuff width, sex, and race on arterial occlusion: implications for blood flow restriction research. *Sports Med*. 2016;46(6):913-921. doi:10.1007/s40279-016-0473-5.
294. Noordin S, McEwen JA, Kragh JF Jr, Eisen A, Masri BA. Surgical tourniquets in orthopaedics. *J Bone Joint Surg Am*. 2009;91(12):2958-2967. doi:10.2106/JBJS.I.00634.
295. AORN Recommended Practices Committee. Recommended practices for the use of the pneumatic tourniquet in the perioperative practice setting. *AORN J*. 2007;86(4):640-655. doi:10.1016/j.aorn.2007.09.004.
296. Mouser JG, Mattocks KT, Buckner SL, et al. High-pressure blood flow restriction with very low load resistance training results in peripheral vascular adaptations similar to heavy resistance training. *Physiol Meas*. 2019;40(3):035003. doi:10.1088/1361-6579/ab0d2a.
297. Segal N, Davis MD, Mikesky AE. Efficacy of blood flow-restricted low-load resistance training for quadriceps strengthening in men at risk of symptomatic knee osteoarthritis. *Geriatr Orthop Surg Rehabil*. 2015;6(3):160-167. doi:10.1177/2151458515583088.
298. Segal NA, Williams GN, Davis MC, Wallace RB, Mikesky AE. Efficacy of blood flow-restricted, low-load resistance training in women with risk factors for symptomatic knee osteoarthritis. *PM R*. 2015;7(4):376-384. doi:10.1016/j.pmrj.2014.09.014.
299. Zeng Z, Centner C, Gollhofer A, König D. Blood-flow-restriction training: validity of pulse ox-

- imetry to assess arterial occlusion pressure. *Int J Sports Physiol Perform.* 2019;14(9):1408-1414. doi: 10.1123/ijspp.2019-0043.
300. Mattocks KT, Jessee MB, Grant Mouser J, et al. The application of blood flow restriction: lessons from the laboratory. *Curr Sports Med Rep.* 2019;17(4):129-134. doi: 10.1249/JSR.0000000000000473.
301. Weatherholt A, VanWye WR, Lohmann J, Owens J. The effect of cuff width for determining limb occlusion pressure: a comparison of blood flow restriction devices. *Int J Exerc Sci.* 2019;12(3):136-143.
302. Loenneke JP, Fahs CA, Rossow LM, et al. Blood flow restriction pressure recommendations: a tale of two cuffs. *Front Physiol.* 2013;4:249. doi: 10.3389/fphys.2013.00249.
303. Buckner SL, Dankel SJ, Counts BR, et al. Influence of cuff material on blood flow restriction stimulus in the upper body. *J Physiol Sci.* 2017;67(1):207-215. doi: 10.1007/s12576-016-0457-0.
304. Hughes L, Rosenblatt B, Gissane C, Paton B, Patterson SD. Interface pressure, perceptual and mean arterial pressure responses to different blood flow restriction systems. *Scand J Med Sci Sports.* 2018;28(7):1757-1765. doi:10.1111/sms.13092.
305. Joshi S, Mahoney S, Pitts L, McCullough S, Hackney K, Jarajapu YPR. Blood flow restriction exercise increases the ACE2/ACE ratio and ACE2 shedding in CD34+ cells in healthy individuals. *FASEB J.* 2019;33(1 Suppl):695.6-695.6.
306. Montgomery R, Paterson A, Williamson C, Florida-James G, Ross MD. Blood flow restriction exercise attenuates the exercise-induced endothelial progenitor cell response in healthy, young men. *Front Physiol.* 2019;10:447. doi: 10.3389/fphys.2019.00447.
307. Mitchell CJ, Churchward-Venne TA, West DWD, et al. Resistance exercise load does not determine training-mediated hypertrophic gains in young men. *J Appl Physiol.* 2012;113(1):71-77. doi: 10.1152/jappphysiol.00307.2012.
308. Shad BJ, Thompson JL, Holwerda AM, et al. One week of step reduction lowers myofibrillar protein synthesis rates in young men. *Med Sci Sports Exerc.* 2019;51(10):2125-2134. doi: 10.1249/MSS.0000000000002034
309. Kilroe S, Fulford J, Jackman S, van Loon L, Wall B. Temporal muscle-specific disuse atrophy during one week of leg immobilization. *Med Sci Sports Exerc.* 2020;52(4):944-954. doi:10.1249/MSS.0000000000002200.
310. Schoenfeld BJ, Ogborn D, Krieger JW. Dose-response relationship between weekly resistance training volume and increases in muscle mass: a systematic review and meta-analysis. *J Sports Sci.* 2017;35(11):1073-1082. doi: 10.1080/02640414.2016.
311. Bagheri R, Rashidlamir A, Motevalli MS, Elliott BT, Mehrabani J, Wong A. Effects of upper-body, lower-body, or combined resistance training on the ratio of follistatin and myostatin in middle-aged men. *Eur J Appl Physiol.* 2019; 119(9):1921-1931. doi:10.1007/s00421-019-04180-z.