

**CURRENT CONCEPTS IN OCCUPATIONAL HEALTH:
MANAGING AN ACUTE INJURY THAT LIMITS WORK PARTICIPATION**

*Trisha Perry, PT, DPT; Anthony Cheung, PT, DPT;
Adrienne Asumbrado, PT, DPT; Katie McBee, PT, DPT*

This 2019 document is a revision of the original document that was rescinded as APTA Guidelines in May 2011 and adopted by the Orthopaedic Section BWC July 11, 2011. It has been published with the following citation:

Perry T, Cheung A, Asumbrado A, McBee K. Current concepts in occupational health: managing an acute injury that limits work participation. *Orthop Phys Ther Pract.* 2019;31(2):101-105.

INTRODUCTION

The purpose of this document is to address physical therapist management for a worker who presents with a neuromusculoskeletal injury incurred on the job, resulting in impairments, activity limitations, and participation restrictions in normal work duties. Related physical therapist interventions and modifications of work methods to prevent workplace recurrence of injury are also addressed. The concepts described for managing acute injuries that result from home or leisure activities would be similar to work-related injuries when work participation is limited, except that non-occupational injuries that are unrelated to employment would not be subject to Occupational Safety and Health Administration (OSHA) requirements.¹

The physical therapist has unique qualifications to facilitate optimal functional outcomes through diagnosis of neuromusculoskeletal conditions and application of interventions to specific body functions and structures affected by the injury. Early physical therapy intervention during the management of the acutely injured worker reduces subsequent use of health care services and downstream costs of care.²⁻⁴ Effective and timely management of the acutely injured worker is enhanced by participation in some form of productive duty, access to workplace-based return-to-work interventions and proactive company approaches, and convenient health care provider services and/or programs.⁵⁻⁷

Inherent in the management of an acute injury that results in work restrictions is frequent and open communication and coordination between the physical therapist and injured worker, other members of the employee health team, and employer representatives as indicated. Clear, concise, functionally relevant information about the injured worker's physical therapist management and recovery progress must be documented and conveyed in a timely manner to necessary stakeholders. Stakeholders may include the injured worker, employer representatives from human resources, safety management, the worker's supervisor and/or a department contact person, occupational health nurse, case manager, adjuster, a physician and/or surgeon, other care providers and the physical therapist.

The following describes a model for managing an acutely injured worker. Concepts discussed are intended to be used in conjunction with the most current versions of the American Physical Therapy Association's Standards of Practice for Physical Therapy,⁸ the Academy of Orthopaedic Physical Therapy's Current Concepts of Orthopaedic Physical Therapy,⁹ Clinical Practice Guidelines,¹⁰ International Classification of Functioning, Disability and Health,¹¹ and nationally recognized occupational health treatment guidelines.

CURRENT CONCEPTS FOR MANAGING AN ACUTE INJURY

1. Reduce local inflammatory response for neuromusculoskeletal pain

Injured worker management during the acute phase is focused on the control and reduction of localized inflammatory response, joint and soft tissue swelling or restriction, and the stabilization and containment of the injury or illness. Immediate, post-trauma intervention lowers the risk of subsequent medical service usage.¹²

Early intervention prevents the negative effects of physical inactivity, disability, depression, and reduces longer-term opioid use and lower-intensity opioid use for musculoskeletal pain.^{3,4,12-15} The role of the physical therapist includes examination and evaluation of an individual for work-related risk factors, impairments, activity limitations, participation restrictions, or other health-related conditions that prevent workers from performing their work duties. An emphasis on instructing the worker in self-management techniques to alleviate symptoms should also be introduced.

2. Validate physical job demands or accommodation options to reduce lost-time

The physical therapist needs to have knowledge of the worker's physical demands for critical work tasks and modified duty options obtained through a job site analysis, video analysis, written physical job demands analysis, or through communication with the employer and/or worker. Early contact between the health care provider and employer to validate physical job demands or accommodation options is an effective workplace intervention to reduce lost-time.⁶

3. Screen for red flags and refer for follow-up when worker is not appropriate for therapy

During the examination and evaluation process, the physical therapist should screen for any red flags and assess the appropriateness for participation in physical therapy. The physical therapist has been equipped with the knowledge and evaluation skills to make appropriate medical referral if the worker is not able to safely participate in physical therapy.¹⁶⁻¹⁸ Communication during this phase typically consists of the worker's status during the initial evaluation noting any impairments, activity limitations, participation restrictions, and whether physical therapy is recommended, or further referral is required.

4. Progress therapeutic management to emphasize daily functional work tasks

Once a diagnosis, prognosis, and plan of care is established, intervention is geared towards improving the worker's ability to move, reducing pain, restoring function, and preventing disability. Therapeutic exercise and functional activity training are the cornerstones of

physical therapist management of the injured worker. The emphasis should be on progressive work and therapeutic activities to increase muscle performance, improve joint integrity and mobility, and improve function for the injured worker. Functional training should include instruction in pacing and body mechanics to improve tolerance for work tasks. Activities and treatment interventions should transition to more vigorous therapeutic activities to prepare the worker for return to usual work and lifestyle activities.

5. Implement evidenced-based interventions into clinical practice

Whenever possible, interventions should be based on evidence supporting its use in order to return injured workers to their jobs safely and in a timely manner. Clinicians are expected to integrate clinical experience with conscientious, explicit, and judicious use of research evidence in order to make clearly informed decisions to help maximize and optimize patient well-being.¹⁹ Clinical practice guidelines and current concepts of orthopaedic physical therapy have been developed by the AOPT based on best available evidence; which provide a good starting point for evaluation, examination, and treatment for commonly encountered clinical scenarios.^{9,10}

6. Identify and address modifiable psychosocial risk factors that may prolong recovery

Since psychosocial risk factors are predictive of future disability with work-related injuries, screening for psychosocial risk factors and integrating behavioral and cognitive modification techniques targeted to address modifiable psychosocial risk factors can reduce future disability.²⁰⁻²² Assessment tools used to help screen for any psychosocial risk factors include the Fear-Avoidance Beliefs Questionnaire (FABQ),²³⁻²⁵ the Fear-Avoidance Components Scale (FACS), the STarT Back,²⁶ the Örebro Musculoskeletal Pain Screening Questionnaire,²⁷ and the Optimal Screening for Prediction of Referral and Outcome (OSPRO).²⁸ Early identification of workers at risk of developing chronic conditions and associated work-related disability is important for appropriate modification of a worker's plan of care and education.²⁴ Psychosocial issues may be as important as physical management in preventing chronicity and understanding disability.²⁹ Interventions in the acute stage addressing these issues may be most useful in reducing fear-avoidance beliefs and promoting return to normal activity.^{20-22,29} Patient education based on a fear avoidance model consists of educating the injured worker in a way such that the worker views his or her pain as a common condition, rather than as a serious disease that needs careful protection.³⁰ Education in combination with exercise have decreased fear-avoidance beliefs and reduced long-term absences due to illness in individuals with low back pain.³¹

7. Promote modified duty with work restrictions to reduce lost productivity

Modern clinical management for most neuromusculoskeletal conditions supports having the injured worker stay at work, with modifications if needed, or return-to-work as soon as the injured worker is medically appropriate.³²⁻³⁷ Workers with neuromusculoskeletal conditions who return-to-work enjoy better health than those who remain off of work.^{33,37,38} Participating in work tasks is noted to be: (1) be therapeutic, (2) help to promote recovery and rehabilitation, (3) lead to better health outcomes, (4) minimize the harmful physical, mental and social effects of long-term sickness absence and worklessness, (5) reduce the chances of chronic disability, long-term

incapacity for work and social exclusion, (6) promote full participation in society, independence and human rights, (7) reduce poverty, and (8) improve quality of life and well-being.³⁸

If the worker is unable to safely participate in normal work duties, reduction of the physical demands with transitional work or temporary modified duty with work restrictions can facilitate early return-to-work and promote work retention.^{32,34,38} Reduced work hours, worksite modifications, and adjustments to job responsibilities are commonly used strategies to provide return-to-work opportunities for workers who can participate in some work duties but have not fully recovered from their injury and are not ready to be fully re-integrated into the workplace.⁵

Physical activity and early return-to-work interventions are not associated with increased risk of recurrent injury if there is compliance with work and activity restrictions.^{32-34,36-42} Modified work programs cut the number of lost work days in half, and injured workers whom are offered modified work duty return-to-work twice as often as those that are not.⁶ Modified work can be introduced in a variety of ways in which each case should be individually assessed and tailored to the injured worker.^{43,44} Transitional work arrangements are a way of facilitating return-to-work and is only meant to be temporary. There is also strong support that a workplace-based return-to-work program can reduce work disability duration and associated costs.^{6,34-36,38} Implementing strategies early in the process with an emphasis on return-to-work, is one of the most effective ways to support positive employment outcomes for workers with impairments.⁷

8. Minimize risks of injury recurrence by facilitating job and work station improvements

During the return-to-work planning, the physical therapist also has a role in minimizing injury recurrence, which may include making sound and practical ergonomic recommendations for work station design, work performance and worker training to improve knowledge of personal responsibilities for fatigue control. Refer to AOPT's Current Concepts in Occupational Health: Work-Related Injury and Illness Prevention for further recommendations along with interventions and parameters related to occupational injury/illness prevention and ergonomic services as provided by physical therapists.⁴⁵

9. Monitor worker response to resumption of normal work tasks and modify as needed

Resumption of normal work tasks for the injured worker is warranted when the functional goals set by the physical therapist have been met or exceeded and the worker has returned to work without any restrictions.⁴² If there is uncertainty on how the injured worker would tolerate a resumption of normal work tasks, the worker may benefit from a trial of full work duty to assess tolerance of normal work activities. The ultimate anticipated goal is the restoration of the injured worker's physical and functional capacity for a safe and expeditious return-to-work. If impairments are still present and causing disability, and the injured worker appears to not be benefiting from physical therapy, the injured worker may then be referred for the need of additional interventions or the appropriateness of an impairment rating.^{16-18,46}

10. Consider referral to another health professional when disability duration exceeds guideline recommendations

If the neuromusculoskeletal problem is not satisfactorily resolved within a limited number of visits per nationally recognized occupational health treatment guidelines, a referral for further examination and evaluation by another health professional may be indicated.^{16,17} Official Disability Guidelines include physical therapy treatment guidelines as a resource that provides an evidence-based starting point for time out of work, serving as an invaluable tool for obtaining the information necessary for effective management of return-to-work following illness/injury and clinical practice recommendations.⁴⁶

An objective Functional Capacity Evaluation (FCE) should be considered when disability duration is excessive and there is not adequate information to substantiate a worker's readiness and ability for a safe return-to-work.^{47,48} An FCE is a comprehensive performance-based medical assessment consisting of a standardized battery of tests in which an injured worker's functional ability is determined and then compared to the physical job demands.^{49,50} An FCE can be used to indicate physical and functional recovery following an injury and guide return-to-work readiness.⁵⁰⁻⁵² Results from an FCE, along with a review of previous treatment progression, provide input into whether a worker can physically participate in work tasks or whether they may require entry into an appropriate work conditioning or work hardening program.

CONCLUSION

The global outcomes of effective physical therapist management of the acutely injured worker are to optimize work performance and minimize the development of work-related occupational disability. Physical therapists are uniquely skilled to manage the rehabilitation of the acutely injured worker and best positioned to assess return-to-work readiness and the timing of such readiness through a thorough evaluation and examination assessing for any impairments, and activity limitations that may hinder involvement in normal work duties. Managing acute injuries in a cost-effective manner relies heavily on collaboration and communication among all involved stakeholders. Early physical therapy intervention and participation in productive work, whether that consists of normal work duties or modified work, is essential in facilitating optimal functional outcomes, promoting quicker return-to-work duties, managing utilization costs; all while reducing the potential detrimental effects to the neuromusculoskeletal system due to physical inactivity. Proper management must also include the identification and intervention of risk factors that may impact positive outcomes or need further medical referral, while gradually advancing the injured worker toward more functional activities and occupation-specific stresses. Through adherence to evidence-based treatment guidelines and timely early intervention, physical therapists play a pivotal role in the management and prevention of recurrent workplace injuries for the acutely injured worker.

REFERENCES

- [1] Occupational Safety and Health Administration. 1904.7 – General recording criteria. | <https://www.osha.gov/laws-regs/regulations/standardnumber/1904/1904.7>. Accessed February 25, 2019.

- [2] Childs J, Fritz JM, Wu SS, et al. Implications of early and guideline adherent physical therapy for low back pain on utilization and costs. *BMC Health Serv Res.* 2015;15:150. doi: 10.1186/s12913-015-0830-3.
- [3] Gatchel RJ, Polatin PB, Noe C, Gardea M, Pulliam C, Thompson J. Treatment and cost effectiveness of early intervention for acute low-back pain patients: a one-year prospective study. *J Occup Rehabil.* 2003;13(1):1-9.
- [4] Whitfill T, Haggard R, Bierner SM, Pransky G, Hassett RG, Gatchel RJ. Early intervention options for acute low back pain patients: a randomized clinical trial with one-year follow-up outcomes. *J Occup Rehabil.* 2010;20(2):256-263. doi: 10.1007/s10926-010-9238-4.
- [5] Ashley J, Cashdollar W, Etcheverry R, Magill K. Transition back to work: policies to support return to work after illness or injury. IMPAQ International. 2017. https://www.dol.gov/odep/topics/pdf/PAP_Transition_Back_to_Work_FINAL_2017-09-07.pdf. Accessed February 25, 2019.
- [6] Cullen KL, Irvin E, Collie A, et al. Effectiveness of workplace interventions in return-to-work for musculoskeletal, pain-related and mental health conditions: an update of the evidence and messages for practitioners. *J Occup Rehabil.* 2018;28(1):1-15. doi: 10.1007/s10926-016-9690-x.
- [7] Waddell G, Burton AK, Kendall N. *Vocational Rehabilitation: What Works, for Whom, and When?* London, UK: The Stationery Office; 2008.
- [8] American Physical Therapy Association. Standards of Practice for Physical Therapy. Updated October 1, 2013. http://www.apta.org/uploadedFiles/APTAorg/About_Us/Policies/Practice/StandardsPractice.pdf. Accessed February 25, 2019.
- [9] Academy of Orthopaedic Physical Therapy. *Current Concepts of Orthopaedic Physical Therapy*, 4th Ed. La Crosse, WI: Academy of Orthopaedic Physical Therapy. 2016.
- [10] American Physical Therapy Association. Clinical Practice Guidelines (CPGs) Developed by APTA. <http://www.apta.org/EvidenceResearch/EBPTools/CPGs/APTA/>. Accessed February 25, 2019.
- [11] World Health Organization. International Classification of Functioning, Disability and Health (ICF). <http://www.who.int/classifications/icf/en/>. Accessed February 25, 2019.
- [12] Gellhorn AC, Chan L, Martin B, Friedly J. Management patterns in acute low back pain: the role of physical therapy. *Spine (Phila Pa 1976).* 2012;37(9):775-782. doi: 10.1097/BRS.0b013e3181d79a09.
- [13] Sun E, Mosfegh J, Rishel CA, Cook CE, Goode AP, George SZ. Association of early physical therapy with long-term opioid use among opioid-naïve patients with musculoskeletal pain. *JAMA Network Open.* 2018;1(8):e185909. doi: 10.1001/jamanetworkopen.2018.5909.
- [14] Horn ME, Fritz JM. Timing of physical therapy consultation on 1-year healthcare utilization and costs in patients seeking care for neck pain: a retrospective cohort. *BMC Health Serv Res.* 2018;18(1):887. doi: 10.1186/s12913-018-3699-0.
- [15] Fritz JM, Magel JS, McFadden M, et al. Early physical therapy vs usual care in patients with recent-onset low back pain: a randomized clinical trial. *JAMA.* 2015;314(14):1459-1467. doi: 10.1001/jama.2015.11648.

- [16] Boissonnault WG, Ross MD. Physical therapists referring patients to physicians: a review of case reports and series. *J Orthop Sports Phys Ther.* 2012;42(5):446-454. doi: 10.2519/jospt.2012.3890. Epub 2012 Jan 25.
- [17] Jette DU, Ardleigh K, Chandler K, McShea L. Decision-making ability of physical therapists: physical therapy intervention or medical referral. *Phys Ther.* 2006;86(12):1619-1629.
- [18] Samsson K, Larsson ME. Physiotherapy screening of patients referred for orthopaedic consultation in primary healthcare – a randomised controlled trial. *Man Ther.* 2014;19(5):386-391. doi: 10.1016/j.math.2013.10.004. Epub 2013 Oct 29.
- [19] Manske RC, Lehecka BJ. Evidence-based medicine/practice in sports physical therapy. *Int J Sports Phys Ther.* 2012;7(5):461-473.
- [20] Hill JC, Fritz JM. Psychosocial influences on low back pain, disability, and response to treatment. *Phys Ther.* 2011;91(5):712-721. doi: 10.2522/ptj.20100280. Epub 2011 Mar 30.
- [21] Pransky G, Gatchel R, Linton SJ, Loisel P. Improving return to work research. *J Occup Rehabil.* 2005;15(4):453-457.
- [22] New Zealand Guidelines Group. Guide to . https://chiro.org/LINKS/GUIDELINES/FULL/NEW_ZEALAND/Guide_to_Assessing/full_text.html. Accessed February 25, 2019.
- [23] Waddell G, Newton M, Henderson I, Somerville D, Main CJ. A Fear-Avoidance Beliefs Questionnaire (FABQ) and the role of fear-avoidance beliefs in chronic low back pain and disability. *Pain.* 1993;52(2):157-168.
- [24] Neblett R, Mayer TG, Hartzell MM, Williams MJ, Gatchel RJ. The Fear-avoidance Components Scale (FACS): development and psychometric evaluation of a new measure of pain-related fear avoidance. *Pain Pract.* 2016;16(4):435-450. doi: 10.1111/papr.12333. Epub 2015 Jul 31.
- [25] Cleland JA, Fritz JM, Brennan GP. Predictive validity of initial fear avoidance beliefs in patients with low back pain receiving physical therapy: is the FABQ a useful screening tool for identifying patients at risk for a poor recovery? *Eur Spine J.* 2008;17(1):70-79.
- [26] Hill JC, Whitehurst DG, Lewis M, et al. Comparison of stratified primary care management for low back pain with current best practice (STarT Back): a randomized controlled trial. *Lancet.* 2011;378(9802):1560-71. doi: 10.1016/S0140-6736(11)60937-9. Epub 2011 Sep 28.
- [27] Westman A, Linton SJ, Öhrvik J, Wahlen P, Leppert J,. Do psychosocial factors predict disability and health at a 3-year follow-up for patients with non-acute musculoskeletal pain? A validation of the Örebro Musculoskeletal Pain Screening Questionnaire. *Eur J Pain.* 2008;12(5):641-649.
- [28] George SZ, Beneciuk JM, Lentz TA, et al. Optimal Screening for Prediction of Referral and Outcome (OSPRO) for musculoskeletal pain conditions: results from the validation cohort. *J Orthop Sports Phys Ther.* 2018;48(6):460-475. doi: 10.2519/jospt.2018.7811. Epub 2018 Apr 7.
- [29] Fritz JM, George SV, Delitto A. The role of fear-avoidance beliefs in acute low back pain: relationships with current and future disability and work status. *Pain.* 2001;94(1):7-15.

- [30] George SV, Bialosky JE, Fritz JM. Physical therapist management of a patient with acute low back pain and elevated fear-avoidance beliefs. *Phys Ther.* 2004;84(6):538-539.
- [31] Louw A, Diener I, Butler DS, Puentedura EJ. The effect of neuroscience education on pain, disability, anxiety, and stress in chronic musculoskeletal pain. *Arch Phys Med Rehabil.* 2011;92(12):2041-2056. doi: 10.1016/j.apmr.2011.07.198.
- [32] Shaw WS, Nelson CC, Woiszwilllo MJ, Gaines B, Peters SE. Early return to work has benefits for relief of back pain and functional recovery after controlling for multiple confounds. *J Occup Environ Med.* 2018;60(10):901-910.
- [33] Stay-at-Work and Return-to-Work Process Improvement Committee. Preventing needless work disability by helping people stay employed. *J Occup Environ Med.* 2006 Sep;48(9):972-87.
- [34] Krause N, Dasinger LK, Neuhauser F. Modified work and return to work: a review of the literature. *J Occup Rehabil.* 1998;8(2):113-139.
- [35] Bernacki EJ, Guidera JA, Schaefer JA, Tsai S. A facilitated early return to work program at a large urban medical center. *J Occup Environ Med.* 2000;42(12):1172-1177.
- [36] Brooker AS, Cole DC, Hogg-Johnson S, Smith J, Frank JW. Modified work: prevalence and characteristics in a sample of workers with soft-tissue injuries. *J Occup Environ Med.* 2001;43(3):276-284.
- [37] Jurisic M, Bean M, Harbaugh J, Cloeren M, Hardy S, Liu H, Nelson C, Christian J. The Personal Physician's Role in Helping Patients With Medical Conditions Stay at Work or Return to Work. *J Occup Environ Med.* 2017 Jun;59(6):e125-e131.
- [38] Waddell G, Burton AK. *Is Work Good for Your Health and Well-Being?* London, UK: The Stationery Office; 2007.
- [39] Hlobil H, Staal JB, Twisk J, et al. The effects of a graded activity intervention for low back pain in occupational health on sick leave, functional status and pain: 12-month results of a randomized controlled trial. *J Occup Rehabil.* 2005;15(4):569-580.
- [40] Bültmann U, Sherson D, Olsen J, Hansen CL, Lund T, Kilsgaard J. Coordinated and tailored work rehabilitation: a randomized controlled trial with economic evaluation undertaken with workers on sick leave due to musculoskeletal disorders. *J Occup Rehabil.* 2009;19(1):81-93. doi: 10.1007/s10926-009-9162-7. Epub 2009 Jan 24.
- [41] Cheng MS, Amick BC 3rd, Watkins MP, Rhea CD. Employer, physical therapist, and employee outcomes in the management of work-related upper extremity disorders. *J Occup Rehabil.* 2002;12(4):257-267.
- [42] Johnston V, Nielsen M, Corbière M, Franche RL. Experiences and perspectives of physical therapists managing patients covered by workers' compensation in Queensland, Australia. *Phys Ther.* 2012;92(10):1306-1315.
- [43] Mathematica Policy Research. Ben-Shalom Y. Steps states can take to help workers keep their jobs after injury, illness, or disability. https://www.dol.gov/odep/topics/pdf/SAW-RTW_PAP_States.pdf. Accessed February 25, 2019.
- [44] United States Department of Labor. S@W/R2W Research & RETAIN Demonstration Projects. Research & Publications. <https://www.dol.gov/odep/topics/SAW-RTW/research-publications.htm>. Accessed February 25, 2019.
- [45] Murphy B, Deal L, Furtak C, Studebaker C, Koehler M [2017]. Current concepts in occupational health: work-related injury/illness prevention and ergonomics guidelines. La Crosse, WI: Academy of Orthopaedic Physical Therapy Academy of Orthopaedic

- Physical Therapy. 2017.
https://www.orthopt.org/uploads/content_files/files/OHSIG%20PREVENTION%20AND%20ERGONOMICS%202017.pdf. Accessed February 25, 2019.
- [46] ODG by MCG. Return to work & medical treatment guidelines.
<https://www.mcg.com/odg/>. Accessed February 25, 2019.
- [47] Academy of Orthopedic Physical Therapy, Occupational Health Special Interest Group. Guideline: occupational health physical therapy: advanced work rehabilitation guidelines.
https://www.orthopt.org/uploads/content_files/OHSIG_Guidelines/OHSIG_guidelines_2/Work_Rehab_Guideline_Final_Draft_4_1_11.pdf. Accessed February 25, 2019.
- [48] Gross DP, Battié MC, Asante AK. Evaluation of a short-form functional capacity evaluation: less may be best. *J Occup Rehabil*. 2007;17(3):422-435.
- [49] American Physical Therapy Association. Glossary of Workers' Compensation Terms.
<http://www.apta.org/Payment/WorkersCompensation/Glossary/>. Accessed February 25, 2019.
- [50] Academy of Orthopedic Physical Therapy, Occupational Health Special Interest Group. Current concepts in functional capacity evaluation: a best practice guideline.
https://www.orthopt.org/uploads/content_files/files/2018%20Current%20Concepts%20in%20OH%20PT-FCE%2006-20-18%20FINAL.pdf. Accessed February 25, 2019.
- [51] Gross DP, Battié MC. Does functional capacity evaluation predict recovery in workers' compensation claimants with upper extremity disorders? *Occup Environ Med*. 2006;63(6):404-410.
- [52] Genovese E, Galper J. *Guide to the Evaluation of Functional Ability*. Chicago, IL: American Medical Association; 2009.